

Visual preferences for direct gaze faces in infant macaques (*Macaca mulatta*) with limited face exposure

Running title: Preference for direct gaze

Elizabeth A. Simpson^{a*}, Annika Paukner^{b,c}, Eric J. Pedersen^d, Pier F. Ferrari^e, Lisa A. Parr^{f,g}

^a Department of Psychology, University of Miami, Coral Gables, Florida, USA

^b *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health, Bethesda, Maryland, USA

^c Department of Psychology, Nottingham Trent University, Nottingham, England

^d Department of Psychology and Neuroscience, University of Colorado Boulder, Boulder, Colorado, USA

^e Institut des Sciences, Cognitives-Marc Jeannerod, Université Claude Bernard, Lyon, France

^f Department of Psychiatry and Behavioral Science, Emory University, Atlanta, Georgia, USA

^g Yerkes National Primate Research Center, Emory University, Atlanta, Georgia, USA

*Corresponding author: simpsons@miami.edu

Word count: 5,517 (body)

Abstract

From birth, human and nonhuman primates attend more to faces with direct gaze compared to averted gaze, and previous studies report that attention to the eyes is linked to the emergence of later social skills. Here we explored whether early experiences influence attraction to eye-contact in infant macaques by examining their attention to face pairs varying in their gaze direction across the first 13 weeks of life. Infants raised by human caretakers had limited conspecific interactions (nursery-reared; $N=16$) and were compared to infants raised in rich social environments (mother-reared; $N=20$). Both groups looked longer to faces and the eyes of direct compared to averted gaze faces. Looking to all faces and eyes also increased with age. Nursery-reared infants did not display age-associated increases in attention to direct-gaze faces specifically, suggesting that, while there may be an initial preference for direct gaze faces from birth, social experiences may support its early development.

Key terms: eye contact, mutual gaze, face, infant development, attention, macaque, vision

Acknowledgements: We thank Stephen J. Suomi and the Laboratory of Comparative Ethology for providing resources and assisting with data collection. This work was supported by the National Institutes of Health [NICHD P01HD064653 to PFF, NIMH R01MH104534 and P50MH000029 to LAP], a National Science Foundation CAREER Award [1653737 to EAS], and the Division of Intramural Research, *Eunice Kennedy Shriver* National Institute of Child Health and Human Development, National Institutes of Health.

Conflicts of Interest: The authors have no conflicts of interest to declare.

Eye-contact is a powerful way of establishing a social connection, signaling readiness for social interaction. In human adults, faces with eye-contact are located more rapidly and processed preferentially compared to those with averted gaze (Conty, Tijus, Hugueville, Coelho, & George, 2006; Crehan & Althoff, 2014; Doi & Shinohara, 2013; Framorando, George, Kerzel, & Burra, 2017; Senju & Johnson, 2009; Yokoyama, Sakai, Noguchi, & Kita, 2014). This preferential attraction to direct-gaze faces is evident early in development, prior to substantial postnatal experience. For example, human newborns look longer to faces with open compared to closed eyes (Batki, Baron-Cohen, Wheelwright, Connellan, & Ahluwalia, 2000) and direct compared to averted gaze (Farroni, Csibra, Simion, & Johnson, 2002), and may even show a rudimentary form of gaze-following (i.e., looking in the same direction as someone else: Farroni, Massaccesi, Pividori, Simion, & Johnson, 2004). While the neural mechanisms remain to be fully understood, eye gaze is processed quickly and automatically, and undoubtedly plays a central role in early communicative development (Hoehl, Reid, Parise, Handl, Palumbo, & Striano, 2009).

Other species also share this preference for direct-gaze faces. Nonhuman primates (NHP), like humans, exhibit faster and longer looking to faces with eye-contact (e.g., adult macaques: Leonard, Blumenthal, Gothard, & Hoffman, 2012). Infant NHP also show these preferences. For example, 2- to 8-month-old chimpanzees look longer to faces with their eyes open compared to closed, and look longer to faces with direct as opposed to averted eye gaze (Hirata, Fuwa, Sugama, Kusunoki, & Fujita, 2010; Myowa-Yamakoshi, Tomonaga, Tanaka, & Matsuzawa, 2003). Similarly, infant macaques show a rapid increase in preference for direct compared to averted gaze faces soon after birth, peaking around 2 months of age (Muschinski, Feczko, Brooks, Collantes, Heitz, & Parr, 2016). Thus, sensitivity to mutual gaze appears to have early roots in both ontogeny and phylogeny.

According to the two-process theory of face processing, infants are equipped with an initial inborn bias to detect and orient towards faces, termed *conspec* (Johnson, Senju, & Tomalski, 2015; Morton & Johnson, 1991). Studies in humans suggest that preferences for certain low-level properties (e.g., high-contrast elements in the locations of the eyes and mouth) likely guide newborns' initial attention to faces through a primitive face-configuration detection system (Farroni, Menon, & Johnson, 2006). Faces with eye-contact are more prototypical—compared to faces with eyes closed or averted—and therefore are the best at capturing infants' attention, likely through a subcortical route operating from birth (Johnson, 2005). Faces with direct gaze have high-contrast eye-regions, which may attract infants' attention, a pattern which is disrupted when faces are inverted, suggesting both gaze perception and the configuration of the face (e.g., top-heavy, symmetrical) drive early preferences (Farroni, Johnson, Menon, Zulian, Faraguna, & Csibra, 2005).

This initial interest in faces is further refined by infants' early experiences, in a second more cortically-driven process termed *conlearn* (Morton & Johnson, 1991). Faces are one of the most common visual stimuli infants encounter, enabling them to rapidly develop expertise (Gliga & Csibra, 2007). Differences in infants' early experiences with faces can have lasting impacts. For example, human infants who have blind parents, and therefore have reduced experience of parental eye-contact and gaze behavior, develop seemingly normal social skills; however, they attend less to eye-gaze cues (Senju et al., 2015), particularly in their blind parents (Ganea, Hudry, Tucker, Charman, Johnson, & Senju, 2018). These findings may indicate that a lack of

experience with direct gaze interactions results in infants paying less attention to the eyes. Further, at 6 to 10 months of age, infants of blind parents do not neurally discriminate—measured with event-related potentials (ERPs)—between direct and averted gaze faces, as do infants of sighted parents (Vernetti, Ganea, Tucker, Charman, Johnson, & Senju, 2018). This finding may indicate that selective brain responses to perceived gaze shifts may depend on parent-infant communicative experiences. Together, these studies suggest that while newborns exhibit a basic preference for faces and especially the eyes in the context of the face, infants' responses to faces and eyes become more sophisticated as they get older, influenced by their early environments. Early eye-detection serves an important communicative function. Indeed, by 3 to 4 months of age, infants attend longer and smile more in social interactions with partners making eye-contact compared to looking away, and, in the presence of someone looking away, try to engage partners in eye-contact (Blass & Camp, 2001; Hains & Muir, 1996; Symons, Hains, & Muir, 1998). The early development of face expertise may in fact be driven by attention to the eyes (Gliga & Csibra, 2007).

To better uncover the influence of experience in shaping early attention, more experimental studies are necessary. Attempts to experimentally manipulate infants' early experiences have largely been done in animals (for a review: Davidson & Clayton, 2016). For example, in the first month of life (between 7-30 days of age), infant monkeys exhibit considerable individual differences in the extent to which they attend to the eye-region of faces: in mother-reared infants, males look longer to the eye region than females (Paukner, Slonecker, Murphy, Wooddell, & Dettmer, 2018), but in nursery-reared infants (who have more limited social experience with conspecifics and are raised by human caretakers), females look more to the eye-region than males (Simpson, Nicolini, Shetler, Suomi, Ferrari, & Paukner, 2016). Maternal rank was positively associated with attention to the eyes, although this was only the case for infants reared by their mothers, suggesting this attentional preference may be transmitted through social interactions, rather than inherited through some biological mechanism (Paukner et al., 2018). In sum, it appears that there are individual differences in attention to faces and the eye region, shaped by infants' early social experiences in just the first month of life. However, it remains unknown whether earlier social experience—in the first weeks of life—likewise influences attention to faces as a function of their gaze direction (i.e., eye-contact or averted-gaze). Further, we currently lack longitudinal studies of eye gaze processing.

In the present study, we were interested in the extent to which infants are initially biased to attend to direct-gaze faces, and how these biases are further refined in the first few months of life as a function of their social experience. We hypothesized that faces with direct gaze are special from birth and take on additional meaning with development. That is, we predicted that with age, infants would exhibit increasing relative interest in faces with eye-contact compared to faces looking away.

We chose to study NHP because, unlike humans, we can carefully control infants' early environment, including social experiences and exposure to faces (e.g., Sugita, 2008). In addition, like humans, macaques are highly gregarious and infants engage in complex face-to-face interactions (Ferrari, Paukner, Ionica, & Suomi, 2009) and show considerable individual differences in social behavior, making them a promising model for the study of disorders such as autism (Feczko, Bliss-Moreau, Walum, Pruett, & Parr, 2016). Macaques and humans share a number of similarities in social attention (Parr, 2011). For example, when viewing faces, both

humans and monkeys spend longer viewing the inner than outer facial features (Dahl, Wallraven, Bülthoff, & Logothetis, 2009), especially the eyes (Gothard, Brooks, & Peterson, 2009). Importantly, infant macaques' looking behavior can be assessed with remote eye tracking (e.g., Paukner, Simpson, Ferrari, Mrozek, & Suomi, 2014). Such studies have revealed that, by 3 weeks of age, macaques can efficiently detect and look longer at faces compared to non-faces (Simpson, Jakobsen, Damon, Suomi, Ferrari, & Paukner, 2017). Further, socially reared macaque infants exhibit early preferences for direct-gaze faces soon after birth (Muschinski et al., 2016) followed by the rapid development of gaze following in the first year of life (Rosati, Arre, Platt, & Santos, 2016), much like human infants. We therefore sought to examine whether macaque infants with fewer species-typical early social experiences (i.e., reared by humans in a neonatal nursery) would exhibit different early patterns of attention to direct and averted gaze faces, compared to infants with more species-typical, rich social experiences, reared by their mothers in social groups.

Methods

The *Eunice Kennedy Shriver* National Institute of Child Health and Human Development Animal Care and Use Committee approved the procedures. We conducted the study in accordance with the Guide for the Care and Use of Laboratory Animals and complied with the Animal Welfare Act.

Subjects

Subjects were 16 healthy, full-term infant rhesus macaques (*Macaca mulatta*), 9 females and 7 males, born in 2015. Infants were healthy and were separated from their mothers on the day of birth (typically by 8am) and reared in a nursery facility by human caretakers for ongoing, unrelated research studies.

In the first months of life, these infants had limited exposure to faces generally, including human and NHP faces. Infants could see other infants housed in adjacent cages, but lacked species-typical exposure to adult conspecifics' faces. Human caretakers were present for 13 hours each day and interacted with infants every 2 hours for feeding and cleaning purposes. Caregivers wore personal protective equipment, including goggles, masks covering the nose and mouth, and hats, so only their eyes were visible (see Paukner, Huntsberry, & Suomi, 2010). While difficult to estimate precisely, these infants had limited face exposure generally (to faces of any species), and especially to conspecifics, compared to infants reared by their mothers in social groups.

Infants were raised identically for the first 5 weeks of life. Once the youngest infant reached 36 days of age, infants were placed into small, same-aged peer groups. Infants were randomly assigned to one of two rearing conditions for unrelated research studies: low-socialization infants ($n = 8$; 4 females), and high-socialization infants ($n = 8$, 5 females). Low-socialization infants continued to be individually housed but assigned to playgroups composed of 3 to 4 peers and put together for 2 hours a day, 5 days a week. High-socialization infants were raised in groups with 3 to 4 peers (for details: Simpson et al., 2016). Therefore, by 3 months of age, infants had extensive experience with same-aged conspecifics, though still lacking species-typical exposure to adult conspecifics.

Infants were tested longitudinally in the 13 weeks after birth. Testing occurred weekly in the first month (days 10, 17, 24, 30), then every-other-week in the second and third months after birth (days 44, 58, 72, 86), for a total of 8 time-points. Infants were individually housed in

incubators (51 cm × 38 cm × 43 cm) for the first two weeks of life and in larger cages (65 cm × 73 cm × 83 cm) thereafter. Both housing arrangements contained an inanimate surrogate mother covered with fleece fabric as well as loose pieces of fleece fabric and various toys. Infants were fed Similac infant formula from birth and additionally Purina Monkey chow (#5054) starting at 2 weeks old. Additional food enrichment (fruit, seeds, nuts) was introduced twice daily when infants were 2 months old. Water was available ad libitum.

We compared these nursery-reared infants to a group of mother-reared infants ($n=20$), all males, healthy, full-term, and tested with eye tracking using the same stimuli (for details, see Muschinski et al., 2016). The notable difference from the nursery-reared infants was that the mother-reared infants had enriched species-typical environments, reared by their mothers in large social groups (~50-100 individuals). Mother-reared infants were tested longitudinally across 14 test sessions, starting after the third day of life and continuing through approximately 5 months of age, so their testing also extended over a longer period than that for the nursery-reared infants.

Materials

Stimuli. At each age, infants viewed two novel pairs of unfamiliar faces, for a total of 16 unique trials per infant (8 time points × 2 trials at each age). These faces were randomly selected for each test from a larger set of 42 female monkey faces. Each infant saw each face only once. The stimuli were the same as those used previously in mother-reared rhesus macaque infants (Muschinski et al., 2016). In each trial, two photos of the same adult female monkey were shown for 10 seconds, one with direct gaze and one with averted gaze, side-by-side, cropped closely around the head, presented on a black background (Figure 1; also, see Figure 2 in Muschinski et al., 2016). The direct-gaze faces were facing forward, with eyes looking straight into the camera. The averted-gaze faces' heads and eyes were angled away from the camera at about 45° to either the left or right. The location of the direct and averted gaze faces were balanced, so they were equally likely to appear on the left and right side of the screen. Faces appeared on a black screen, sized 28 × 51 cm (1280 × 720 pixels).

[Figure 1 here]

Eye tracking. Eye movements were recorded via corneal reflection using a Tobii TX300 eye tracker with a remote 58.4 cm monitor and integrated eye tracking technology and a 60 Hertz sampling rate. We used Tobii Studio software (Tobii Technology, Sweden) to collect and summarize the data. An area of interest (AOI) was drawn around each face, and each eye region (see the bottom panel of Figure 1). The sizes of the AOIs did not vary systematically across direct and averted face conditions, for either the face or eye AOIs, $t(19) = 1.49$, $p = .152$ and $t(19) = 1.16$, $p = .262$, respectively. We used the Tobii Filter to extract fixations, defined as occurring within a window of 37 pixels for at least 50 ms.

Procedures

An experimenter held the infant wrapped in soft fleece fabric at a distance of approximately 60 cm from the screen. Each infant was calibrated prior to each test using a 5-point calibration procedure to Tobii Studio's pre-set locations; individual calibration points judged to be unreliable were repeated until acceptable. Following calibration, each infant viewed two face pairs (4 face images total) each day.

Data Analysis

Overall, we followed the general data analysis strategy used by Muschinski et al. (2016). We calculated two dependent variables for each AOI (face and eyes) separately: (1) the proportion¹ of total looking time (the sum of fixation times) to the AOI out of the total time looking to the screen, and (2) the sum of fixation times (in seconds) to a AOI (see Supporting Information, including Figures S1, S2, and S3, and Tables S1 and S2). We analyzed these data with linear mixed models in SPSS v.23. All models included a fixed effect for stimulus category of gaze direction (direct vs. averted gaze), a random intercept for subject, a random slope for age, and an unstructured covariance matrix was used for the random effects.

Results

Main effects of stimulus category

First, we ran models using this basic structure to test for main effects of stimulus category (gaze direction) on our dependent variables in the nursery-reared infants. Stimulus category was dummy coded 1 for direct gaze, 0 for averted gaze.

In the model predicting proportion of total looking time to the face, the intercept was statistically significant, $b = .26$, $SE = .01$, 95% CI: [.23, .29], $p < .001$, indicating that infants spent approximately a quarter of the time they were looking at the screen looking at the face AOI of averted gaze faces. The coefficient for gaze direction was also significant, $b = .05$, $SE = .02$, 95% CI: [.02, .08], $p = .005$, indicating that infants spent more of their time looking at direct gaze faces than averted gaze faces (Figure 2). Similarly, in the model predicting proportion of total looking time to the eyes, both the intercept ($b = .06$, $SE = .01$, 95% CI: [.05, .08], $p < .001$) and the coefficient for gaze direction ($b = .03$, $SE = .01$, 95% CI: [.01, .05], $p = .006$) were significant, indicating that infants spent more of their time looking at the eyes on direct gaze faces than they did looking at the eyes on averted gaze faces.

[Figure 2 here]

Longitudinal effects of stimulus category

Next, we added a fixed effect for week and a week \times stimulus type (gaze direction) interaction to the models to test for longitudinal changes. We rescaled week by subtracting a constant of 2 so that the intercepts in the following models correspond to week 2, the earliest week of testing. Following Muschinski et al. (2016), we fit three full-factorial models of this type for each analysis: one with a linear growth term, one that added a quadratic growth term, and one that added a cubic growth term. Then, we used chi-square deviance tests to determine which of the nested models best fit the data. In all cases except for one, the model with the linear growth term fit the data best, so those are the models we report here. The one exception was the cubic model predicting percentage of time looking at the face—however, none of the terms in this model were statistically significant, indicating that the model was simply overfit due to the large number of included terms. Therefore, we retained the linear model instead.

Full model results are presented in Tables 1 and 2. In the model predicting the proportion of time looking to the face, there were significant main effects for both week ($b =$

¹ Analyzing proportion outcomes in a linear model can lead to model predictions outside the range of 0-1. To verify that this issue did not affect our results here, we re-ran all of the proportion models with a logit-transformed outcome variable, which circumvents the issue without increasing error rates (Warton & Hui, 2011). None of the effects in the logit-transformed models meaningfully differed from the proportion models, so we report the results of the proportion models here for ease of interpretation and consistency with Muschinski et al. (2016).

.008, $p = .026$) and gaze direction ($b = .08$, $p = .014$). Thus, in addition to a predicted increase in looking proportion over time generally, the model predicts that subjects spent an additional 8% of their time looking at direct gaze faces than at averted gaze faces. The interaction between gaze direction and week was not significant, indicating that the effect of gaze direction did not change over development (Figure 3). In the model predicting the proportion of time looking to the eyes, neither week ($b = .005$, $p = .070$) nor gaze direction ($b = .03$, $p = .114$) were statistically significant, but they trended in the same direction as in the face model.

[Figure 3 here]

Comparison with mother-reared infants

In all of their longitudinal analyses for mother-reared infants, Muschinski et al. (2016) found significant interactions between age and gaze direction in predicting fixation time and proportion of looking time. In some of their models, they also found significant non-linear effects of age (i.e., quadratic and cubic). In contrast, we found no significant interactions or non-linear effects of age for our longitudinal analyses in nursery-reared infants, which indicate that there may be differences in developmental trajectories between mother-reared and nursery-reared infants. We obtained the raw data from Muschinski et al. (2016) for the first 13 weeks of testing their mother-reared infants ($N = 20$), and we combined these data with our more recently collected nursery-reared infant data to directly test for overall effects of mother- vs. nursery-rearing. That is, beyond the apparent differences in growth trajectories between the two data sets, do mother-reared infants and nursery-reared infants differ in their overall preferences for direct and averted gaze faces?

To test this, we re-ran our four main effects models—fixation proportion and fixation time for face and eyes AOIs—on the combined dataset with an added predictor for rearing condition (for fixation time results, see Supporting Information). Each model contained a dummy-coded predictor for gaze direction (1 = direct gaze, 0 = averted gaze), rearing condition (1 = mother-reared; 0 = nursery-reared), and their interaction. Muschinski et al. (2016) began testing in the first week of life, but we rescaled the week variable to match our own (i.e., we subtracted a constant of 2 across both datasets). Thus, the intercept in each model referred to nursery-reared infants looking at averted faces in their second week of life. Full model results are displayed in Table 3 (face AOI) and Table 4 (eyes AOI). Particularly noteworthy is that for fixation proportion to the face AOI (Figure 4), mother-reared infants, compared to nursery-reared infants, showed an increased preference for direct-gaze faces, as indicated by the significant gaze direction \times rearing condition interactions. Re-running the model with the dummy code for rearing condition switched confirmed that, in mother-reared infants, there was a significant preference for direct gaze faces compared to averted gaze faces ($b = .13$, $SE = .02$, 95% CI: [.09, .16], $p < .001$).

[Figure 4 here]

Additionally, the significant simple effects of rearing condition in both of the fixation proportion models (face and eyes AOIs) indicate that mother-reared infants spent a greater portion of their time looking at averted gaze faces than did nursery-reared infants (recall that the intercepts of these models refer to nursery-reared infants looking at averted faces). Taken together with the interaction results mentioned above, this indicates a general tendency for mother-reared infants to spend a greater proportion of their time than nursery-reared infants

looking at faces and eyes *in general*, with mother-reared infants also having an even greater preference for direct gaze faces *in particular*.

Finally, we directly tested for differences in growth trajectories between mother-reared and nursery-reared infants by re-running the models with an added fixed effect for week, as well as its interactions with gaze direction and rearing condition. We note that given our relatively small sample size here, the results of these models should be interpreted with caution. In the proportion of looking time model for the face AOI, there was a small but significant week \times stimulus type \times rearing condition interaction ($p = .005$) indicating that mother-reared infants had a stronger preference for direct gaze faces than did nursery-reared infants that gradually decreased (about 3% per week) over time (which can be seen in the upper-left panel of Figure 3). None of the other three models indicated differences in growth trajectories between datasets (full results in Tables S1 and S2). Hence, our models do not conclusively provide evidence that mother-reared and nursery-reared infants differ in their growth trajectories in preferences for direct gaze faces over the first 13 weeks of life, though these models may be underpowered to detect these differences.

Discussion

The present study explored whether early experiences influence attraction to eye-contact in infant rhesus macaques. Using remote eye tracking, we examined infants' attention patterns to face pairs varying in their gaze direction (direct or averted) longitudinally across the first 13 weeks of life. We found that infants who were raised by human caretakers, and therefore had limited contact with conspecifics, spent longer (and a greater proportion of time) looking to direct gaze compared to averted gaze conspecific faces, and especially the eyes. In addition, regardless of gaze direction, nursery-reared infants' looking to faces, and especially the eyes, increased with age. These findings parallel those reported previously in infant monkeys raised with their mothers in rich social environments with conspecifics (Muschinski et al., 2016). These findings suggest that there may be an initial preference for direct gaze faces, and eyes, from birth that is present even when opportunities for social interactions are limited (Sugita, 2008), although some minimal level of face exposure may still be necessary. These results are consistent with the proposal that there may be an "eye direction detector" mechanism (Baron-Cohen, 1994) that develops even with very limited social experience.

However, we did not find evidence that nursery-reared infants had an increased interest specifically in direct-gaze faces with age, suggesting that, while there may be an initial preference for direct gaze faces from birth, social experiences seem to support this attraction and may be necessary for it to grow stronger across the first weeks of life. Further, nursery-reared infants appeared overall less attentive to the faces and eyes compared to mother-reared infants, suggesting overall decreased social interest in conspecifics among infants with more limited social experiences. Similarly, chimpanzees with impoverished early social environments attended less to videos of conspecifics (and instead attended more to nonsocial videos) compared to chimpanzees reared with more socialization (Kano, Shepherd, Hirata, & Call, 2018). These findings are in line with a functional architecture account of development, which proposes that infants' initial predispositions may be supported through early social experiences with caregivers (Murray, De Pascalis, Bozicevic, Hawkins, Sclafani, & Ferrari, 2016). For face processing, more specifically, our findings support the conspec-conlearn two-process theory (Johnson et al., 2015; Morton & Johnson, 1991). That is, infant monkeys appear to be born with

a preference for direct-gaze faces over averted gaze faces, but early social experiences can boost this preference in the first weeks and months of life. In a limited social context, infants may not have as many opportunities to learn the value of direct-gaze faces, so they do not show growing interest in them with age, as do the infants who had more enriched social experiences.

Indeed, studies in both human and nonhuman primates suggest that mutual gaze between infants and caregivers may promote the development of healthy social skills (Dettmer et al., 2016; Murray et al., 2016; Rayson, Bonaiuto, Ferrari, & Murray, 2017). Human 4-month-olds exhibit more facial mimicry when interacting with social partners engaging in eye-contact compared to those without eye-contact (de Klerk, Hamilton, & Southgate, 2018). Human infants who engage in more mutual gaze with their parents go on to develop more attentional control (Niedźwiecka, Ramotowska, & Tomalski, 2018), suggesting that sensitivity to eye-contact may be an important early marker of healthy social development. Sensitivity to the eyes also helps infants to use others, who may be more experienced, as sources of information, for example, to orient to important things in the environment (e.g., social partners, food, or threats; Emery, Lorincz, Perrett, Oram, & Baker, 1997; Teufel, Gutmann, Pirow, & Fischer, 2010). A rich social environment—particularly one that includes joint attention interactions—provides infants with learning opportunities to understand the links between others' gaze and relevant items in the environment (Corkum & Moore, 1998; Gredebäck, Fikke, & Melinder, 2010).

Atypical face and eye processing has been linked to heightened risk of psychopathy (Bedford, Pickles, Sharp, Wright, & Hill, 2015) and autism spectrum disorders (ASD; Gangi et al., 2018; Shultz, Klin, & Jones, 2018). For example, newborns at risk for ASD are less attentive to faces, particularly direct-gaze faces, compared to low-risk newborns (Di Giorgio et al., 2016). Further, at 2 months of age, infants who later are diagnosed with ASD attend to the eye region of faces similar to non-ASD infants, but from 2 to 6 months, ASD infants show a decline in attention to the eye region compared to low-risk control infants, who show an increase in attention to the eyes (Jones & Klin, 2013). The present study found that infants with limited early social experience still exhibit normative attentional preferences to face and eye regions of direct-gaze faces initially in the first weeks of life, but that, with age, their attention to direct-gaze faces and eyes specifically did not increase to a greater extent than their interest in faces generally, as in infants with more species-typical social environments. That is, unlike what previous research has found for mother-reared infants (Muschinski et al., 2016), here we found that nursery-reared infants did not display a pattern of increasing interest in direct gaze faces with age. It is also worth noting, though, that we did not find statistical support for this rearing difference in growth trajectories, so this finding should be interpreted with caution. While this finding might, at first, seem to signify a dysfunction or unhealthy pattern of attention predictive of later disorder, it may, alternatively, be adaptive for infants to develop phenotypes that best suit their particular environments. Drawing parallels between human disorders and the present findings, therefore, should be done with care, mindful of these limitations. Nonetheless, these findings suggest that the developmental trajectories of different types of social attention may be informative indices of normative social development.

Our findings are also consistent with studies of more general face processing. For example, face-deprived infant macaques—not exposed to faces or face-like stimuli after birth—still exhibit attentional preferences for faces compared to non-faces, suggesting initial face biases appear intact even in face-deprived environments (Sugita, 2008). However, experience

with faces may be necessary for the formation of specialized cortical processing for faces, given that face-deprived infant macaques exhibit impaired neural specialization for faces, lacking face patch regions in the superior temporal sulcus that are selectively responsive to faces (Arcaro, Schade, Vincent, Ponce, & Livingstone, 2017). Thus, while initial attentional predispositions may be present independent of experience, early exposure to faces further refines the specialization of this system.

Attention to others' attentional states is a foundation precursor for later social cognitive skills, including gaze-following (i.e., looking where another individual is looking; Frischen, Bayliss, & Tipper, 2007). While a rudimentary form of gaze following is found in human newborns, it is rather inflexible and limited (e.g., requiring motion of the pupil, as well as mutual gaze prior to the motion; Farroni et al., 2004). In humans and NHP, the capacity to systematically follow the direction of others' attention emerges more consistently in infancy around 6 months of age and continues to improve beyond that, becoming more flexible and accurate well into the second year of life (for a review, see Grossman, 2017).

Furthermore, social experiences influence later gaze-following abilities. In human infants, own-race compared to other-race faces are more reliably followed during gaze following tasks at 7 months of age (Xiao et al., 2018). Similarly, in the present study, adult monkey faces were arguably more familiar to the mother-reared infants than the nursery-reared infants. Future work with human face stimuli, or same-aged peer faces, oriented with eye-contact or averted gaze, could be used to see if nursery infants are more likely to discriminate between these types of stimuli and show stronger preferences for direct gaze faces in this context.

Previous studies have found that infant monkeys with more social experience with humans are better at following the direction of humans' eye gaze (i.e., looking where another individual looks) at 7 months of age (Simpson et al., 2016). Early social experiences appear to support the development of infants' gaze following competence, possibly due to infants' increased exposure to social cues, thereby enhancing infants' interest or skill in social interactions.

A number of questions remain, including: Are there long-term consequences of failing to be exposed to social gaze? Studies in rhesus monkeys suggest that infants' early social skills predict their later social relationships. For example, rhesus monkeys with better face recognition and social interaction skills in infancy develop more prosocial behaviors as adults (Sclafani et al., 2016). Similarly, rhesus monkeys with better neonatal imitation skills in the first week of life develop more dominance and higher juvenile social status (Kaburu, Paukner, Simpson, Suomi, & Ferrari, 2016; Wooddell, Simpson, Murphy, Dettmer, & Paukner, in press). These findings suggest that early social capacities may lay the groundwork for later emerging more complex social competencies (for a recent review in humans, see Shultz et al., 2018).

The present study is not without limitations. Mother-reared infants were all male, limiting our ability to test whether there are sex differences. Previous studies suggest female neonates engage in more eye-contact compared to male neonates, in both humans (Hittelman & Dickes, 1979) and macaques (Simpson et al., 2016), although the developmental trajectories of each sex remain, to our knowledge, unexplored. While we found no sex differences in the nursery-reared infants (alone or combined with mother-reared infants), we lacked power and therefore cannot conclude that such differences do not exist. We also followed nursery-reared infants for only the first 13 weeks of life, but there may be developmental changes that occur beyond that

point. Future research with more balanced sex ratios and over more extended periods of development will therefore be important.

In addition, for our stimuli, we used pictures rather than videos or a live individual, as these enabled us to have more experimental control. However, static images are more limited in their ecological validity and may not produce as strong of effects as interacting with a live partner (e.g., Pönkänen, Alhoniemi, Leppänen, & Hietanen, 2010; Risko, Richardson, & Kingstone, 2016). Furthermore, it is unclear to what extent our findings would extend to other types of stimuli, including male faces and faces of other ages or species. For example, nursery-reared infants, who have more experience with same-aged peer faces, may show superior processing with more familiar types of faces, such as those of conspecific infant faces relative to adult faces (Simpson, Suomi, & Paukner, 2016).

Finally, there were methodological differences in our testing of mother-reared and nursery-reared infants (e.g., different eye-tracking systems) that may have influenced data quality and therefore limit our comparisons. While future research using more comparable equipment across the two groups would be ideal, the present study provides initial evidence of different developmental patterns across these groups.

Nonetheless, the present study suggests that early attentional preferences for direct-gaze faces may emerge independent of early infant experiences, but that social experiences appear to also play an important role in supporting the development of social attention.

References

- Arcaro, M. J., Schade, P. F., Vincent, J. L., Ponce, C. R., & Livingstone, M. S. (2017). Seeing faces is necessary for face-domain formation. *Nature Neuroscience*, *20*(10), 1404.
- Batki, A., Baron-Cohen, S., Wheelwright, S., Connellan, J., & Ahluwalia, J. (2000). Is there an innate gaze module? Evidence from human neonates. *Infant Behavior and Development*, *23*(2), 223-229. doi: 10.1016/S0163-6383(01)00037-6
- Bedford, R., Pickles, A., Sharp, H., Wright, N., & Hill, J. (2015). Reduced face preference in infancy: A developmental precursor to callous-unemotional traits? *Biological Psychiatry*, *78*(2), 144-150. doi: 10.1016/j.biopsych.2014.09.022
- Blass, E. M., & Camp, C. A. (2001). The ontogeny of face recognition: Eye contact and sweet taste induce face preference in 9- and 12-week-old human infants. *Developmental Psychology*, *37*(6), 762-774.
- Conty, L., Tijus, C., Hugueville, L., Coelho, E., & George, N. (2006). Searching for asymmetries in the detection of gaze contact versus averted gaze under different head views: a behavioural study. *Spatial Vision*, *19*(6), 529-545. doi: 10.1163/156856806779194026
- Corkum, V. & Moore, C. (1998). The origins of joint visual attention in infants. *Developmental Psychology*, *34*, 28-38.
- Crehan, E. T., & Althoff, R. R. (2015). Measuring the stare-in-the-crowd effect: a new paradigm to study social perception. *Behavior Research Methods*, *47*(4), 994-1003. doi: 10.3758/s13428-014-0514-7
- Dahl, C. D., Wallraven, C., Bülthoff, H. H., & Logothetis, N. K. (2009). Humans and macaques employ similar face-processing strategies. *Current Biology*, *19*(6), 509-513. doi: 10.1016/j.cub.2009.01.061
- Davidson, G. L., & Clayton, N. S. (2016). New perspectives in gaze sensitivity research. *Learning & Behavior*, *44*(1), 9-17. doi: 10.3758/s13420-015-0204-z
- DeCasper, A. J., & Fifer, W. P. (1980). Of human bonding: Newborns prefer their mothers' voices. *Science*, *208*, 1174-1176.
- de Klerk, C. C., Hamilton, A. F. D. C., & Southgate, V. (2018). Eye contact modulates facial mimicry in 4-month-old infants: an EMG and fNIRS study. *Cortex*, *106*, 93-103.
- Dettmer, A. M., Kaburu, S. S., Simpson, E. A., Paukner, A., Sclafani, V., Byers, K. L., ... & Suomi, S. J. (2016). Neonatal face-to-face interactions promote later social behaviour in infant rhesus monkeys. *Nature Communications*, *7*, 11940. doi: 10.1038/ncomms11940
- Di Giorgio, E., Frasnelli, E., Salva, O. R., Scattoni, M. L., Puopolo, M., Tosoni, D., ... & Persico, A. (2016). Difference in visual social predispositions between newborns at low- and high-risk for autism. *Scientific Reports*, *6*, 26395.
- Doi, H., & Shinohara, K. (2013). Task-irrelevant direct gaze facilitates visual search for deviant facial expression. *Visual Cognition*, *21*(1), 72-98. doi: 10.1080/13506285.2013.779350
- Emery, N. J., Lorincz, E. N., Perrett, D. I., Oram, M. W., & Baker, C. I. (1997). Gaze following and joint attention in rhesus monkeys (*Macaca mulatta*). *Journal of Comparative Psychology*, *111*, 286-293.
- Farroni, T., Csibra, G., Simion, F., & Johnson, M. H. (2002). Eye contact detection in humans from birth. *Proceedings of the National Academy of Sciences*, *99*(14), 9602-9605. doi: 10.1073/pnas.152159999

- Farroni, T., Johnson, M. H., Menon, E., Zulian, L., Faraguna, D., & Csibra, G. (2005) Newborns' preference for face-relevant stimuli: Effects of contrast polarity. *Proceedings of the National Academy of Sciences*, *102*(47), 17245–17250.
- Farroni, T., Massaccesi, S., Pividori, D., Simion, F., & Johnson, M. H. (2004). Gaze following in newborns. *Infancy*, *5*(1), 39-60. doi: 10.1207/s15327078in0501_2
- Farroni, T., Menon, E., & Johnson, M. H. (2006). Factors influencing newborns' preference for faces with eye contact. *Journal of Experimental Child Psychology*, *95*(4), 298-308. doi: 10.1016/j.jecp.2006.08.001
- Feczko, E. J., Bliss-Moreau, E., Walum, H., Pruett Jr, J. R., & Parr, L. A. (2016). The macaque social responsiveness scale (MSRS): a rapid screening tool for assessing variability in the social responsiveness of rhesus monkeys (*Macaca mulatta*). *PloS ONE*, *11*(1), e0145956.
- Ferrari, P. F., Paukner, A., Ionica, C., & Suomi, S. J. (2009). Reciprocal face-to-face communication between rhesus macaque mothers and their newborn infants. *Current Biology*, *19*(20), 1768-1772. doi: 10.1016/j.cub.2009.08.055
- Framorando, D., George, N., Kerzel, D., & Burra, N. (2017). Straight gaze facilitates face processing but does not cause involuntary attentional capture. *Visual Cognition*, 1-11. doi: 10.1080/13506285.2017.1285840
- Frischen, A., Bayliss, A. P., & Tipper, S. P. (2007). Gaze cueing of attention: visual attention, social cognition, and individual differences. *Psychological Bulletin*, *133*(4), 694. doi: 10.1037/0033-2909.133.4.694
- Ganea, N., Hudry, K., Tucker, L., Charman, T., Johnson, M. H., & Senju, A. (2018). Development of adaptive communication skills in infants of blind parents. *Developmental Psychology*. <http://eprints.bbk.ac.uk/22166/>
- Gangi, D. N., Schwichtenberg, A. J., Iosif, A. M., Young, G. S., Baguio, F., & Ozonoff, S. (2018). Gaze to faces across interactive contexts in infants at heightened risk for autism. *Autism*, *22*(6), 763-768. doi: 10.1177/1362361317704421.
- Gelman, A., & Stern, H. (2006). The difference between “significant” and “not significant” is not itself statistically significant. *The American Statistician*, *60*(4), 328-331.
- Gliga, T., & Csibra, G. (2007). Seeing the face through the eyes: A developmental perspective on face expertise. *Progress in Brain Research*, *164*, 323-339. doi: 10.1016/S0079-6123(07)64018-7
- Gothard, K. M., Brooks, K. N., & Peterson, M. A. (2009). Multiple perceptual strategies used by macaque monkeys for face recognition. *Animal Cognition*, *12*(1), 155-167. doi: 10.1007/s10071-008-0179-7
- Gredebäck, G., Fikke, L. & Melinder, A. (2010). The development of joint visual attention: a longitudinal study of gaze following during interactions with mothers and strangers. *Developmental Science*, *13*, 839–848.
- Grossmann, T. (2017). The eyes as windows into other minds: an integrative perspective. *Perspectives on Psychological Science*, *12*(1), 107-121. doi: 10.1177/1745691616654457
- Hains, S. M., & Muir, D. W. (1996). Infant sensitivity to adult eye direction. *Child Development*, *67*(5), 1940–1951.

- Hirata, S., Fuwa, K., Sugama, K., Kusunoki, K., & Fujita, S. (2010). Facial perception of conspecifics: chimpanzees (*Pan troglodytes*) preferentially attend to proper orientation and open eyes. *Animal Cognition*, *13*(5), 679-688. doi: 10.1007/s10071-010-0316-y
- Hittelman, J. H., & Dickes, R. (1979). Sex differences in neonatal eye contact time. *Merrill-Palmer Quarterly of Behavior and Development*, *25*(3), 171-184.
- Hoehl, S., Reid, V. M., Parise, E., Handl, A., Palumbo, L., & Striano, T. (2009). Looking at eye gaze processing and its neural correlates in infancy—Implications for social development and autism spectrum disorder. *Child Development*, *80*(4), 968-985.
- Johnson, M. H. (2005). Subcortical face processing. *Nature Reviews Neuroscience*, *6*(10), 766-774. doi: 0.1038/nrn1766
- Johnson, M. H., Dziurawiec, S., Ellis, H., & Morton, J. (1991). Newborns' preferential tracking of face-like stimuli and its subsequent decline. *Cognition*, *40*(1-2), 1-19. doi: 10.1016/0010-0277(91)90045-6
- Johnson, M. H., Senju, A., & Tomalski, P. (2015). The two-process theory of face processing: modifications based on two decades of data from infants and adults. *Neuroscience & Biobehavioral Reviews*, *50*, 169-179. doi: 10.1016/j.neubiorev.2014.10.009
- Jones, W., & Klin, A. (2013). Attention to eyes is present but in decline in 2-6-month-old infants later diagnosed with autism. *Nature*, *504*(7480), 427-431.
- Kaburu, S. S., Paukner, A., Simpson, E. A., Suomi, S. J., & Ferrari, P. F. (2016). Neonatal imitation predicts infant rhesus macaque (*Macaca mulatta*) social and anxiety-related behaviours at one year. *Scientific Reports*, *6*, 34997.
- Kano, F., Shepherd, S. V., Hirata, S., & Call, J. (2018). Primate social attention: Species differences and effects of individual experience in humans, great apes, and macaques. *PLoS ONE*, *13*(2), e0193283.
- Leonard, T. K., Blumenthal, G., Gothard, K. M., & Hoffman, K. L. (2012). How macaques view familiarity and gaze in conspecific faces. *Behavioral Neuroscience*, *126*(6), 781-791. doi: 10.1037/a0030348
- Lickliter, R., & Bharick, L. E. (2016). Using an animal model to explore the prenatal origins of social development. In N. Reissland, & B. Kisilevsky (Eds.), *Fetal Development: Research on Brain and Behavior, Environmental Influences, and Emerging Technologies*. pp. 3-14. Switzerland: Springer.
- Morton, J., & Johnson, M. H. (1991). CONSPEC and CONLERN: a two-process theory of infant face recognition. *Psychological Review*, *98*(2), 164-181. doi: 10.1037/0033-295X.98.2.164
- Murray, L., De Pascalis, L., Bozicevic, L., Hawkins, L., Sclafani, V., & Ferrari, P. F. (2016). The functional architecture of mother-infant communication, and the development of infant social expressiveness in the first two months. *Scientific Reports*, *6*, 39019. doi: 10.1038/srep39019
- Muschinski, J., Feczko, E., Brooks, J. M., Collantes, M., Heitz, T. R., & Parr, L. A. (2016). The development of visual preferences for direct versus averted gaze faces in infant macaques (*Macaca mulatta*). *Developmental Psychobiology*, *58*(8), 926-936. doi: 10.1002/dev.21421

- Myowa-Yamakoshi, M., Tomonaga, M., Tanaka, M., & Matsuzawa, T. (2003). Preference for human direct gaze in infant chimpanzees (*Pan troglodytes*). *Cognition*, *89*(2), 113-124. doi: 10.1016/S0010-0277(03)00071-4
- Niedźwiecka, A., Ramotowska, S., & Tomalski, P. (2018). Mutual gaze during early mother-infant interactions promotes attention control development. *Child Development*. Advance Online Publication. doi: 10.1111/cdev.12830
- Nagy, E. (2011). The newborn infant: A missing stage in developmental psychology. *Infant and Child Development*, *20*(1), 3-19. doi: 10.1002/icd.683
- Paukner, A., Huntsberry, M. E., & Suomi, S. J. (2010). Visual discrimination of male and female faces by infant rhesus macaques. *Developmental Psychobiology*, *52*, 54-61. doi: 10.1002/dev.20412
- Paukner, A., Simpson, E. A., Ferrari, P. F., Mrozek, T., & Suomi, S. J. (2014). Neonatal imitation predicts how infants engage with faces. *Developmental Science*, *17*(6), 833-840. doi: 10.1111/desc.12207
- Paukner, A., Slonecker, E. M., Murphy, A. M., Wooddell, L. J., & Dettmer, A. M. (2018). Sex and rank affect how infant rhesus macaques look at faces. *Developmental Psychobiology*, *60*(2), 187-193. doi: 10.1002/dev.21579
- Pönkänen, L. M., Alhoniemi, A., Leppänen, J. M., & Hietanen, J. K. (2010). Does it make a difference if I have an eye contact with you or with your picture? An ERP study. *Social Cognitive and Affective Neuroscience*, *6*(4), 486-494.
- Rayson, H., Bonaiuto, J. J., Ferrari, P. F., & Murray, L. (2017). Early maternal mirroring predicts infant motor system activation during facial expression observation. *Scientific Reports*, *7*, 11738. doi: 10.1038/s41598-017-12097-w
- Reeb-Sutherland, B. C., Fifer, W. P., Byrd, D. L., Hammock, E. A. D., Levitt, P., & Fox, N. A. (2011). One-month-old human infants learn about the social world while they sleep. *Developmental Science*, *14*, 1134–1141.
- Reid, V. M., Dunn, K., Young, R. J., Amu, J., Donovan, T., & Reissland, N. (2017). The human fetus preferentially engages with face-like visual stimuli. *Current Biology*, *27*(12), 1825-1828.e3. doi: 10.1016/j.cub.2017.05.044
- Risko, E. F., Richardson, D. C., & Kingstone, A. (2016). Breaking the fourth wall of cognitive science: Real-world social attention and the dual function of gaze. *Current Directions in Psychological Science*, *25*(1), 70-74. doi: 10.1177/0963721415617806
- Rosati, A. G., Arre, A. M., Platt, M. L., & Santos, L. R. (2016). Rhesus monkeys show human-like changes in gaze following across the lifespan. In Proc. R. Soc. B (Vol. 283, No. 1830, p. 20160376). *Proceedings of the Royal Society B*, *283*, 20160376. doi: 10.1098/rspb.2016.0376
- Sclafani, V., Del Rosso, L. A., Seil, S. K., Calonder, L. A., Madrid, J. E., Bone, K. J., ... & Parker, K. J. (2016). Early predictors of impaired social functioning in male rhesus macaques (*Macaca mulatta*). *PLoS ONE*, *11*(10), e0165401.
- Senju, A., & Johnson, M. H. (2009). The eye contact effect: mechanisms and development. *Trends in Cognitive Sciences*, *13*(3), 127-134. doi: 10.1016/j.tics.2008.11.009
- Senju, A., Vermetti, A., Ganea, N., Hudry, K., Tucker, L., Charman, T., & Johnson, M. H. (2015). Early social experience affects the development of eye gaze processing. *Current Biology*, *25*(23), 3086-3091. doi: 10.1016/j.cub.2015.10.019

- Shultz, S., Klin, A., & Jones, W. (2018). Neonatal transitions in social behavior and their implications for autism. *Trends in Cognitive Sciences*, 22(5), 452-469. doi: 10.1016/j.tics.2018.02.012
- Sifre, R., Olson, L., Gillespie, S., Klin, A., Jones, W., & Shultz, S. (2018). A longitudinal investigation of preferential attention to biological motion in 2-to 24-month-old infants. *Scientific Reports*, 8(1), 2527. doi: 10.1038/s41598-018-20808-0
- Simpson, E. A., Jakobsen, K. V., Damon, F., Suomi, S. J., Ferrari, P. F., & Paukner, A. (2017). Face detection and the development of own-species bias in infant macaques. *Child Development*, 88, 103-113. doi: 10.1111/cdev.12565
- Simpson, E. A., Miller, G. M., Ferrari, P. F., Suomi, S. J., & Paukner, A. (2016). Neonatal imitation and early social experience predict gaze following abilities in infant macaques. *Scientific Reports*, 6, 20233. doi: 10.1038/srep20233
- Simpson, E. A., Nicolini, Y., Shetler, M., Suomi, S. J., Ferrari, P. F., & Paukner, A. (2016). Experience-independent sex differences in newborn macaques: Females are more social than males. *Scientific Reports*, 6, 19669. doi: 10.1038/srep19669
- Simpson, E. A., Suomi, S. J., & Paukner, A. (2016). Evolutionary relevance and experience contribute to face discrimination in infant macaques (*Macaca mulatta*). *Journal of Cognition and Development*, 17(2), 285-299.
- Sugita, Y. (2008). Face perception in monkeys reared with no exposure to faces. *Proceedings of the National Academy of Sciences*, 105(1), 394-398. doi: 10.1073/pnas.0706079105
- Sugita, Y. (2009). Innate face processing. *Current Opinion in Neurobiology*, 19(1), 39-44. Doi: 10.1016/j.conb.2009.03.001
- Symons, L., Hains, S., & Muir, D. (1998). Look at me: 5-month-old infant's sensitivity to very small deviations in eye-gaze during social interactions. *Infant Behavior and Development*, 21, 531-536.
- Teufel, C., Gutmann, A., Pirow, R., & Fischer, J. (2010). Facial expressions modulate the ontogenetic trajectory of gaze-following among monkeys. *Developmental Science*, 13(6), 913-922. doi: 10.1111/j.1467-7687.2010.00956.x
- Vernetti, A., Ganea, N., Tucker, L., Charman, T., Johnson, M. H., & Senju, A. (2018). Infant neural sensitivity to eye gaze depends on early experience of gaze communication. *Developmental Cognitive Neuroscience*, 34, 1-6.
- Vouloumanos, A., Hauser, M. D., Werker, J. F., & Martin, A. (2010). The tuning of human neonates' preference for speech. *Child Development*, 81, 517-527. doi: 10.1111/j.1467-8624.2009.01412.x
- Warton, D. I., & Hui, F. K. (2011). The arcsine is asinine: the analysis of proportions in ecology. *Ecology*, 92(1), 3-10.
- Wooddell, L. J., Simpson, E. A., Murphy, A. M., Dettmer, A. M., & Paukner, A. (in press). Interindividual differences in neonatal sociality and emotionality predict juvenile social status in rhesus monkeys. *Developmental Science*. Advance Online Publication. doi: 10.1111/desc.12749
- Xiao, N. G., Wu, R., Quinn, P. C., Liu, S., Tummeltshammer, K. S., Kirkham, N. Z., ... & Lee, K. (2018). Infants rely more on gaze cues from own-race than other-race adults for learning under uncertainty. *Child Development*, 89(3), e229-e244.

Yokoyama, T., Sakai, H., Noguchi, Y., & Kita, S. (2014). Perception of direct gaze does not require focus of attention. *Scientific Reports*, 4, 3858. doi: 10.1038/srep03858

Figure Legend

Figure 1. Direct gaze (top left) and averted gaze (top right) sample stimuli. Areas of interest (AOI) are shown around the head and eye zone regions (bottom).

Figure 2. Overall fixation proportion collapsed across age in nursery-reared infants. Infants looked for a greater proportion of time to the direct (dark blue) compared to averted (light blue) gaze faces, for both the face region (left plot) and eye region (right plot), $p_s < .05$. Solid horizontal lines indicate medians, the bottom and top of the boxes indicate 25th and 75th percentiles, respectively, and whiskers indicate the most extreme data point that is no more than $1.5 \times$ interquartile range (IQR) below the lower quartile and above the upper quartile. Dots indicate outliers, more than $1.5 \times$ IQR.

Figure 3. Fixation proportion across development (in weeks) in mother-reared (left graphs) and nursery-reared (right graphs) infants. Infants displayed increases in their proportions of time looking to the areas of interest (AOI) in the zones around the face (top graphs) and eyes (bottom graphs) with age, $p_s < .05$. Direct-gaze faces are represented with dark blue and averted-gaze faces are represented with light blue. Points reflect means and error bars reflect standard errors of the means.

Figure 4. Mean fixation proportion collapsed across age in mother-reared (left graph) and nursery-reared (right graph) infants, for the areas of interest (AOI) in the zones around the face and eyes, for direct-gaze (dark blue) and averted-gaze (light blue) faces. There was a stimulus type \times rearing interaction for the face looking time proportion, $p_s < .05$, indicating that mother-reared infants displayed a larger difference in attention between direct and averted gaze faces compared to nursery-reared infants. Error bars reflect standard error of the mean.

Tables

Table 1. *Longitudinal results for face*

Parameter	Fixation time				Fixation proportion			
	b	SE	95% CI	p	b	SE	95% CI	p
Intercept	.62	.12	[.37, .86]	< .001	.22	.02	[.17, .26]	< .001
Week	.06	.02	[.03, .10]	.001	.008	.003	[.001, .014]	.026
Direct	.16	.15	[-.13, .44]	.292	.08	.03	[.02, .13]	.014
Week * Direct	-.01	.02	[-.06, .03]	.630	-.005	.005	[-.014, .005]	.314

Note: Results of two linear mixed models, one predicting fixation time and one predicting fixation proportion. Models included a random intercept for subject, and a random slope for age (week), and an unstructured covariance matrix was used for the random effects. “Direct” refers to a dummy code for gaze direction (1 = direct gaze; 0 = averted gaze).

Table 2. *Longitudinal results for eyes*

Parameter	Fixation time				Fixation proportion			
	b	SE	95% CI	p	b	SE	95% CI	p
Intercept	.14	.07	[.01, .28]	.036	.05	.01	[.02, .07]	.001
Week	.03	.01	[.003, .05]	.028	.005	.003	[-.0004, .01]	.070
Direct	.06	.10	[-.12, .25]	.509	.03	.02	[-.007, .07]	.114
Week * Direct	.008	.02	[-.02, .04]	.620	.00	.003	[-.006, .006]	.974

Note: Results of two linear mixed models, one predicting fixation time and one predicting fixation proportion. Models included a random intercept for subject, and a random slope for age (week), and an unstructured covariance matrix was used for the random effects. “Direct” refers to a dummy code for gaze direction (1 = direct gaze; 0 = averted gaze).

Table 3. *Face AOI overall comparison with mother-reared infants*

Parameter	Fixation time				Fixation proportion			
	b	SE	95% CI	p	b	SE	95% CI	p
Intercept	.90	.09	[.71, 1.08]	< .001	.26	.02	[.23, .29]	< .001
Direct	.10	.10	[-.09, .29]	.312	.05	.02	[.01, .09]	.023
Mother	.21	.12	[-.03, .46]	.085	.08	.02	[.04, .12]	< .001
Direct * Mother	.32	.12	[.08, .56]	.008	.08	.03	[.02, .13]	.005

Note: Results of two linear mixed models, one predicting fixation time and one predicting fixation proportion. Models included a random intercept for subject, and a random slope for age, and an unstructured covariance matrix was used for the random effects. “Direct” refers to a dummy code for gaze direction (1 = direct gaze; 0 = averted gaze). “Mother” refers to a dummy code for rearing condition (1 = mother reared; 0 = nursery reared).

Table 4. *Eyes AOI overall comparison with mother-reared infants*

Parameter	Fixation time				Fixation proportion			
	b	SE	95% CI	p	b	SE	95% CI	p
Intercept	.29	.06	[.17, .40]	< .001	.07	.01	[.05, .10]	< .001
Direct	.10	.06	[-.01, .21]	.070	.03	.01	[.003, .06]	.028
Mother	.08	.08	[-.07, .23]	.310	.04	.02	[.01, .08]	.012
Direct * Mother	.10	.07	[-.04, .23]	.173	.03	.02	[-.003, .07]	.072

Note: Results of two linear mixed models, one predicting fixation time and one predicting fixation proportion. Models included a random intercept for subject, and a random slope for age, and an unstructured covariance matrix was used for the random effects. “Direct” refers to a dummy code for gaze direction (1 = direct gaze; 0 = averted gaze). “Mother” refers to a dummy code for rearing condition (1 = mother reared; 0 = nursery reared).

Supporting Information

Visual preferences for direct gaze faces in infant macaques (*Macaca mulatta*) with limited face exposure

Methods

Rearing Group Differences in Data Collection

Unfortunately, we were unable to use the same eye-tracking system for testing of nursery-reared and mother-reared infants. However, both eye-tracking systems have successfully been used previously with infant rhesus macaque monkeys (e.g., mother-reared: Parr et al., 2016a, 2016b; nursery-reared: Damon et al., 2017; Paukner et al., 2018; Simpson et al., 2016, 2017; Slonecker et al., 2018). The two groups of infants were tested in different ways, to ensure they were as calm and comfortable as possible, each with the most familiar caregiver: mother-reared infants clung to their lightly anesthetized mothers, while nursery-reared infants were held by a familiar human experimenter. Both nursery-reared and mother-reared infants underwent a standard 5-point calibration, then viewed the same stimuli, and were free to look as much or as little as they wanted. We currently have no way of directly comparing calibration quality across the two systems, but can anecdotally report that good calibration was achieved prior to data collection in both groups. (Without sufficient calibration quality, no data would have been collected.) Nursery infants were generally very attentive; only 3% of trials had no looking to the screen. All 16 nursery-reared infants contributed usable data at all ages. In the mother-reared sample, not all subjects could be tested at all ages, so between 5 and 15 infants contributed data at each age (for details, see Table 1 in Muschinski et al., 2016). For both mother-reared and nursery-reared infants, if an infant was sleepy or fussy, another attempt was made to test the infant on another day in the same week. Data were extracted using the same definition of a fixation across the two rearing groups (as reported on p. 5). For mother-reared infants, the data included 192 trials and 4,440 fixations at 23 time points (8 trials and 193 fixations per time point). For nursery-reared infants, the data included 250 trials and 3,764 fixations at 8 time points (31 trials and 470 fixations per time point). Mean fixation durations were .15 seconds for mother-reared infants ($SD=.18$) and .36 seconds for nursery-reared infants ($SD=.29$), $t(5894)=32.89$, $p<.001$, suggesting further work is necessary to assess differences in data quality across these infant groups and across these eye-tracking systems (Wass, Smith, & Johnson, 2013; for a report of data quality in nursery-reared infants compared to human infants using the TX300 eye-tracker, see Paukner, Johnson, & Simpson, under review).

Data Analysis

We calculated two dependent variables for each AOI (face and eyes) separately: (1) the proportion of total looking time (the sum of fixation times) to the AOI out of the total time looking to the screen (reported in the main paper), and (2) the sum of fixation times (in seconds) to a AOI (reported below).

Fixation Time Results

Main effects of stimulus category

First, we ran models using this basic structure to test for main effects of stimulus category (gaze direction) on our dependent variables in the nursery-reared infants. Stimulus category was dummy coded 1 for direct gaze, 0 for averted gaze.

In the model predicting fixation time to the face, the intercept was statistically significant, $b = .97$, $SE = .07$, 95% CI: [.82, 1.11], $p < .001$, indicating that, overall, infants spent a non-zero amount of time looking at faces. The coefficient for stimulus was not statistically significant, $b = .10$, $SE = .09$, 95% CI: [-.07, .27], $p = .255$, indicating that there was not a main effect for stimulus type (Gaze direction) on fixation time. Similarly, in the model predicting fixation time to the eyes, the intercept was statistically significant, $b = .14$, $SE = .06$, 95% CI: [.01, .28], $p = .042$, whereas the coefficient for stimulus was not, $b = .10$, $SE = .06$, 95% CI: [-.01, .21], $p = .070$ (Figure S1).

Longitudinal effects of stimulus category

Next, we added a fixed effect for week and a week \times stimulus type interaction to the models to test for longitudinal changes. We rescaled week by subtracting a constant of 2 so that the intercepts in the following models correspond to week 2, the earliest week of testing. Following Muschinski et al. (2016), we fit three full-factorial models of this type for each analysis: one with a linear growth term, one that added a quadratic growth term, and one that added a cubic growth term. Then, we used chi-square deviance tests to determine which of the nested models best fit the data. In all cases except for one, the model with the linear growth term fit the data best, so those are the models we report here. The one exception was the cubic model predicting percentage of time looking at the face—however, none of the terms in this model were statistically significant, indicating that the model was simply overfit due to the large amount of included terms. Therefore, we retained the linear model instead.

Full model results are presented in Table 1 (face) and Table 2 (eyes). In the model predicting fixation time for the face, the only notable significant effect was for week ($b = .06$, $p = .001$), indicating that, for each additional week (the data have a range of 11 weeks), the model predicts that fixation time on the face increases .06 seconds (Figure S2). Though this effect is rather small per one-week segment, over the 11 weeks it predicts an approximate doubling of the fixation time predicted at week 1 (the intercept, $b = .62$). In this model, there was not a significant effect of stimulus type ($b = .16$, $p = 2.92$). The model predicting fixation time for the eyes contained the same general results, with only a significant effect for week, but the effect was half as strong ($b = .03$, $p = .028$) as in the model predicting fixation time for the face (see Table 2 for details).

Comparison with mother-reared infants

We obtained the raw data from Muschinski et al. (2016) for the first 13 weeks of testing their mother-reared infants ($N = 20$), and we combined these data with our more recently collected nursery-reared infant data to directly test for overall effects of mother- vs. nursery-rearing. That is, beyond the apparent differences in growth trajectories between the two data sets², do mother-reared infants and nursery-reared infants differ in their overall preferences for direct and averted gaze faces?

² Importantly, the difference in the significance of the interactions and non-linear effects between the two datasets does not necessarily imply a statistically significant difference between the datasets in growth trajectories (Gelman & Stern, 2006). However, given the relatively small sample size here, we did not have adequate power to directly test for these differences with models including stimulus \times rearing condition \times dataset \times age interactions, nor more complicated models including the non-linear age terms.

To test this, we re-ran our four main effects models (fixation time and fixation proportion for face and eyes AOIs) on the combined dataset with an added predictor for rearing condition. Each model contained a dummy-coded predictor for stimulus category (gaze direction; 1 = direct gaze, 0 = averted gaze), rearing condition (1 = mother-reared; 0 = nursery-reared), and their interaction. Thus, the intercept in each model referred to nursery-reared infants looking at averted faces. Full model results are displayed in Table 3 (face AOI) and Table 4 (eyes AOI). Particularly noteworthy is that both for fixation time (Figure S3) and fixation proportion (Figure 4) for the face AOI, mother-reared infants, compared to nursery-reared infants, showed an increased preference for direct-gaze faces, as indicated by the significant stimulus category \times rearing condition interactions. Re-running these two models with the dummy code for rearing condition switched confirmed that, in mother-reared infants, there was a significant preference for direct gaze faces compared to averted gaze faces in the fixation proportion model ($b = .13$, $SE = .02$, 95% CI: [.09, .16], $p < .001$).

References

- Damon, F., Méary, D., Quinn, P. C., Lee, K., Simpson, E. A., Paukner, A., ... & Pascalis, O. (2017). Preference for facial averageness: Evidence for a common mechanism in human and macaque infants. *Scientific Reports*, *7*, 46303.
- Parr, L. A., Brooks, J. M., Jonesteller, T., Moss, S., Jordano, J. O., & Heitz, T. R. (2016a). Effects of chronic oxytocin on attention to dynamic facial expressions in infant macaques. *Psychoneuroendocrinology*, *74*, 149-157.
- Parr, L. A., Murphy, L., Feczko, E., Brooks, J., Collantes, M., & Heitz, T. R. (2016b). Experience-dependent changes in the development of face preferences in infant rhesus monkeys. *Developmental Psychobiology*, *58*(8), 1002-1018.
- Paukner, A., Johnson, S., & Simpson, E. A. (under review). Accuracy evaluation of the Tobii TX300 in human and monkey infants.
- Paukner, A., Slonecker, E. M., Murphy, A. M., Wooddell, L. J., & Dettmer, A. M. (2018). Sex and rank affect how infant rhesus macaques look at faces. *Developmental Psychobiology*, *60*(2), 187-193.
- Simpson, E. A., Jakobsen, K. V., Damon, F., Suomi, S. J., Ferrari, P. F., & Paukner, A. (2017). Face detection and the development of own-species bias in infant macaques. *Child Development*, *88*(1), 103-113.
- Simpson, E. A., Nicolini, Y., Shetler, M., Suomi, S. J., Ferrari, P. F., & Paukner, A. (2016). Experience-independent sex differences in newborn macaques: Females are more social than males. *Scientific Reports*, *6*, 19669.
- Slonecker, E. M., Simpson, E. A., Suomi, S. J., & Paukner, A. (2018). Who's my little monkey? Effects of infant-directed speech on visual retention in infant rhesus macaques. *Developmental Science*, *21*(2), e12519.
- Wass, S. V., Smith, T. J., & Johnson, M. H. (2013). Parsing eye-tracking data of variable quality to provide accurate fixation duration estimates in infants and adults. *Behavior Research Methods*, *45*(1), 229-250.

Figures




Figure S1. Overall fixation time (in seconds) collapsed across age in nursery-reared infants. There were no differences in the duration of time looking to direct (dark blue) and averted (light blue) gaze faces (left plot) or eyes (right plot), $p_s > .05$. Solid horizontal lines indicate medians, the bottom and top of the boxes indicate 25th and 75th percentiles, respectively, and whiskers indicate the most extreme data point that is no more than $1.5 \times$ interquartile range (IQR) below the lower quartile and above the upper quartile. Dots indicate outliers, more than $1.5 \times$ IQR.




Figure S2. Fixation time (in seconds) across development (in weeks) in mother-reared (left graphs) and nursery-reared (right graphs) infants. Infants displayed increases in attention to the face (top graphs) and eyes (bottom graphs) with age, $p_s < .05$. Direct-gaze faces are represented with dark blue and averted-gaze faces are represented with light blue. Points reflect means and error bars reflect standard errors of the means.




Figure S3. Mean fixation time (in seconds) collapsed across age in mother-reared (left graph) and nursery-reared (right graph) infants, for the areas of interest (AOI) in the zones around the face and eyes, for direct-gaze (dark blue) and averted-gaze (light blue) faces. There was a stimulus type \times rearing interaction for the face region looking time, $p_s < .05$, indicating that mother-reared infants displayed a larger difference in attention between direct and averted gaze faces compared to nursery-reared infants. Error bars reflect standard error of the mean.

Tables

Table S1. Face AOI longitudinal comparison with mother-reared infants

Parameter	Fixation time				Fixation proportion			
	b	SE	95% CI	p	b	SE	95% CI	p
Intercept	.61	.12	[.38, .84]	< .001	.22	.03	[.17, .27]	< .001
Week	.06	.02	[.02, .11]	.003	.01	.01	[-.003,.02]	.166
Gaze Direction	.16	.16	[-.16, .47]	.336	.08	.04	[.01,.14]	.035
Rearing	.43	.17	[.09, .77]	.013	.05	.04	[-.02, .12]	.174
Week *								
Rearing	-.06	.03	[-.12, .01]	.081	.01	.01	[-.01,.02]	.362
Week * Gaze								
Direction	-.01	.03	[-.06, .04]	.660	-.004	.01	[-.02, .01]	.391
Gaze Direction								
* Rearing	.60	.24	[.13, 1.06]	.013	.18	.05	[.07, .28]	.001
Week * Direct *								
Rearing	-.06	.04	[-.14, .01]	.105	-.02	.01	[-.04, -.01]	.005

Note: Results of two linear mixed models, one predicting fixation time and one predicting fixation proportion. Models included a random intercept for subject, and a random slope for age, and an unstructured covariance matrix was used for the random effects. "Gaze Direction" refers to a dummy code for stimulus category (1 = direct gaze; 0 = averted gaze). "Rearing" refers to a dummy code for rearing condition (1 = mother reared; 0 = nursery reared).

Table S2. Eyes AOI longitudinal comparison with mother-reared infants

Parameter	Fixation time				Fixation proportion			
	b	SE	95% CI	p	b	SE	95% CI	p
Intercept	.14	.07	[.00,.29]	.050	.05	.02	[.01, .08]	.010
Week	.03	.01	[.005, .05]	.019	.01	.003	[-.003, .01]	.066
Gaze Direction	.06	.10	[-.13, .25]	.513	.03	.02	[-.02, .08]	.191
Rearing	.19	.11	[-.03,.40]	.084	.04	.03	[-.008, .09]	.099
Week * Mother	-.02	.02	[-.06, .01]	.232	.001	.004	[-.01, .01]	.858
Week * Gaze								
Direction	.01	.02	[-.02, .04]	.623	.0001	.004	[-.01, .01]	.978
Gaze Direction								
* Rearing	.19	.14	[-.09, .47]	.184	.05	.03	[-.01 .12]	.123
Week * Direct *								
Rearing	-.02	.02	[-.07, .02]	.286	-.01	.005	[-.02, .005]	.283

Note: Results of two linear mixed models, one predicting fixation time and one predicting fixation proportion. Models included a random intercept for subject, and a random slope for age, and an unstructured covariance matrix was used for the random effects. "Gaze Direction" refers to a dummy code for stimulus category (1 = direct gaze; 0 = averted gaze). "Rearing" refers to a dummy code for rearing condition (1 = mother reared; 0 = nursery reared).