


# Journal Pre-proof

A brief psychological overview of disordered gaming

Kagan Kircaburun (Conceptualization)<ce:contributor-role>Writing - Original Draft<ce:contributor-role>Writing - Review and Editing), Halley M Pontes (Conceptualization)<ce:contributor-role>Writing - Review and Editing), Vasileios Stavropoulos (Conceptualization)<ce:contributor-role>Writing - Review and Editing), Mark D Griffiths (Conceptualization) (Supervision)<ce:contributor-role>Writing - Review and Editing)



PII: S2352-250X(20)30033-6  
DOI: <https://doi.org/10.1016/j.copsyc.2020.03.004>  
Reference: COPSYC 986

To appear in: *Current Opinion in Psychology*

Please cite this article as: Kircaburun K, Pontes HM, Stavropoulos V, Griffiths MD, A brief psychological overview of disordered gaming, *Current Opinion in Psychology* (2020), doi: <https://doi.org/10.1016/j.copsyc.2020.03.004>

This is a PDF file of an article that has undergone enhancements after acceptance, such as the addition of a cover page and metadata, and formatting for readability, but it is not yet the definitive version of record. This version will undergo additional copyediting, typesetting and review before it is published in its final form, but we are providing this version to give early visibility of the article. Please note that, during the production process, errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

© 2019 Published by Elsevier.

## **A brief psychological overview of disordered gaming**

Kagan Kircaburun, Halley M. Pontes, Ph.D\*, Vasileios Stavropoulos, PhD, Mark D. Griffiths, PhD

University of Tasmania, Launceston, Tasmania AUSTRALIA\*

\*Corresponding author:

### **Highlights**

- Prevalence of gaming disorder (GD) in representative samples range between 1%-5%
- There are many methodological limitations in gaming disorder assessment
- GD can co-occur with other psychiatric conditions and other addictive behaviors
- Internet gaming disorder and internet addiction are different nosological entities
- GD has had treatment success with pharmacotherapy and cognitive-behavioral therapy

### **Abstract**

In the latest (eleventh) revision of the International Classification of Diseases (ICD-11), the World Health Organization (WHO) recognized Gaming Disorder (GD) as an official diagnostic entity. Furthermore, in the latest (fifth) edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5), the American Psychiatric Association (APA) proposed Internet Gaming Disorder (IGD) as a tentative disorder in need of further study. The present review provides a brief analysis of the field. Even though there has been an ongoing debate concerning the proposed diagnostic criteria, there are now a number of assessment tools that have been developed using the diagnostic frameworks devised by the WHO and APA which have provided greater accuracy and consistency in IGD research. The prevalence rates of IGD reported in representative samples have ranged from approximately 1% to 5%. However, the discrepancy in the prevalence rates are mainly due to the reliance on non-representative samples, inconsistent assessment, and conceptual heterogeneity. In terms of treatment approaches, the literature suggests that pharmacological treatment and cognitive behavioral therapy-based treatments have been successfully employed to reduce the symptoms of IGD.

Despite the latest clinical advances in IGD research, there are still major drawbacks in treatment and existing intervention studies due to key limitations relating to sample sizes in treatment studies, small effect sizes, and scarcity of research on intervention studies. Taken together, these issues highlight the need for further studies into disordered gaming.

**Keywords:** internet gaming disorder, gaming disorder, gaming addiction, problematic gaming

### **Classification and diagnosis of disordered gaming**

In the past three decades, increased scholarly research on problematic gaming has led to formal diagnostic criteria for a now recognized disorder [1]. In 2013, the American Psychiatric Association [2] introduced ‘Internet Gaming Disorder’ (IGD), as a tentative disorder in the 5<sup>th</sup> edition of the *Diagnostic and Statistical Manual of Mental Disorders* (DSM-5). IGD was defined as “persistent and recurrent use of the internet to engage in games, often with other players, leading to clinically significant impairment or distress” (p. 795) [2]. More recently, the World Health Organization [3,4] followed by acknowledging ‘Gaming Disorder’ (GD) as a formal diagnosis in the 11<sup>th</sup> revision of the *International Classification of Diseases* (ICD-11).

The WHO outlined that GD manifests itself when the gaming behavior pattern is so severe that it negatively affects an individual’s personal, social, and/or educational/occupational activities in the previous 12-month period [3]. Similarly, the DSM-5 suggested that the provision of the IGD diagnosis required that, five out of nine criteria must be endorsed over a 12-month period (see Table 1 for the WHO and APA criteria). The nine IGD criteria proposed by the APA have been subject to extensive conceptual debate and empirical scrutiny [5,6], prompting concerns regarding the legitimacy of disordered gaming as a mental health issue [7,8]. Furthermore, IGD psychometric studies have illustrated inter-

criterion differences in diagnostic power [9-11]. Interestingly, the WHO [3,12] proposed a different set of three core criteria (see Table 1) to assess GD (with much less psychometric scrutiny to date).

Nevertheless, skepticism considering the acceptance of GD/IGD as a bona fide addictive disorder remains. Indicatively, the possibility of the proposed criteria leading to the over-diagnosis of passionate gamers as disordered has been highlighted [6]. These have been accompanied by broader concerns regarding over-pathologizing and defining new non-problematic behaviors as behavioral addictions including IGD [13]. Despite the continuing debates in the field, empirical evidence supports the sensitivity and specificity for most of the proposed nine symptoms of IGD in both clinical interview and cross-sectional designs [9,14].

-Table 1-

### **Prevalence and assessment of disordered gaming**

The prevalence of IGD has varied across studies mainly due to the various definitions, instruments, and/or self-selected samples used [15]. According to recent studies, prevalence rates of disordered gaming have been found to vary between 1% to 15%, with studies employing nationally representative samples reporting lower rates ranging from 1.2% to 5.5% across several countries [6,9,16-19]. In a recent meta-analysis, the estimated adolescent rate for GD was 4.6% globally [19,20].

Despite these discrepancies, significant progress has been made because psychometric tools employed prior to the introduction of IGD in DSM-5 involved several weaknesses [21; see Table 2]. Since then, a number of robust psychometric tools based on the nine IGD criteria in the DSM-5 have been developed [9,15,22-24]. These IGD-based assessment tools have been psychometrically assessed across countries and over time indicating the sufficient psychometric performance of the nine criteria suggested by DSM-5 [25,26].

The latest wave of advances in the psychometric assessment of disordered gaming was prompted by the introduction of the WHO's criteria for GD. More specifically, the Gaming Disorder Test (GDT) [10] is a brief standardized psychometric tool that includes four items assessing the key defining features of GD as specified in the ICD-11. The first three items of the GDT were developed to map on the following clinical criteria: (i) impaired control over gaming (i.e., *"I have had difficulties controlling my gaming activity"*), (ii) increased priority given to gaming (*"I have given increasing priority to gaming over other life interests and daily activities"*), and (iii) continuation despite negative consequences (*"I have continued gaming despite the occurrence of negative consequences"*). The fourth item of the GDT reflects the experience of major problems in life when the severity of GD is markedly high, reflecting the potential functional impairments that GD can cause at extreme levels (*"I have experienced significant problems in life [e.g., personal, family, social, education, occupational] due to the severity of my gaming behavior"*).

A recent systematic review study evaluated a total of 32 English-language psychometric tools for disordered gaming published across 320 studies using 462,249 participants [27]. In their study, King and colleagues [27] suggested the Assessment of Internet and Computer Addiction Scale-Gaming (AICA-Sgaming), Game Addiction Scale (GAS-7), Internet Gaming Disorder Test (IGDT-10), Internet Gaming Disorder Scale–Short-Form (IGDS9-SF), and the Internet Gaming Disorder Scale (IGD-9) were the most psychometrically consistent tools in the assessment of disordered gaming. Despite these important developments, existing scholarly controversies and concerns [28] imply that further large-scale research is still needed to bridge the existing gaps in the field.

-Table 2-

### **Etiological factors in disordered gaming**

Many studies have attempted to provide empirical insight concerning the etiology of disordered gaming in light of key individual differences such as personality factors and psychiatric comorbidities. A recent review that identified 21 studies suggesting that disordered gaming was negatively correlated with extraversion, conscientiousness, agreeableness, histrionic traits, openness to experience, grit, oppositional traits, and self-demeaning traits [29]. The same review concluded that disordered gaming was positively correlated with negative valence, neuroticism, sensation seeking, inhibition, introversion, egotism, narcissism, sadism, Type D personality traits, negative affectivity, detachment, antagonism, disinhibition, psychoticism, novelty-seeking, harm avoidance, and schizotypal traits [29]. However, the authors emphasized that the reviewed studies presented with several types of limitations including (i) sampling problems (e.g., non-probability sampling, sampling homogeneity, low sample sizes), (ii) measurement problems (e.g., use of non-validated and modified measurements), and (iii) lack of longitudinal data.

Additionally, several studies have explored the relationship between disordered gaming and psychiatric comorbidities. A recent review study examining 24 studies identified significant correlations between disordered gaming and depression, anxiety, attention deficit hyperactivity disorder (ADHD), social phobia/anxiety, and obsessive-compulsive symptoms [30]. The authors concluded that disordered gaming was most highly associated with anxiety and least associated with social anxiety/phobia. However, direction of these associations remains unclear [30]. Nevertheless, the authors emphasized the homogeneity of the geographical distribution of the research in disordered gaming, indicating comorbidity of disordered gaming and psychiatric distress is an emerging global problem.

More recently, Burleigh, Griffiths, Sumich, Stavropoulos, and Kuss [31] reviewed 20 studies and reported that disordered gaming can co-occur with other potential behavioral addictions (e.g., social media addiction, internet addiction, and gambling disorder) mostly

among adolescents, and potential substance addictions (e.g., alcohol, nicotine, and caffeine use disorder) mostly among adult gamers. The authors argued that the co-occurrence of different addictions might be related to the use of (i) maladaptive coping strategies (e.g., emotional avoidance) as a means to avoid unpleasant affective states and associated mental disorders, and (ii) diminished emotional regulation which leads to engagement in risky behaviors including elevated substance use [31]. A recent largescale study comparing the co-occurrence of psychiatric symptoms in gamers assessed with both the APA and WHO diagnostic frameworks for disordered gaming found that both diagnostic frameworks were relatively consistent in predicting the potential psychopathological symptoms associated with disordered gaming, further supporting the utility of the APA and WHO diagnostic frameworks in the assessment of disordered gaming and its accompanying comorbidities [32].

### **Differential diagnosis of gaming disorder and excessive internet behaviors**

Before the inclusion of IGD in the DSM-5, scholars argued whether internet addiction should have been considered as a separate disorder [33,34]. More than two decades ago, Griffiths [35] argued that individuals are not addicted *to* the internet but to the specific activities *on* the internet. Therefore, internet gaming addicts should not be classed as internet addicts but disordered gamers who use the internet to play games, indicating that IGD should be considered as disordered gaming rather than a sub-type of internet addiction [36].

Recent empirical research concerning online addictions has separated unspecified internet use disorder and disordered use of specific online activities [37]. Such research highlights that different types of addictions to unspecified/specific activities present with shared and unique individual difference predictors, indicating that unspecified internet use disorder and specific internet use disorders (e.g., social media, gaming, gambling, pornography use, and shopping) are conceptually different behaviors [37]. Another cross-cultural study investigating the relationship between generalized and specific internet addiction using data

from Germany, Taiwan, Sweden, and China concluded that internet addiction, internet gaming addiction, internet shopping addiction, social media addiction, and internet pornography use addiction were all overlapping but distinct forms of behaviors [38]. Therefore, it appears to be well established that disordered gaming and internet addiction are different nosological entities, an important distinction that can facilitate correct clinical assessment and identification of disordered gaming among gamers.

### **Treatment of disordered gaming**

In the light of the emergence of GD related clinical cases, several types of treatment have been reported [39,40]. In some of these attempts, pharmacological treatment approaches administering different drugs including bupropion, escitalopram, methylphenidate, and atomoxetine were employed. These approaches have been reported as successfully decreasing IGD symptoms with 6 to 12 week courses of medication trials (based on the use of drugs traditionally targeting depression or ADHD; [41,42]). Nevertheless, with the exception of two studies using randomized designs with control groups, most of these findings were compromised by the absence of control groups [41-44].

Besides pharmacotherapy, psychological treatment approaches have also been employed to treat GD. Several attempts with variations of cognitive-behavioral therapy (CBT) including mindfulness, gaming-specific CBT, CBT focusing on craving, and standard CBT have been reported to have promising results [43-48]. These studies comprised both randomized and non-randomized controlled trials and all of them successfully managed to reduce individuals' time spent on gaming and disordered gaming symptoms. Interestingly, combined pharmacological and CBT IGD interventions have been accompanied with more efficient and successful results than using only medicine or only psychotherapy [43]. Despite



these positive developments, IGD treatment studies present with several limitations ([39,40]; see Table 3).

-Table 3-

### **Conclusion and further studies**

Various inconsistencies and psychometric weaknesses have been reported by previous studies [27]. Assessment and measurement consistency in regards to the officially introduced criteria in DSM-5 [2] and ICD-11 [3] is essential to avoid major limitations in GD research, which will facilitate researchers in examining GD and its psychological and social detrimental effects on society. As noted above, methodological shortcomings have been reported for almost all IGD treatment studies and there are still large inconsistencies on the efficacy and treatment effectiveness of the interventions. Therefore, further largescale research is still needed to bridge the existing gaps in the field. Finally, the adoption of the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) checklist methodology is required to have a better quality of reporting of observational IGD studies and their strengths, weaknesses, and generalizability [49].

### **Credit Author Statement**

**Kagan Kircaburun:** Conceptualization, Writing - Original Draft, Writing - Review & Editing.

**Halley M. Pontes:** Conceptualization, Writing - Review & Editing.

**Vasileios Stavropoulos:** Conceptualization, Writing – Reviewing & Editing.

**Mark D. Griffiths:** Conceptualization, Supervision, Writing - Review & Editing.

### **Declaration of interests**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Journal Pre-proof

## References

- of interest
- of outstanding interest

## References

1. Griffiths, MD, Kuss, DJ, King, DL: **Video game addiction: Past, present and future.** *Curr Psych Rev* 2012, **8**:308-318.
2. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (5th ed.)*; Author: Arlington, VA, 2013.
3. World Health Organization. ICD-11 Beta Draft: Gaming Disorder. Available online: <http://apps.who.int/classifications/icd11/browse/f/en#/http%3a%2f%2fid.who.int%2fid%2fentity%2f1448597234> (accessed on March 13).
4. World Health Organization. Gaming disorder. Available online: <https://www.who.int/features/qa/gaming-disorder/en/> (accessed on March 13).
5. Dowling, NA: **Issues raised by the DSM-5 internet gaming disorder classification and proposed diagnostic criteria.** *Addiction* 2014, **109**:1408-1409.
6. Griffiths, MD, Van Rooij, AJ, Kardefelt-Winther, D, Starcevic, V, Király, O, Pallesen, S, Müller, K, Dreier, M, Carras, M, Prause, N, et al.: **Working towards an international consensus on criteria for assessing Internet Gaming Disorder: A critical commentary on Petry et al. (2014).** *Addiction* 2016, **111**:167-175.
7. Aarseth, E, Bean, AM, Boonen, H, Colder, CM, Coulson, M, Das, D, Deleuze, J, Dunkels, E, Edman, J, Ferguson, CJ, et al.: **Scholars' open debate paper on the World Health Organization ICD-11 Gaming Disorder proposal.** *J Beh Add* 2016, **6**:267-270.
8. Griffiths, MD, Kuss, DJ, Lopez-Fernandez, O, Pontes, HM: **Problematic gaming exists and is an example of disordered gaming: Commentary on: Scholars' open debate paper on the World Health Organization ICD-11 Gaming Disorder proposal (Aarseth et al.).** *J Beh Add* 2017, **6**:296-301.
9. Lemmens, JS, Valkenburg, PM, Gentile, DA: **The Internet Gaming Disorder Scale.** *Psych Ass* 2015, **27**:567-582.
10. Pontes, HM, Schivinski, B, Brzozowska-Woś, M, Stavropoulos, V: **Laxer clinical criteria for gaming disorder may hinder future efforts to devise an efficient diagnostic approach: A tree-based model study.** *J Clin Med* 2019, **8**:1730.

11. Schivinski, B, Brzozowska-Woś, M, Buchanan, EM, Griffiths, MD, Pontes, HM: **Psychometric assessment of the Internet Gaming Disorder diagnostic criteria: An Item Response Theory study.** *Add Beh Rep* 2018, **8**:176-184.
12. World Health Organization. 6C51 Gaming disorder. Available online: <https://icd.who.int/browse11/l-m/en#/http://id.who.int/icd/entity/1448597234> (accessed on February 27).
13. Billieux, J, Schimmenti, A, Khazaal, Y, Maurage, P, Heeren, A: **Are we overpathologizing everyday life? A tenable blueprint for behavioral addiction research.** *J Beh Add* 2015, **4**:119-123.
14. Ko, CH, Yen, JY, Chen, SH, Wang, PW, Chen, CS, Yen, CF: **Evaluation of the diagnostic criteria of Internet Gaming Disorder in the DSM-5 among young adults in Taiwan.** *Journal of Psychiatric Research* 2014, **53**:103-110.
15. Pontes, HM, Griffiths, MD: **Measuring DSM-5 Internet Gaming Disorder: Development and validation of a short psychometric scale.** *Comp Hum Beh* 2015, **45**:137-143.
16. Pontes, HM, Macur, M, Griffiths, MD: **Internet Gaming Disorder among Slovenian primary schoolchildren: Findings from a nationally representative sample of adolescents.** *J Beh Add* 2016, **5**:304-310.
17. Wartberg, L, Kriston, L, Thomasius, R: **Internet gaming disorder and problematic social media use in a representative sample of German adolescents: Prevalence estimates, comorbid depressive symptoms and related psychosocial aspects.** *Comp Hum Beh* 2020, **103**:31-36.
18. Wu, AMS, Chen, JH, Tong, KK, Yu, S, Lau, JTF: **Prevalence and associated factors of Internet gaming disorder among community dwelling adults in Macao, China.** *J Beh Add* 2018, **7**:62-69.
- 19. Fam, JY: **Prevalence of internet gaming disorder in adolescents: A meta-analysis across three decades.** *Scan J Psych* 2018, 10.1111/sjop.12459. This study reviewed the prevalence rates of Internet Gaming Disorder among adolescents. A total of 16 studies met the inclusion criteria for this review and the authors reported that the prevalence of Internet Gaming Disorder among adolescents was 4.6%, with male adolescents showing higher prevalence rates (6.8%) than female adolescents (1.3%).
20. Olson, DH, Barnes, H: **Family Communication Scale.** *University of Minnesota* 1995.

21. King, DL, Haagsma, MC, Delfabbro, PH, Gradisar, M, Griffiths, MD: **Toward a consensus definition of pathological video-gaming: A systematic review of psychometric assessment tools.** *Clin Psych Rev* 2013, **33**:331-342.
22. Király, O, Slezcka, P, Pontes, HM, Urbán, R, Griffiths, MD, Demetrovics, Z: **Validation of the ten-item Internet Gaming Disorder Test (IGDT-10) and evaluation of the nine DSM-5 Internet Gaming Disorder criteria.** *Add Beh* 2017, **64**:253–260.
23. Pontes, HM, Király, O, Demetrovics, Z, Griffiths, MD: **The conceptualisation and measurement of DSM-5 Internet Gaming Disorder: The development of the IGD-20 Test.** *PLoS ONE* 2014, **9**:e110137.
24. Van Rooij, AJ, Schoenmakers, TM, van de Mheen, D: **Clinical validation of the C-VAT 2.0 assessment tool for gaming disorder: A sensitivity analysis of the proposed DSM-5 criteria and the clinical characteristics of young patients with ‘video game addiction’.** *Add Beh* 2017, **64**:269-274.
25. Stavropoulos, V, Bamford, L, Beard, C, Gomez, R, Griffiths, MD: **Test-retest measurement invariance of the nine-item Internet Gaming Disorder scale in two countries: A preliminary longitudinal study.** *Int J Ment Health Add* 2019, 10.1007/s11469-019-00099-w.
26. Stavropoulos, V, Beard, C, Griffiths, MD, Buleigh, T, Gomez, R, Pontes, HM: **Measurement invariance of the Internet Gaming Disorder Scale–Short-Form (IGDS9-SF) between Australia, the USA, and the UK.** *Int J Ment Health Add* 2018, **16**:377-392.
- 27. King, DL, Chamberlain, SR, Carragher, N, Billieux, J, Stein, D, Mueller, K, Potenza, MN, Rumpf, HJ, Saunders, J, Starcevic, V, et al.: **Screening and assessment tools for gaming disorder: A comprehensive systematic review.** *Clin Psych Rev* 2020, 10.1016/j.cpr.2020.101831:101831. This study reviews a total of 32 psychometric tests developed to assess Gaming Disorder and Internet Gaming Disorder that were published across 320 studies. Although the review suggests that no single tool emerged as the clearly optimal choice, the AICA-Sgaming, GAS-7, IGDT-10, IGDS9-SF, and Lemmens IGD-9 scales had greater evidential support for their psychometric properties.
28. Kuss, DJ, Griffiths, MD, Pontes, HM: **Chaos and confusion in DSM-5 diagnosis of Internet Gaming Disorder: Issues, concerns, and recommendations for clarity in the field.** *J Beh Add* 2017, **6**:103-109.

29. Şalvarlı, Şİ, Griffiths, MD: **Internet gaming disorder and its associated personality traits: A systematic review using PRISMA guidelines.** *Int J Ment Health Add* 2019, 10.1007/s11469-019-00081-6.
- 30. González-Bueso, V, Santamaría, J, Fernández, D, Merino, L, Montero, E, Ribas, J: **Association between Internet Gaming Disorder or pathological video-game use and comorbid psychopathology: A comprehensive review.** *Int J Environ Res* 2018, **15**:668. The goal of this study was to review systematically the literature on Internet Gaming Disorder (IGD) to determine its association with different types of psychopathology. After reviewing 24 studies, the authors found that Internet Gaming Disorder is often associated with anxiety, depression, attention deficit hyperactivity disorder, and with social phobia/anxiety and obsessive-compulsive symptoms.
31. Burleigh, TL, Griffiths, MD, Sumich, A, Stavropoulos, V, Kuss, DJ: **A systematic review of the co-occurrence of Gaming Disorder and other potentially addictive behaviors.** *Curr Add Rep* 2019, 10.1007/s40429-019-00279-7.
- 32. Montag, C, Schivinski, B, Sariyska, R, Kannen, C, Demetrovics, Z, Pontes, HM: **Psychopathological Symptoms and Gaming Motives in Disordered Gaming—A Psychometric Comparison between the WHO and APA Diagnostic Frameworks.** *J Clin Med* 2019, **8**:1691. This study compared the American Psychiatric Association diagnostic criteria for Internet Gaming Disorder against the diagnostic framework for Gaming Disorder developed by the World Health Organization. The authors found that both diagnostic frameworks are relatively consistent in predicting psychopathological symptoms and that minor prevalence rates discrepancies exist across the two diagnostic frameworks.
33. Griffiths, MD, King, DL, Demetrovics, Z: **DSM-5 Internet Gaming Disorder needs a unified approach to assessment.** *Neuropsychiatry* 2014, **4**:1-4.
34. Petry, NM, O'Brien, CP: **Internet Gaming Disorder and the DSM- 5.** *Addiction* 2013, **108**:1186–1187.
35. Griffiths, MD: **Internet addiction-time to be taken seriously?** *Add Res* 2000, **8**:413-418.
36. Griffiths, MD: **Conceptual issues concerning internet addiction and internet gaming disorder: Further critique on Ryding and Kaye (2017).** *Int J Ment Health Add* 2018, **16**:233-239.

37. Sindermann, C, Sariyska, R, Lachmann, B, Brand, M, Montag, C: **Associations between the dark triad of personality and unspecified/specific forms of Internet-use disorder.** *J Beh Add* 2018, **7**:985-992.
38. Montag, C, Bey, K, Sha, P, Li, M, Chen, YF, Liu, WY, Zhu, YK, Li, CB, Markett, S, Keiper, J, et al.: **Is it meaningful to distinguish between generalized and specific Internet addiction? Evidence from a cross-cultural study from Germany, Sweden, Taiwan and China.** *Asia Pac Psych* 2014, **7**:20-26.
39. King, DL, Delfabbro, PH: **Internet Gaming Disorder Treatment: A Review of Definitions of Diagnosis and Treatment Outcome.** *The J Clin Psych* 2014, **70**:942-955.
- 40. Zajac, K, Ginley, MK, Chang, R, Petry, NM: **Treatments for internet gaming disorder and internet addiction: A systematic review.** *Psych Addict Behav* 2017, **31**:979-994. This study conducted a systematic review of the evidence on the treatment of Internet Gaming Disorder. The study identified a total of 22 studies evaluating treatments for Internet Gaming Disorder using medication, cognitive behavioral psychotherapy, and other interventions and psychosocial treatments. The authors concluded that the existing methodological flaws among such studies prevent robust conclusions about the efficacy of any treatment developed for Internet Gaming Disorder.
41. Nam, B, Bae, S, Kim, SM, Hong, JS, Han, DH: **Comparing the effects of Bupropion and Escitalopram on excessive internet game play in patients with major depressive disorder.** *Clin Psychopharmacol Neurosci* 2017, **15**:361-368.
42. Park, JH, Lee, YS, Sohn, JH, Han, DH: **Effectiveness of atomoxetine and methylphenidate for problematic online gaming in adolescents with attention deficit hyperactivity disorder.** *Hum Psychopharmacol Clin Exp* 2016, **31**:427-432.
43. Kim, SM, Han, DH, Lee, YS, Renshaw, PF: **Combined cognitive behavioral therapy and bupropion for the treatment of problematic on-line game play in adolescents with major depressive disorder.** *Comp Hum Beh* 2012, **28**:1954-1959.
44. Song, J, Park, JH, Han, DH, Roh, S, Son, JH, Choi, TY, Lee, H, Kim, TH, Lee, YS: **A comparative study of the effects of bupropion and escitalopram on Internet gaming disorder.** *Psych Clin Neuro* 2016, **70**:527-535.
45. Li, W, Garland, EL, McGovern, P, O'Brien, JE, Tronnier, C, Howard, MO: **Mindfulness-oriented recovery enhancement for internet gaming disorder in U.S. adults: A stage I randomized controlled trial.** *Psych Addict Behav* 2017, **31**:393-402.

46. Torres-Rodríguez, A, Griffiths, MD, Carbonell, X, Oberst, U: **Treatment efficacy of a specialized psychotherapy program for Internet Gaming Disorder.** *J Beh Add* 2018, 10.1556/2006.7.2018.111:1-14.
47. Yao, YW, Chen, PR, Li, CSR, Hare, TA, Li, S, Zhang, JT, Liu, L, Ma, SS, Fang, XY: **Combined reality therapy and mindfulness meditation decrease intertemporal decisional impulsivity in young adults with Internet Gaming Disorder.** *Comp Hum Beh* 2017, **68**:210-216.
48. Zhang, JT, Yao, YW, Potenza, MN, Xia, CC, Lan, J, Liu, L, Wang, LJ, Liu, B, Ma, SS, Fang, XY: **Effects of craving behavioral intervention on neural substrates of cue-induced craving in Internet Gaming Disorder.** *Neuroimage Clin* 2016, **12**:591-599.
49. von Elm, E, Altman, DG, Egger, M, Pocock, SJ, Gøtzsche, PC, Vandenbroucke, JP, for the, SI: **The Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) Statement: Guidelines for Reporting Observational Studies.** *Ann Intern Med* 2007, **147**:573-577.



## TABLES

**Table 1.** Official diagnostic criteria for gaming disorder

	<i>Diagnostic criteria</i>
IGD (APA, 2013)	
1	Excessive preoccupation with gaming.
2	Experiencing withdrawal symptoms when unable to engage in gaming.
3	Increasing levels of gaming over time.
4	Experiencing relapse when attempting to cease or reduce the behavior.
5	Losing interest in previous hobbies because of gaming.
6	Continuing to engage in gaming despite problems.
7	Deceiving significant others about the amount of time spent on gaming.
8	Using gaming to achieve a positive mood.
9	Risking, jeopardizing, or losing a job or relationship due to gaming.
GD (WHO, 2019)	
1	Impaired control over gaming (e.g., onset, frequency, intensity, duration, termination, context).
2	Increasing priority given to gaming to the extent that gaming takes precedence over other life interests and daily activities.
3	Continuation or escalation of gaming despite occurrence of negative consequences.

*Note.* IGD = Internet Gaming Disorder; GD = Gaming Disorder; APA = American Psychiatric Association; WHO = World Health Organization.

**Table 2.** Weaknesses of gaming disorder psychometric assessment instruments

Weaknesses (King, Haagsma, Delfabbro, Gradisar, & Griffiths, 2013)	
1	Inconsistency in the core criteria used for disordered gaming.
2	Lack of a temporal dimension in the assessment of disordered gaming.
3	Variety in the cut-off scores adopted to identify disordered gaming.
4	Inadequate inter-rater reliability and predictive validity.
5	Low consistency in the dimensionality of such tools.

Journal Pre-proof

**Table 3.** Limitations of gaming disorder treatment and intervention studies

Limitations (King & Delfabbro, 2014)	
1	The majority of the (reviewed) treatment studies did not tend to use an equivalent diagnostic method for disordered gaming.
2	Formative change in diagnostic status at post-treatment tended to not be assessed.
3	Inadequate follow-up duration was used to assess relapse and remission.
4	Researchers limited posttreatment assessment mostly to disordered gaming symptomatology, comorbidity, and frequency of gaming.
Limitations (Zajac et al., 2017)	
1	Methodological flaws (e.g., small sample sizes, lack of control groups, lack of treatment adherence information).
2	A lack of consistent definitions of gaming disorder and assessment tools.
Limitations (Zajac et al., 2020)	
1	Pharmacological treatment research is inconclusive with the drugs being promising but remaining in early evaluation stages.
2	Cognitive-behavioral therapy treatment warrants more research because of the mixed results reported on its effectiveness.
3	Specific weaknesses of prior studies, including lack of appropriate control groups, non-random assignment to treatment conditions, and small sample sizes, prevent strong and conclusive inferences about the efficacy of disordered gaming treatments.