

1 A Narrative Review on Female Physique Athletes: The Physiological  
2 and Psychological Implications of Weight Management Practices

3 Nura Alwan<sup>1</sup>, Samantha L. Moss<sup>2</sup>, Kirsty J. Elliott-Sale<sup>3</sup>, Ian G.  
4 Davies<sup>1</sup>, and Kevin Enright<sup>1</sup>

5

6 <sup>1</sup>School of Sport Studies, Leisure and Nutrition, Liverpool John  
7 Moores University, UK.

8 <sup>2</sup>Department of Sport and Exercise Sciences, University of Chester,  
9 UK.

10 <sup>3</sup>Musculoskeletal Physiology Research Group, Sport, Health and  
11 Performance Enhancement Research Centre, Nottingham Trent  
12 University, UK.

13

14 **Running head:** Health considerations in female physique athletes

15

16 **Corresponding author:**

17 Nura Alwan

18 <sup>1</sup>School of Sport Studies, Leisure and Nutrition,

19 Liverpool John Moores University,

20 Barkhill Road,

21 Liverpool,

22 L17 6BD

23 United Kingdom

24 Email: [N.alwan@2016.ljmu.ac.uk](mailto:N.alwan@2016.ljmu.ac.uk)

25

**26 Abstract**

27 Physique competitions are events in which aesthetic appearance and  
28 posing ability are valued above physical performance. Female  
29 physique athletes are required to possess high lean body mass and  
30 extremely low fat mass in competition. As such, extended periods of  
31 reduced energy intake and intensive training regimens are utilised with  
32 acute weight loss practices at the end of the pre-competition phase.  
33 This represents an increased risk for chronic low energy availability  
34 and associated symptoms of *Relative Energy Deficiency in Sport*,  
35 compromising both psychological and physiological health. Available  
36 literature suggests that a large proportion of female physique athletes  
37 report menstrual irregularities (*e.g.*, amenorrhea and oligomenorrhea),  
38 which are unlikely to normalise immediately post-  
39 competition. Furthermore, the tendency to reduce intakes of numerous  
40 essential micronutrients is prominent among those using restrictive  
41 eating patterns. Following competition reduced resting metabolic rate,  
42 and hyperphagia, are also a concern for these female athletes, which  
43 can result in frequent weight cycling, distorted body image and  
44 disordered eating/eating disorders. Overall, female physique athletes  
45 are an understudied population and the need for more robust studies to  
46 detect low energy availability and associated health effects is  
47 warranted. This narrative review aims to define the natural female  
48 physique athlete, explore some of the physiological and psychological  
49 implications of weight management practices experienced by female  
50 physique athletes and propose future research directions.

51

52 **Keywords**

53 Fat loss, low energy availability, physique events, body composition,

54 nutrition

55

56

57

58

59

60

61

62

63

64

65

66

67

68

69

70

71

72

## 73 **Background**

74 Physique competitions are events in which competitors are judged on  
75 aesthetic appearance rather than on physical performance. Natural  
76 (*i.e.*, drug-free) physique competitions have evolved dramatically in  
77 recent years, with a growth in organisations, contests and classes  
78 (Halliday et al., 2016). The International Federation of Body Building  
79 and Fitness (IFBB) hosts over 2,000 competitions annually, in 196  
80 affiliated countries. Approximately 1,300 female and male athletes  
81 competed at the World Fitness Championships in 2017 (Rowbottom,  
82 2017) and this number is anticipated to increase, with around 1,000  
83 new members joining the sport each year (Parish et al., 2010).

84 Female physique (FP) athletes have aspirations of achieving a lean and  
85 muscular body composition for competitive success (Halliday et al.,  
86 2016). Preparing for a natural physique competition provides a myriad  
87 of health benefits including improvement in cardiovascular status  
88 (Kistler et al., 2014; Robinson et al., 2015), muscle strength  
89 (Campbell et al., 2018), increasing feelings of accomplishment and  
90 transient improvements in self-esteem (Aspridis et al., 2014; Baghurst  
91 et al., 2014; Probert et al., 2007). Despite these positive outcomes,  
92 numerous unfavorable consequences also exist, including, but not  
93 limited to: diminished levels of reproductive hormones (Hulmi et al.,  
94 2016) and symptoms of disordered eating and eating disorders  
95 (DE/ED) (Walberg and Johnston, 1991). Available research on FP  
96 athletes reveals prolonged periods of sustained energy restriction and  
97 intensive training regimens in an attempt to acquire and maintain a  
98 lean body composition, indicating an increased risk of low energy  
99 availability (LEA) and its associated effects (Fagerberg, 2017). For a

100 thorough understanding of the existence, aetiologies and clinical  
101 consequences of LEA, readers are directed to the review by Loucks et  
102 al. (2011).

103 Prolonged periods of LEA with or without disordered eating,  
104 menstrual dysfunction and low bone mineral density is termed the  
105 Female Athlete Triad (Triad), representing a medical condition  
106 observed in females who perform high levels of physical activity  
107 (Manore, 2007). In order to describe the wide range of physiological,  
108 psychological and performance-related impairments associated with  
109 LEA, the International Olympic Committee introduced the concept of  
110 Relative Energy Deficiency in Sport (RED-S) in 2014 (Mountjoy et  
111 al., 2014). Considering the health risks of RED-S, and the increasing  
112 participation of females in physique events, the purpose of this  
113 narrative review was three-fold: 1. to define the natural female  
114 physique athlete; 2. to explore the physiological and psychological  
115 implications of the weight management practices experienced by the  
116 natural FP athlete; 3. to address future research directions.

117

118

119

120

121

122

123

124

## 125 **Literature Search**

126 A literature search was conducted using databases: PubMed, Web of  
127 Science, Google Scholar, and SPORTDiscus (via EBSCO) up to 10<sup>th</sup>  
128 September 2018. Despite slight variation in the terminology used for  
129 ‘physique athlete’ in the literature, synonyms were included in the  
130 search strategy. Various combinations of the following search terms  
131 were used, for the search: ‘physique athlete’ OR ‘fitness competitor’  
132 OR ‘bodybuilding’ OR ‘competitive body-builder’ OR ‘figure athlete’  
133 AND (contest or competition OR dieting OR dietary intake or nutrition  
134 OR macronutrient OR micronutrient OR training OR body  
135 composition OR peak week OR practices OR weight loss OR weight  
136 regain). Several other search terms associated with health outcomes  
137 included: ‘physique athlete’ OR ‘fitness competitor’ OR  
138 ‘bodybuilding’ OR ‘competitive body-builder’ OR ‘figure athlete’  
139 AND (energy availability OR menstrual cycle OR bone, OR eating OR  
140 body image). Any additional articles relevant to the scope of this  
141 narrative review were obtained through PubMed via the function  
142 “similar articles” or from the reference lists of the included studies.

143 Criteria for inclusion were: *i*) studies published in English language  
144 and in peer-reviewed articles within the past 30 years (*i.e.*, theses or  
145 conference abstracts were not eligible), *ii*) studies involving human  
146 participants, *iii*) studies with participants who were currently engaging  
147 in or had previously been engaged in physique competitions, across  
148 any category (*i.e.*, bikini fitness, wellness fitness, and figure), *iv*)  
149 studies using female participants, or studies using both female and  
150 male participants, and *v*) studies investigating at least one of the  
151 following: body composition, nutritional intake, micronutrients,

152 training strategies, psychology, menstrual cycle, hormonal markers,  
153 bone mineral density, energy availability, and weight  
154 loss/management practices). Exclusion criteria were studies that  
155 reported use of performance-enhancing drugs, and only male  
156 participants.

157

### 158 **Definition of the natural female physique athlete**

159 Benjamin and Joseph Weider established the first organisation which  
160 specialised solely in bodybuilding events, known as the IFBB (Vallet,  
161 2017). To date, the IFBB is one of the most influential amateur sports  
162 organisations in the bodybuilding sphere and is an official signatory of  
163 the World Anti-Doping Code where athletes participate in random  
164 drug testing programs, such as urinalysis and polygraph tests for  
165 prohibited substances (IFBB, 2014).

166 Although bodybuilding was traditionally a male dominated sport, the  
167 growth of female competitors has increased significantly in recent  
168 times (Spendlove et al., 2015). This increase in popularity is largely  
169 due to the introduction of new female-specific physique categories  
170 (*e.g.*, Fitness, Body Fitness and Bikini Fitness) since 1995 (Spendlove  
171 et al., 2015; Tajrobehkar, 2016). As these new categories allowed  
172 ‘smaller’ competitors to enter the sport, and reduced the emphasis on  
173 muscle mass, it has encouraged healthier practices, indirectly  
174 attracting more women from mainstream society than in previous  
175 decades (Tajrobehkar, 2016).

176 Female physique athletes are assessed on aesthetic appearance and  
177 posing ability whereby high lean body mass (LBM) and low fat mass

178 (FM) are key markers of performance (Kleiner et al., 1994).  
179 Competitions involve comparison rounds; wherein athletes are  
180 instructed to perform poses, and a final round; in which top ranked  
181 athletes perform an individual posing routine (Steele et al., 2018). The  
182 intricate scoring system assesses athlete features, such as symmetry,  
183 muscularity, size and presentation (*i.e.*, personal confidence, facial  
184 beauty, and skin condition) (Choi, 2003; Obel, 1996). Unlike other  
185 weight-restricted sports (*e.g.*, male bodybuilding, wrestling and  
186 boxing), in which weight categories are utilised, FP athletes are  
187 allocated to categories based on their subjective assessment of the  
188 amount of LBM and FM, and are then further sub-classified by height  
189 (Fry et al., 1991). At one end of the continuum (*i.e.*, bikini fitness),  
190 athletes typically have less LBM and higher FM, whilst at the other  
191 end (*i.e.*, physique), athletes are diametrically opposed with high LBM  
192 and a corresponding low FM (Fig.1).

193

194 **[Insert Figure 1 near here]**

195

### 196 **Body composition in competition**

197 Typically, an annual schedule for the physique athlete is divided into  
198 an off-season phase and a pre-competition phase (Hackett et al., 2013).  
199 Within the off-season phase, physique athletes manipulate resistance  
200 training variables including volume, intensity and frequency for the  
201 purpose of gaining LBM (Spendlove et al., 2015). This period can last  
202 years and is characterised by a positive energy balance, in conjunction  
203 with a high protein intake to stimulate muscle anabolism (Phillips,



204 2004; Campbell et al., 2018). In the pre-competition phase, the  
205 majority of athletes attempt to reduce FM and preserve LBM using a  
206 combination of rigorous resistance and aerobic training, while  
207 manipulating their nutritional intake to achieve a negative energy  
208 balance (Hackett et al., 2013; Petrizzo et al., 2017). The pre-  
209 competition phase lasts between 12 and 24 weeks (Mitchell et al.,  
210 2018) and athletes are likely to compete between two to three times  
211 per year (Chappell et al., 2018). Usually, the pre-competition phase is  
212 followed by a recovery phase (a transition to off-season), during which  
213 athletes increase total energy intake and decrease total training load  
214 (Hulmi et al., 2016). Previous research reports the magnitude of weight  
215 loss is in the range of 6-10 kg over a 18-24 week period (Table 1). This  
216 suggests that FP athletes pursue a gradual approach to weight loss (~  
217 0.4 kg per week), similar to male bodybuilding and physique athletes  
218 (~ 0.6-0.8 kg per week) (Chappell et al., 2018; Kistler et al., 2014;  
219 Robinson et al., 2015; Rossow et al., 2013). In the end stages of the  
220 pre-competition phase, FP athletes achieve 8.6 - 16% body fat (Hulmi  
221 et al., 2016; Rohrig et al., 2017; Tinsley et al., 2018; Trexler et al.,  
222 2017), which is exceptionally lower than the recommended values for  
223 female athletes (Sundgot-Borgen and Garthe, 2011).

224

### 225 **Strategies to manipulate body composition during competition** 226 **week**

227 Whilst FP athletes employ a gradual approach to fat loss, acute weight  
228 loss practices occur during the competition week (Helms et al., 2014).  
229 Peer-reviewed articles suggest fluid, salt, and carbohydrate  
230 manipulation is commonly practiced to reduce body water content in

231 order to enhance muscle definition on competition day (Mitchell et al.,  
232 2017; Shephard, 1994). Over a third of twenty-two FP athletes  
233 practiced water manipulations (36 %), whereas more than two-quarters  
234 practiced carbohydrate manipulations (77 %) (Chappell and Simper,  
235 2018). Water loading, followed by water restriction is allegedly used  
236 to modify renal hormones and encourage urination beyond the period  
237 of increased fluid intake, resulting in reduced body water (Helms et  
238 al., 2014; Mitchell et al., 2017). The physiological effects of water  
239 loading have only been investigated in male combat sport athletes with  
240 a purpose of making-weight (Crighton et al., 2016; Reale et al., 2018),  
241 as opposed to physique athletes trying to enhance their aesthetic  
242 appearance. The acute weight loss experienced early in competition  
243 week (~7-5 days prior to competition) is likely to be mediated by  
244 glycogen depletion prior to a carbohydrate loading protocol (Chappell  
245 and Simper, 2018). Female physique athletes reduce their  
246 carbohydrate intake from 4.1- 4.5 g·kg<sup>-1</sup>·d<sup>-1</sup> before entering the pre-  
247 competition phase, to 1.2 - 2.7 g·kg<sup>-1</sup>·d<sup>-1</sup> at the end stages of pre-  
248 competition phase (Halliday et al., 2016; Rohrig et al., 2017). In one  
249 case, daily carbohydrate intake was reduced to ~ 0.3 g·kg<sup>-1</sup>·d<sup>-1</sup>, three  
250 days prior to competition (Tinsley et al., 2018). From the available  
251 evidence, it appears that during the pre-competition phase, FP athletes  
252 fall considerably below the carbohydrate recommendations for  
253 moderate volume training (5-7 g·kg<sup>-1</sup>·d<sup>-1</sup>) (Manore, 2002). Addressing  
254 the distribution of carbohydrate intake throughout the day and in  
255 relation to training, could provide further insights into the strategies  
256 used to optimise body composition (Slater and Phillips, 2011).

257 Based on limited data, the efficacy and safety of competition week  
258 strategies in physique events are still unknown, but might be  
259 detrimental to athlete health (Chappell and Simper, 2018; Helms et al.,  
260 2014) by increasing the risks associated with hyponatremia and  
261 glycogen depletion (Slater and Phillips, 2011).

262

### 263 **Health implications for the female physique athlete**

264 Physique athletes typically reduce their total energy intake to induce  
265 gradual weight loss over a prolonged period of time, and progress  
266 towards acute weight loss methods, such as restrictive diets (energy  
267 availability [EA]  $< 30 \text{ kcal} \cdot \text{kg}^{-1} \text{ FFM} \cdot \text{d}^{-1}$ , where FFM = fat free mass),  
268 in the latter stages of the pre-competition phase (Sundgot-Borgen et  
269 al., 2013; Fagerberg et al., 2017). As such, FP athletes face major  
270 health-related challenges in an attempt to reach and maintain a lean  
271 body composition.

272

#### 273 *Reduced energy availability in female physique athletes*

274 Current literature on FP athletes has documented prolonged periods of  
275 LEA, specifically during the pre-competition phases. Halliday and  
276 colleagues (Halliday et al., 2016) showed that during a 20-week pre-  
277 competition phase, the estimated mean EA was categorised as low in  
278 the initial ( $27.9 \text{ kcal} \cdot \text{kg}^{-1} \text{ FFM} \cdot \text{d}^{-1}$ ) and latter ( $23.3 \text{ kcal} \cdot \text{kg}^{-1} \text{ FFM} \cdot \text{d}^{-1}$ )  
279 stages of the phase, respectively. In this study (Halliday et al., 2016),  
280 total energy intake and exercise energy expenditure were self-reported  
281 and reproductive function was not measured. Similarly, Tinsley et al.  
282 (2018) documented caloric intakes of between  $18.2$  and  $31.1 \text{ kcal} \cdot \text{kg}^{-1}$

283 FFM·d<sup>-1</sup> in a FP athlete (during two different pre-competition phases)  
284 indicating extreme caloric restriction (Manore, 2002). Although EA  
285 was not objectively quantified, the authors estimated that the athlete  
286 fell below the threshold of EA for the maintenance of normal  
287 physiological function based on total energy intake and body  
288 composition data. Self-report research designs are not uncommon in  
289 the literature on physique athletes and, as such should be interpreted  
290 with caution (Fagerberg, 2017). Therefore, EA data in FP athletes  
291 remains questionable considering the lack of sensitive and relevant  
292 screening tools (Heikura et al., 2018). Nonetheless, aforementioned  
293 studies highlight that FP athletes may induce sub-optimal EA and  
294 shows the importance for future studies on this topic to utilise more  
295 robust measures of total energy intake and exercise energy expenditure  
296 in order to accurately evaluate EA (Elliott-Sale et al., 2018; Fagerberg,  
297 2017).

298

### 299 *Nutrient deficiency*

300 Bodybuilding diets are traditionally characterised as restrictive and  
301 monotonous, as they often limit food variability (Kleiner et al., 1994).  
302 As a consequence, compromised micronutrient status is often observed  
303 in the pre-competition phase among FP athletes (Slater and Phillips,  
304 2011). Calcium, iron, zinc and sodium intakes have been shown to  
305 decrease significantly, to less than two-thirds (~ 67%) of the  
306 Recommended Daily Allowance (RDA) (Newton et al., 1993;  
307 Walberg-Rankin and Gwazdauskas, 1993) in the absence of dietary  
308 supplements during the pre-competition phase. These results may be  
309 attributed to restricted energy intake combined with the elimination of  
310 sodium and dairy products from the diet (Steen, 1991). Considering

311 that weight loss trends/dietary fads typically change over time, it is  
312 worth noting that the applicability of the aforementioned studies might  
313 be limited (Spendlove et al., 2015).

314 More recently, Ismaeel et al. (2017) showed that FP athletes who used  
315 extreme restrictive eating patterns consumed significantly less protein  
316 ( $123 \pm 23$  g *cf.*  $65 \pm 16$  g,  $p = 0.02$ ), sodium ( $4,060 \pm 397$  mg *cf.*  $2,636$   
317  $\pm 1,028$  mg,  $p = 0.03$ ), vitamin E ( $10 \pm 2$  mg *cf.*  $6 \pm 1$  mg,  $p = 0.03$ )  
318 and vitamin C ( $170 \pm 47$  mg *cf.*  $66 \pm 27$  mg,  $p = 0.02$ ) than athletes  
319 who permitted dietary flexibility (Ismaeel et al., 2017). These  
320 differences may be caused by the large variation in total energy intake  
321 ( $1,965 \pm 259$  kcal·d<sup>-1</sup> *cf.*  $1,455 \pm 541$  kcal·d<sup>-1</sup>) consumed by each  
322 group. While the study (Ismaeel et al., 2017) included dietary  
323 supplements in the micronutrient analysis, it did not specify whether  
324 individuals were in the pre-competition or off-season phase.  
325 Nevertheless, these results identify potential risks for deficiencies in  
326 essential nutrients for FP athletes, which could suppress the immune  
327 function and cause increased susceptibility to illnesses and infections,  
328 especially for those engaging in restrictive eating patterns (Sundgot-  
329 Borgen and Garthe, 2011). As the majority of studies assessing  
330 micronutrient status have also used self-report methods (Ismaeel et al.,  
331 2017; Kleiner et al., 1994; Newton et al., 1993; Walberg-Rankin and  
332 Gwazdauskas, 1993; Walberg and Johnston, 1991), it is prudent that  
333 future measures are clarified using biomarkers in blood or urine  
334 samples.

335

336

337 *Menstrual irregularities, endocrine effects and bone health in female*  
338 *physique athletes*

339 Many active women with LEA develop various forms of reproductive  
340 dysfunction, including oligomenorrhea, amenorrhea and luteal phase  
341 defects (Manore, 2002). Low energy availability causes alterations in  
342 the hypothalamic-pituitary-ovarian axis, namely diminished secretion  
343 of luteinizing hormone and follicle stimulating-hormone, which  
344 subsequently reduces oestrogen production. The final consequence is  
345 typically described as functional hypothalamic amenorrhea (West,  
346 1998). Previous research has shown that 82-86% of females (non-  
347 contraceptive users) who entered at least one physique competition  
348 were either oligomenorrheic or amenorrheic (Walberg-Rankin and  
349 Gwazdauskas, 1993; Walberg and Johnston, 1991). Similarly, case  
350 studies have also observed amenorrhea (Hulmi et al., 2016; Petrizzo et  
351 al., 2017; Rohrig et al., 2017), with some reporting delays in  
352 menstruation of up to 71 weeks post-competition (Halliday et al.,  
353 2016; Kleiner et al., 1994; Kleiner et al., 1990).

354 Changes to reproductive and metabolic hormones in FP athletes have  
355 been observed in the pre-competition phase, including decreases in  
356 oestradiol, testosterone, thyroid stimulating hormone, triiodothyronine  
357 (T3) and leptin (Table 1). These hormones were normalised within 4 -  
358 16 weeks post-competition, when supported by an increased intake of  
359 protein ( $\sim 2. \text{g} \cdot \text{kg}^{-1} \cdot \text{d}^{-1}$ ) and greater EA (Hulmi et al., 2016; Trexler et  
360 al., 2017) with the exception of serum T3 and testosterone (Hulmi et  
361 al., 2016), which were only partially recovered 12-16 weeks after  
362 competition. As such, the suppression of these key metabolic  
363 hormones persist further into the recovery phase, possibly due to the

364 effects of dropping below the EA threshold regardless of altered  
365 exercise regimen, as previously described by Loucks and Heath  
366 (1994). More longitudinal data is required on endocrine and metabolic  
367 function beyond the 16 weeks post-competition to better understand  
368 the time-course for full restoration.

369 Regular menstrual cycles are often used as a surrogate marker of long-  
370 term LEA; however, the use of hormonal contraceptives may  
371 obfuscate this relationship (Heikura et al., 2018). Hormonal  
372 contraceptives provide negative feedback to the hypothalamus and  
373 pituitary glands, leading to suppression of follicle stimulating-  
374 hormone, luteinizing hormone and gonadotropin-releasing hormone,  
375 and continuous down-regulation of endogenous oestrogen and  
376 progesterone (Elliott-Sale et al., 2013). Previous data in FP athletes  
377 have failed to investigate female sex hormones (*i.e.*, oestrogen and  
378 progesterone) (Trexler et al., 2017), did not include hormonal  
379 contraceptive users (Halliday et al., 2016; Rohrig et al., 2017; Tinsley  
380 et al., 2018) or grouped all oral contraceptive users together, making  
381 the interpretation difficult (Elliott-Sale et al., 2013). Considering the  
382 high prevalence of hormonal contraceptive use (Hulmi et al., 2016),  
383 there is great concern that FP athletes, who are experiencing chronic  
384 LEA, are going undetected, as hormonal contraceptive use maintains  
385 regular menstrual cycles. To this end, there is a need for studies to  
386 determine whether the FP athletes, who are using hormonal  
387 contraceptives, are at increased risk of endocrine dysfunction.

388 Although it is not unusual for bone mineral density to be compromised  
389 during calorie restriction and reduced body mass, it is possible that the  
390 minimal changes observed in bone mineral density ( $1.062\text{-}1.204\text{g}\cdot\text{cm}^3$ )

391 (Van der Ploeg et al., 2001; Hulmi et al., 2016; Petrizzo et al., 2017)  
392 is explained by the high-impact and weight-bearing activities  
393 performed in their training regimens (Zanker et al., 2004). As a result,  
394 this may have served to retain bone-mineral density compartment  
395 (Layne & Nelson, 1999).

396

### 397 *Weight cycling*

398 Female physique athletes often experience rapid weight gain following  
399 competitions (Andersen et al., 1995; Walberg-Rankin and  
400 Gwazdauskas, 1993) with one study reporting uncontrollable binge  
401 eating behaviour, reflecting a hyperphagic effect to intensive weight  
402 loss protocols (Trexler et al., 2017). This practice is commonly known  
403 as ‘weight cycling’ (*i.e.*, repeated cycles of weight loss and regain).  
404 Previous research has reported unfavorable metabolic parameters  
405 including a decline in resting metabolic rate (RMR) (reduced between  
406 155 and 226 kcal) (Rohrig et al., 2017; Tinsley et al., 2018) during pre-  
407 competition phase and weight regain of up to 8.6 kg at 4 weeks post-  
408 competition refeeding in females (Walberg-Rankin and Gwazdauskas,  
409 1993). The RMR suppression is possibly induced by the dietary  
410 restriction during weight loss resulting in alterations in leptin levels,  
411 thyroid status and sympathetic nervous system activity (Stiegler and  
412 Cunliffe, 2006). Conversely, recent case studies have shown that some  
413 FP athletes use a “reverse dieting” technique, in order to avoid those  
414 implications (Trexler et al., 2014). This strategy requires athletes to  
415 slowly increase their energy intake in an effort to limit any rapid  
416 increases in FM, and to prevent reductions in RMR (Trexler et al.,  
417 2014). However, the effort to “reverse” (*i.e.*, slowly increase) energy



418 intake requires considerable discipline to curb the increases in appetite  
419 sensations (Greenway, 2015), and therefore the authors speculate,  
420 whether such a strategy is achievable. Future research on “reverse  
421 dieting” technique in the recovery phase is warranted.

422

423 *Disordered eating /Eating Disorders behaviours*

424 Considering that appearance is a major criterion to judge performance  
425 of FP athletes, the increased risk of DE/ED in this population is  
426 perhaps unsurprising. Important risk and trigger factors of poor eating  
427 habits in FP athletes may include the focus on aesthetic appearance as  
428 the primary performance marker in competition (Sundgot-Borgen and  
429 Torstveit, 2004), the peer/media pressure which can elicit body  
430 dissatisfaction (Hausenblas et al., 2013) and the influences from  
431 coaches with inadequate nutrition knowledge (Sundgot-Borgen,  
432 1994). There is also evidence that FP athletes are particularly  
433 vulnerable to DE/ED and body image dissatisfaction because of the  
434 preoccupation with being muscular and lean (Devrim et al., 2018).

435 For example, a cross-sectional study by Walberg and Johnston (1991)  
436 compared 12 aspiring and retired FP athletes with 103 recreational  
437 weight-lifters on the Eating Disorder Inventory. Results revealed that  
438 FP athletes had significantly greater food obsessions (67%),  
439 uncontrolled urges to eat (58%) and felt more terrified of becoming fat  
440 (58%; all  $p < 0.05$ ). The use of laxatives, for weight loss, (17% *cf.*  
441 15%) and binge eating (50% *cf.* 62%) were similar between the groups.

442 In another study, Andersen et al. (1998) reported that ten out of twenty-  
443 six FP athletes experienced binge eating episodes in the recovery phase

444 and eighteen out of twenty-six FP athletes displayed body and weight  
445 dissatisfaction, reiterating that there is a high risk of eating and body  
446 image-related problems within the sport (Pope et al., 1997).  
447 Nevertheless, the small sample size and the lack of any comparative  
448 group analysis by Andersen et al. (1998) somewhat limits the  
449 interpretation. To the authors' knowledge, no quantitative data  
450 examining disordered eating behaviours exists for a large cohort of  
451 natural FP athletes.

452 Furthermore, it is difficult to capture sensitive data using questionnaire  
453 methods concerning mental health and well-being without a  
454 confirmatory interview (Andersen et al., 1998). Athletes may be  
455 anxious of revealing inappropriate eating practices in fear of being  
456 negatively judged, which could prevent honest disclosure.  
457 Nevertheless, there is a plausible link between participation in  
458 physique sports and DE behaviours. Further research is warranted to  
459 explore the psychopathological and behavioural outcomes in these  
460 athletes. Understanding the experiences and perceptions of weight  
461 management and eating behaviours across the pre-competition,  
462 recovery and off-season phases might be of particular importance.  
463 Using validated screening tools to detect DE and EDs and follow-up  
464 interviews will allow researchers to collect comprehensive data that  
465 could inform practice.

466

#### 467 **Conclusions and future research**

468 The ultimate determinant of competitive success in physique events is  
469 a high degree of muscularity and minimal FM. As such, FP athletes

470 engage in both prolonged energy restriction and intensive training  
471 regimens in order to meet these demands. Some FP athletes may be  
472 vulnerable to chronic LEA and associated physiological and  
473 psychological health effects, even during the recovery phase. Despite  
474 an increased participation in physique events, there is paucity in the  
475 literature on FP athletes. Future research should therefore:

- 476 *i)* identify the weight management strategies and DE/ED  
477 behaviours of FP athletes, in order to determine the risks  
478 of LEA in this population;
- 479 *ii)* explore such strategies using a qualitative approach, to  
480 enable FP athletes to express and elaborate on their  
481 experiences of weight management, eating behaviours  
482 and psycho-physiological health implications;
- 483 *iii)* investigate endocrine and micronutrient changes in FP  
484 athletes using objective biomarkers, to assess whether  
485 these individuals are in chronic states of LEA throughout  
486 the season;
- 487 *iv)* develop effective, safe and evidence-based nutritional  
488 recovery guidelines to minimise any long-term health  
489 implications.

490

#### 491 **Practical Application Statement**

492 At present, it is difficult to draw upon practical applications from the  
493 existing literature. FP athletes are an understudied population, and  
494 methodological limitations exist. A primary issue is that the majority  
495 of cited reports are case studies or observational studies with small

496 sample sizes, which may be insufficient for drawing definite  
497 conclusions on the possible physiological and psychological health  
498 implications among natural FP athletes. More research will have a  
499 valuable impact upon the advice and strategies provided by coaches  
500 and practitioners who work with these athletes.

501 It is worth noting that many female athletes are reluctant to discuss  
502 their weight management practices and health histories with sport  
503 science/health professionals (Manore, 2002), making this population  
504 difficult to research (Aspridis et al., 2014), and may explain the small  
505 sample sizes reported by previous studies (Halliday et al., 2016;  
506 Ismaeel et al., 2017; Petrizzo et al., 2017). Therefore, it is imperative  
507 that coaches and sport science/health professionals working with  
508 physique athletes build trusting relationships and respect their desires  
509 to be lean, with a view to achieve an optimum body composition and  
510 health outcomes, through a collaborative relationship.

511

#### 512 **Novelty statement**

513 This is the first review to summarise the common physiological and  
514 psychological health implications among female physique athletes.

515

#### 516 **Acknowledgements**

517 The authors would like to thank Dr Jim McVeigh, Director of Public  
518 Health at Liverpool John Moores University for his valuable input to  
519 this review.

520

**521 Author's contributions**

522 The lead author on this paper was NA. SM, KE-S, ID and KE  
523 participated in the drafting of the manuscript. All authors read and  
524 approved the final manuscript.

525

**526 Declaration of funding sources**

527 No funding was received for the preparation of this manuscript

528

**529 Conflicts of interest**

530 The authors declare that they have no conflicts of interest

531

532

533

534

535

536

537

538

539

540

541

542

543

544

545

546

547

548

549

550

551

552

553

554

555

556

557

558

559

560

561 **References**

562

563 Andersen, R. E., Barlett, S. J., Morgan, G. D., and Brownell, K. D.  
 564 (1995). Weight loss, psychological, and nutritional patterns in  
 565 competitive male body builders. *International Journal of*  
 566 *Eating Disorders*, 18(1), 49-57.

567

568 Andersen, R. E., Brownell, K. D., Morgan, G. D., and Bartlett, S. J.  
 569 (1998). Weight loss, psychological, and nutritional patterns in  
 570 competitive female bodybuilders. *Eating Disorders*, 6(2),  
 571 159-167.

572

573 Aspridis, A., O'Halloran, P., and Liamputtong, P. (2014). Female  
 574 Bodybuilding: Perceived Social and Psychological Effects of  
 575 Participating in the Figure Class. *Women in Sport and*  
 576 *Physical Activity Journal*, 22(1), 24-29.

577

578 Baghurst, T., Parish, A., and Denny, G. (2014). Why Women Become  
 579 Competitive Amateur Bodybuilders. *Women in Sport and*  
 580 *Physical Activity Journal*, 22(1), 5-9.

581

582

583 Campbell, B. I., Aguilar, D., Conlin, L., Vargas, A., Schoenfeld, B. J.,  
 584 Corson, A., and Couvillion, K. (2018). Effects of High Versus  
 585 Low Protein Intake on Body Composition and Maximal  
 586 Strength in Aspiring Female Physique Athletes Engaging in  
 587 an 8-Week Resistance Training Program. *International*  
 588 *Journal of Sport Nutrition and Exercise Metabolism*, 1-6.

589

590 IFBB Elite Pro Categories. (2017). Retrieved from  
 591 <https://eliteproifbb.com/our-disciplines/categories/>.

592

593 Chappell, A., and Simper, T. (2018). Nutritional Peak Week and  
 594 Competition Day Strategies of Competitive Natural  
 595 Bodybuilders. *Sports*, 6(4), 126.

596

597 Chappell, A. J., Simper, T., and Barker, M. E. (2018). Nutritional  
 598 strategies of high level natural bodybuilders during  
 599 competition preparation. *Journal of the International Society*  
 600 *of Sports Nutrition*, 15, 4.

601

602 Choi, P. Y. L. (2003). Muscle matters: maintaining visible differences  
 603 between women and men. *Sexualities, Evolution and Gender*,  
 604 5(2), 71-81.

605

606 Crighton, B., Close, G. L., and Morton, J. P. (2016). Alarming weight  
 607 cutting behaviours in mixed martial arts: a cause for concern  
 608 and a call for action. *British Journal of Sports Medicine*, 50(8),  
 609 446-447.

610

611 Devrim, A., Bilgic, P and Hongo, N. (2018). Is there any relationship  
 612 between body image perception, eating disorders, and muscle  
 613 dysmorphic disorders in male bodybuilders? *American Journal*  
 614 *of Men's Health*, 12(5), 1746-1758

615

- 616  
617 Elliott-Sale, K. J., Smith, S., Bacon, J., Clayton, D., McPhilimey, M.,  
618 Goutianos, G., and Sale, C. (2013). Examining the role of oral  
619 contraceptive users as an experimental and/or control group in  
620 athletic performance studies. *Contraception*, 88(3), 408-412.  
621
- 622 Elliott-Sale, K. J., Tenforde, A. S., Parziale, A. L., Holtzman, B., and  
623 Ackerman, K. E. (2018). Endocrine Effects of Relative Energy  
624 Deficiency in Sport. *International Journal of Sport Nutrition  
625 and Exercise Metabolism*, 28, 335-349.  
626
- 627 Fagerberg, P. (2017). Negative Consequences of Low Energy  
628 Availability in Natural Male Bodybuilding: A Review.  
629 *International Journal of Sport Nutrition and Exercise  
630 Metabolism*, 0(0), 1-18.  
631
- 632 Fry, A. C., Ryan, A. J., Schwab, R. J., Powell, D. R., and Kraemer, W.  
633 J. (1991). Anthropometric characteristics as discriminators of  
634 body-building success. *Journal of Sports Sciences*, 9(1), 23-  
635 32.  
636
- 637 Greenway, F. L. (2015). Physiological adaptations to weight loss and  
638 factors favouring weight regain. *International Journal Of  
639 Obesity*, 39, 1188.  
640
- 641 Hackett, D. A., Johnson, N. A., and Chow, C. M. (2013). Training  
642 practices and ergogenic aids used by male bodybuilders.  
643 *Journal of Strength and Conditioning Research*, 27(6), 1609-  
644 1617.  
645
- 646 Halliday, T. M., Loenneke, J. P., and Davy, B. M. (2016). Dietary  
647 Intake, Body Composition, and Menstrual Cycle Changes  
648 during Competition Preparation and Recovery in a Drug-Free  
649 Figure Competitor: A Case Study. *Nutrients*, 8(11), 740.  
650
- 651 Hausenblas, H. A., Campbell, A., Menzel, J. E., Doughty, J., Levin,  
652 M. and Thompson, J. K. (2013). Media effects of experimental  
653 presentation of the ideal physique on eating disorder  
654 symptoms: A meta-analysis of laboratory studies. *Clinical  
655 Psychology Review*, 33(1), 168-181.  
656
- 657 Heikura, I. A., Uusitalo, A. L. T., Stellingwerff, T., Bergland, D.,  
658 Mero, A. A., and Burke, L. M. (2018). Low Energy  
659 Availability Is Difficult to Assess but Outcomes Have Large  
660 Impact on Bone Injury Rates in Elite Distance Athletes.  
661 *International Journal of Sport Nutrition and Exercise  
662 Metabolism*, 28(4), 403-411.  
663
- 664 Helms, E. R., Aragon, A.A., Fitschen PJ. (2014). Evidence-based  
665 recommendations for natural bodybuilding contest  
666 preparation: nutrition and supplementation. *Journal of the  
667 International Society of Sports Nutrition*, 12(11), 20.  
668
- 669 Hulmi, J. J., Isola, V., Suonpaa, M., Jarvinen, N. J., Kokkonen, M.,  
670 Wennerstrom, A., and Hakkinen, K. (2016). The Effects of

- 671 Intensive Weight Reduction on Body Composition and Serum  
672 Hormones in Female Fitness Competitors. *Frontiers in*  
673 *Physiology*, 7, 689.  
674
- 675 IFBB. (2014). IFBB anti-doping area. Retrieved from  
676 <https://ifbb.com/wada-anti-doping-rules/>.  
677
- 678 Ismaeel, A., Weems, S., and Willoughby, D. (2017). A Comparison of  
679 the Nutrient Intakes of Macronutrient-based Dieting and Strict  
680 Dieting Bodybuilders. *International Journal of Sport*  
681 *Nutrition and Exercise Metabolism*, 28(4), 502-508.  
682
- 683
- 684 Kistler, B. M., Fitschen, P. J., Ranadive, S. M., Fernhall, B., and  
685 Wilund, K. R. (2014). Case study: Natural bodybuilding  
686 contest preparation. *International Journal of Sport Nutrition*  
687 *and Exercise Metabolism*, 24(6), 694-700.  
688
- 689 Kleiner, S., Bazzarre, TL., and Ainsworth, B. E. (1994). Nutritional  
690 status of nationally ranked elite bodybuilders. *International*  
691 *Journal of Sport Nutrition.*, 4(1), 54-69.  
692
- 693 Kleiner, S. M., Bazzarre, T. L., and Litchford, M. D. (1990). Metabolic  
694 profiles, diet, and health practices of championship male and  
695 female bodybuilders. *Journal of the American Dietetic*  
696 *Association*, 90(7), 962-967.  
697
- 698 Layne, J. E., and Nelson, M. E. (1999). The effects of progressive  
699 resistance training on bone density: a review. *Medicine and*  
700 *Science in Sports and Exercise*, 31(1), 25-30.  
701
- 702 Loucks, A. B., and Heath, E. M. (1994). Induction of low-T3  
703 syndrome in exercising women occurs at a threshold of energy  
704 availability. *American Journal of Physiology*, 266(3 Pt 2),  
705 817-823.  
706
- 707 Loucks, A. B., Kiens, B., and Wright, H. H. (2011). Energy availability  
708 in athletes. *Journal of Sports Sciences*,  
709 29 Suppl 1, S7-15.  
710
- 711 Manore, M. M. (2002). Dietary recommendations and athletic  
712 menstrual dysfunction. *Sports Medicine*, 32(14), 887-901.  
713
- 714 Mitchell, L., Hackett, D., Gifford, J., Estermann, F., and O'Connor, H.  
715 (2017). Do Bodybuilders Use Evidence-Based Nutrition  
716 Strategies to Manipulate Physique? *Sports*, 5(4), 76.  
717
- 718 Mitchell, L., Slater, G., Hackett, D., Johnson, N., and O'Connor, H.  
719 (2018). Physiological implications of preparing for a natural  
720 male bodybuilding competition. *European Journal of Sport*  
721 *Sciences*, 18(5), 619-629.  
722
- 723 Mountjoy, M., Sundgot-Borgen, J., Burke, L., Carter, S., Constantini,  
724 N., Lebrun, C., and Ljungqvist, A. (2014). The IOC consensus  
725 statement: beyond the Female Athlete Triad—Relative



- 726 Energy Deficiency in Sport (RED-S). *British Journal of*  
727 *Sports Medicine*, 48(7), 491-497.
- 728
- 729 Newton, L. E., Hunter, G., Bammon, M., and Roney, R. (1993).  
730 Changes in Psychological State and Self-Reported Diet  
731 During Various Phases of Training in Competitive  
732 Bodybuilders. *The Journal of Strength and Conditioning*  
733 *Research*, 7(3), 153-158.
- 734
- 735 Obel, C. (1996). Collapsing Gender in Competitive Bodybuilding:  
736 Researching Contradictions and Ambiguity in Sport.  
737 *International Review for the Sociology of Sport*, 31(2), 185-  
738 202.
- 739
- 740 Parish, T., Bahgurst, T., and Turner, R. (2010). Becoming competitive  
741 amateur bodybuilders: Identification of contributors.  
742 *Psychology of Men and Masculinity*, 11(2), 152-159.
- 743
- 744 Petrizzo, J., DiMenna, F. J., Martins, K., Wygand, J., and Otto, R. M.  
745 (2017). Case Study: The Effect of 32 Weeks of Figure-Contest  
746 Preparation on a Self-Proclaimed Drug-free Female's Lean  
747 Body and Bone Mass. *International Journal of Sport Nutrition*  
748 *and Exercise Metabolism*, 27(6), 543-549.
- 749
- 750 Phillips, S. M. (2004). Protein requirements and supplementation in  
751 strength sports. *Nutrition*, 20(7), 689-695.
- 752
- 753
- 754 Pope, H. G., Gruber, A. J., Choi, P., Olivardia, R., and Phillips, K. A.  
755 (1997). Muscle Dysmorphia: An Underrecognized Form of  
756 Body Dysmorphic Disorder. *Psychosomatics*, 38(6), 548-557.
- 757
- 758
- 759 Probert, A., Palmer, F., and Leberman, S. (2007). The Fine Line: An  
760 insight into 'risky' practices of male and female competitive  
761 bodybuilders. *Annals of Leisure Research*, 10(3-4), 272-290.
- 762
- 763 Reale, R., Slater, G., Cox, G. R., Dunican, I. C., and Burke, L. M.  
764 (2018). The Effect of Water Loading on Acute Weight Loss  
765 Following Fluid Restriction in Combat Sports Athletes.  
766 *International Journal of Sport Nutrition and Exercise*  
767 *Metabolism*, 28(6), 565-573.
- 768
- 769 Robinson, S. L., Lambeth-Mansell, A., Gillibrand, G., Smith-Ryan,  
770 A., and Bannock, L. (2015). A nutrition and conditioning  
771 intervention for natural bodybuilding contest preparation:  
772 Case study. *Journal of the International Society of Sports*  
773 *Nutrition*, 12(1), 20.
- 774
- 775 Rohrig, B. J., Pettitt, R.W., Pettitt, C.D., and Kanzenback, T.L. (2017).  
776 Psychophysiological tracking of a female physique competitor  
777 through competition preparation. *International Journal of*  
778 *Exercise Science*, 10(2), 301-311.
- 779

- 780 Rossow, L. M., Fukuda, D. H., Fahs, C. A., Loenneke, J. P., and Stout,  
781 J. R. (2013). Natural bodybuilding competition preparation  
782 and recovery: a 12-month case study. *International Journal of*  
783 *Sports Physiology and Performance*, 8(5), 582-592.  
784
- 785 Rowbottom, M. (2017, December 1). IFBB competitors place their  
786 bets for World Fitness Championships. Retrieved from  
787 [https://www.insidethegames.biz/articles/1058593/ifbb-](https://www.insidethegames.biz/articles/1058593/ifbb-competitors-place-their-bets-for-world-fitness-championships)  
788 [competitors-place-their-bets-for-world-fitness-](https://www.insidethegames.biz/articles/1058593/ifbb-competitors-place-their-bets-for-world-fitness-championships)  
789 [championships](https://www.insidethegames.biz/articles/1058593/ifbb-competitors-place-their-bets-for-world-fitness-championships).  
790
- 791 Shephard, R. J. (1994). Electrolyte manipulation in female body-  
792 builders. *British Journal of Sports Medicine*, 28(1), 60-61.  
793
- 794 Slater, G., and Phillips, S. M. (2011). Nutrition guidelines for strength  
795 sports: sprinting, weightlifting, throwing events, and  
796 bodybuilding. *Journal of Sports Sciences*, 29 Suppl 1, S67-77.  
797
- 798 Steen, S. N. (1991). Precontest strategies of a male bodybuilder.  
799 *International Journal of Sport Nutrition*, 1(1), 69-78.  
800
- 801 Spendlove, J., Mitchell, L., Gifford, J., Hackett, D., Slater, G., Cobley,  
802 S., and O'Connor, H. (2015). Dietary Intake of Competitive  
803 Bodybuilders. *Sports Medicine*, 45(7), 1041-1063.  
804
- 805 Steele, I., Pope Jr, H., and Kanayama, G. Weightlifting. (2018). In I.D.  
806 Glick, S. Todd., and D. Kamis (Ed.), *The ISSP Manual of*  
807 *Sports Psychiatry* (pp. 91-116). New York: Routledge.  
808
- 809 Stiegler, P., and Cunliffe, A. (2006). The Role of Diet and Exercise for  
810 the Maintenance of Fat-Free Mass and Resting Metabolic Rate  
811 During Weight Loss. *Sports Medicine*, 36(3), 239-262.  
812
- 813 Sundgot-Borgen, J. (1994). Risk and trigger factors for the  
814 development of eating disorders in female elite athletes.  
815 *Medicine and Science in Sports and Exercise*, 26(4), 414-419.  
816
- 817 Sundgot-Borgen, J., and Garthe, I. (2011). Elite athletes in aesthetic  
818 and Olympic weight-class sports and the challenge of body  
819 weight and body compositions. *Journal of Sports Sciences*, 29  
820 *Suppl 1*, 101-114.  
821
- 822 Sundgot-Borgen, J., Meyer, N. L., Lohman, T. G., Ackland, T. R.,  
823 Maughan, R. J., Stewart, A. D., and Muller, W. (2013). How  
824 to minimise the health risks to athletes who compete in  
825 weight-sensitive sports review and position statement on  
826 behalf of the Ad Hoc Research Working Group on Body  
827 Composition, Health and Performance, under the auspices of  
828 the IOC Medical Commission. *British Journal of Sports*  
829 *Medicine*, 47(16), 1012-1022.  
830
- 831 Sundgot-Borgen, J., and Torstveit, M. K. (2004). Prevalence of eating  
832 disorders in elite athletes is higher than in the general  
833 population. *Clinical Journal of Sport Medicine*, 14(1), 25-32  
834

- 835 Tajrobehkar, B. (2016). Flirting With the Judges: Bikini Fitness  
836 Competitors' Negotiations of Femininity in Bodybuilding  
837 Competitions. *Sociology of Sport Journal*, 33(4), 294-304.  
838
- 839 Tinsley, G. M., Trexler, E. T., Smith-Ryan, A. E., Paoli, A., Graybeal,  
840 A. J., Campbell, B. I., and Schoenfeld, B. J. (2018). Changes  
841 in Body Composition and Neuromuscular Performance  
842 Through Preparation, 2 Competitions, and a Recovery Period  
843 in an Experienced Female Physique Athlete. *The Journal of*  
844 *Strength and Conditioning Research, Publish Ahead of Print.*  
845
- 846 Trexler, E. T., Hirsch, K. R., Campbell, B. I., and Smith-Ryan, A. E.  
847 (2017). Physiological Changes Following Competition in  
848 Male and Female Physique Athletes: A Pilot Study.  
849 *International Journal of Sport Nutrition and Exercise*  
850 *Metabolism*, 27(5), 458-466.  
851
- 852 Trexler, E. T., Smith-Ryan, A. E., and Norton, L. E. (2014). Metabolic  
853 adaptation to weight loss: implications for the athlete. *Journal*  
854 *of the International Society of Sports Nutrition*, 11(1), 7.  
855
- 856 Van der Ploeg, G. E., Brooks, A. G., Withers, R. T., Dollman, J.,  
857 Leaney, F., and Chatterton, B. E. (2001). Body composition  
858 changes in female bodybuilders during preparation for  
859 competition. *European Journal of Clinical Nutrition*, 55(4),  
860 268-277.  
861
- 862 Vallet, G. (2017). The gendered economics of bodybuilding.  
863 *International Review of Sociology*, 27(3), 525-545.  
864
- 865 Walberg-Rankin, J. E. C., and Gwazdauskas, F. C. (1993). Diet and  
866 weight changes of female bodybuilders before and after  
867 competition. *International Journal of Sport Nutrition.*, 3(1),  
868 87-102.  
869
- 870 Walberg, J. L., and Johnston, C. S. (1991). Menstrual function and  
871 eating behavior in female recreational weight lifters and  
872 competitive body builders. *Medicine and Science in Sports*  
873 *and Exercise*, 23(1), 30-36.  
874
- 875 West, R. V. (1998). The Female Athlete. *Sports Medicine*, 26(2), 63-  
876 71.  
877
- 878 Zanker, C. L., Osborne, C. Cooke, C. B., Oldroyd, B., and Trustcott,  
879 J. G. (2004). Bone density, body composition and menstrual  
880 history of sedentary female former gymnastics, aged 20-32  
881 years *Osteoporosis International*, 15(2), 145-154.  
882  
883  
884  
885  
886  
887

**Table 1:** Overview of the recent studies of reproductive health of female physique athletes.

Study	N	Body weight change (Body Fat %)		Time period (weeks)	TEST		E <sub>2</sub>	T <sub>3</sub>	T <sub>4</sub>	CORT	Ghrelin	LP	TSH	IN	Method for menstrual status	Absence of menstruation	Bone mass density (DXA)							
		C P	R C		C P	R C												C P	R C	C P	R C	C P	R C	C P
Haliday et al. 2016	1 ♀	-8.3kg; (15.1- 8.6%)	+5.2kg; (8.6- 14.8%)	20 CP; 20 RC	-	-	-	-	-	-	-	-	-	-	-	-	-	Self-report	9 weeks pre- and up to 71 weeks post- competition	NA				
Hulmi et al. 2016	27 ♀	-7.8kg (23.1- 12.7%)	+6.1kg (12.7- 20.1%)	20 CP; 17.5 RC	↓	(↑)	↓	↑	↓	(↑)	↓	↑	-	-	-	↓	↑	↓	↑	-	-	Serum and self-report	11.5% pre- competition and 28% post- competition	↓CP; ↑RC
Trexler al. 2017	8 ♀ 7 ♂	-	+3.9kg (12.5- 14.9%)	4-6 RC	-	↑	-	-	-	↑↓	↑	↓	-	-	-	↑	-	Saliva	-	-				
Petrizzo et al. 2017	1 ♀	-7.7kg (24.4- 11.3%)	-	24 CP	-	-	-	-	-	-	-	-	-	-	-	-	-	Self-report	Oligomenorrhea	No change				
Rohrig et al. 2017	1 ♀	-10.1kg (30.5- 15.9%)	-	24 CP	↑↓	-	↑↓	-	-	-	-	↑↓	-	-	↓	-	↓	-	-	-	Serum and self-report	8 weeks pre- competition	-	
Tinsley et al., 2018	1 ♀	-6 kg (20.3- 11.6%)	+6.8kg (11.6- 18.8%)	18 CP (1) 7 CP (2) 9 RC	-	-	-	-	-	-	-	-	-	-	-	-	-	Self-report	12 weeks pre- competition (1) and up to 12 weeks post- competition (2)	NA				

♀ indicates female physique athletes, ♂ indicates male physique athletes, ↑↓ indicates fluctuation, CP indicates the pre-competition phase, RC indicates recovery phase, ( ) indicates not recovered to initial baseline values, (1) indicates 1<sup>st</sup> competition and (2) indicates a 2<sup>nd</sup> competition. TEST = Testosterone, E2 = Estradiol, T<sub>3</sub> = Triiodothyronine, T<sub>4</sub> = Thyroxine, CORT = Cortisol; TSH= Thyroid stimulating hormone; LP= Leptin, IN = Insulin; DXA = Dual-energy X-ray absorptiometry. NA = Information not available.

**Figure 1:** An overview of the current female categories in women's physique competitions. The categories are progressive steps along a continuum between lean body mass and fat mass. 'Dry' refers to dehydration and the subsequent reduction in body water (Chappell et al., 2018). The number of height classes in each category is determined by the popularity of the single category. This figure was drawn using information retrieved from the International Federation of Bodybuilding and Fitness website (FBB Elite Pro Categories, 2017).

