

High glucose load and endotoxemia among overweight and obese Arab women with and without diabetes

An observational study

Dara Al-Disi, PhD^a, Mohammed Ghouse Ahmed Ansari, MSc^b, Shaun Sabico, MD, PhD^b, Kaiser Wani, MSc^b, Syed Danish Hussain, MSc^b, Mona M. Elshafie^a, Philip McTernan, PhD^c, Nasser M. Al-Daghri, PhD^{b,*}

Abstract

Dietary intake influences gut microbiota activity. Nevertheless, there is a lack of evidence available that illustrates the acute effects of high glucose meal on metabolic endotoxemia. The present study assessed the acute impact of high glucose meal on endotoxemia and other clinical parameters in Saudi females with varying degrees of glycemia.

The subjects were 64 consenting pre-menopausal women, grouped into 3: control [n = 14 lean, non-T2DM, BMI = 22.2 ± 2.2 kg/m²]; overweight [n = 16, non-T2DM, BMI = 28.5 ± 1.5 kg/m²] and T2DM [n = 34, BMI = 35.2 ± 7.7 kg/m²]. After an overnight fast, all subjects were given a standardized high-glucose (75 g) meal. Anthropometrics were taken and blood samples were withdrawn at baseline and postprandial (0, 2 and 4-hours), serum glucose, endotoxin and lipid profile were quantified.

At baseline, total cholesterol, LDL-cholesterol, triglycerides and serum glucose levels were significantly higher (*P* values < .01) whereas significantly lower HDL-cholesterol levels (*P* < .01) were observed in T2DM subjects compared to other groups. Baseline endotoxin levels were highest in the overweight group (3.2 ± 1.1 mmol/L) as compared to control (2.0 ± 0.5 mmol/L) and T2DM (2.7 ± 1.2 mmol/L) (*P* = .046). HDL-cholesterol, LDL-cholesterol and triglycerides, significantly decreased in the T2DM group after 2 hours (*P* values < .05), whereas unremarkable changes observed in other groups. Lastly, endotoxin levels significantly increased only in the overweight group (3.2 ± 1.1 vs 4.2 ± 1.4 mmol/L; *P* < .05), 4 hours postprandial.

High glucose meal elevates endotoxemia only among overweight subjects and impairs dysbiosis.

Abbreviations: BMI = body mass index, CBCD = chair for biomarkers of chronic diseases, GLM = general linear model, HDL = high-density lipoprotein cholesterol, KSA = Kingdom of Saudi Arabia, LAL = limulus amoebocyte lysate, LDL = low-density lipoprotein cholesterol, LPS = lipopolysaccharides, PCC = primary care center, SPSS = statistical package for the social sciences, T2DM = type 2 diabetes mellitus, USA = United States of America, WHR = waist-hip ratio.

Keywords: Arab women, endotoxin, high sugar meal, metabolic endotoxemia

1. Introduction

The prevalence of obesity and diabetes across the globe has become the most devastating health impediments that have progressively increased over a decade.^[1–3] Numerous studies have endorsed that genetics, lifestyle, dietary habits and environmental factors play a major role in their pathogene-

sis.^[4–6] In recent years, mounting evidence has highlighted that the fluctuations in the gut microbiota (dysbiosis) cause an exponential increase in bacterial endotoxin [lipopolysaccharides (LPS)] levels, which potentiate some inflammatory markers thought to underpin a number of common metabolic diseases.^[7–10] Endotoxin translocates from the “leaky” gut lumen to the blood, leading to a condition termed “metabolic endotoxe-

Editor: Gaurav Malhotra.

This research was funded by the Research Supporting Project (RSP-2020/21), King Saud University, Riyadh, Saudi Arabia. The funder had no role in the design of the study; in the collection, analyses, or interpretation of data; in the writing of the manuscript, or in the decision to publish the results.

The authors declare no conflict of interest.

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

^a Department of Community Health Sciences, College of Applied Medical Sciences, King Saud University, ^b Riyadh Biochemistry Department, College of Science, King Saud University, Riyadh, Saudi Arabia, ^c School of Science and Technology, Department of Biosciences, Nottingham Trent University, Nottingham, NG1 8NS, UK.

* Correspondence: Nasser M. Al-Daghri, Biochemistry Department, College of Science, King Saud University, Riyadh 11451, Kingdom of Saudi Arabia (e-mail: ndaghri@ksu.edu.sa).

Copyright © 2020 the Author(s). Published by Wolters Kluwer Health, Inc.

This is an open access article distributed under the Creative Commons Attribution License 4.0 (CCBY), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

How to cite this article: Al-Disi D, Ansari MG, Sabico S, Wani K, Hussain SD, Elshafie MM, McTernan P, Al-Daghri NM. High glucose load and endotoxemia among overweight and obese Arab women with and without diabetes: an observational study. *Medicine* 2020;99:46(e23211).

Received: 12 March 2020 / Received in final form: 8 October 2020 / Accepted: 19 October 2020

<http://dx.doi.org/10.1097/MD.00000000000023211>

mia,^[11] and elevated circulatory endotoxin levels are associated with an array of chronic non-communicable diseases like type 2 diabetes mellitus (T2DM),^[12] obesity,^[13,14] atherosclerosis^[15,16] and liver disease.^[17] Concurrently, several studies have examined the impact of pre-/probiotics on gut microbiota or endotoxins levels and related metabolic diseases.^[7,18,19] Furthermore, biomedical industries have shown an intensive commercial interest in the role of the human microbiome and factors alleviating it, by using pre-/pro-biotics.^[20,21]

It is well known that diet and feeding habits appear to play a predominant role in modulating endotoxin levels over the course of the day.^[22,23] Our previous study pointed out that metabolic endotoxemia facilitated by high-fat meal has been exacerbated in subjects with T2DM, accompanied by a parallel rise in cardio-metabolic risk profile compared to their healthy counterparts.^[24] This cardiometabolic profile includes lipids such as triglycerides, LDL- and HDL-cholesterol, levels of which are significantly linked to endotoxin-induced sub-chronic inflammation observed in patients with metabolic diseases.^[24] To the best of our knowledge, the effect of high-glucose feed on the metabolic endotoxemia, however, remains to be fully understood. Hence, the current study sought to determine the postprandial effects of a high-glucose diet on endotoxemia and other serological parameters in adult Saudi T2DM, overweight females with their healthy counterparts.

2. Materials and methods

2.1. Participant and study design

In this observational study, 64 Saudi pre-menopausal women were randomly recruited from different primary care centres (PCCs) in Riyadh, Saudi Arabia. A research team member was deployed to each PCC with full knowledge of the protocol to be shared with the assigned PCC physician and nurse. The study has been approved by the Ethics Committee of the College of Medicine King Saud University (No. 10–173), Riyadh, Saudi Arabia, prior to the commencement of the research and all patients gave written consent.

2.2. Exclusion criteria

Subjects with chronic conditions such as asthma, hypertension, history of cardiac, kidney or liver disease, in addition, subjects with known long-standing diabetes and/or receiving anti-diabetic medication, those with fasting glucose levels >11 mmol/L, or with fasting triglycerides levels >4 mmol/L were excluded from the study.

2.3. Anthropometry and blood collections

Subjects were requested to visit their respective PCCs after an overnight fast (≥ 10 hours). Fasting blood samples were collected at baseline to determine subjects eligibility for the study and to assess glucose control as well as lipid profiles. Weight (kg) and height (cm) were recorded using an international standard scale (Digital Pearson Scale, ADAM Equipment Inc., (USA)). Waist circumference was measured at the level of the iliac crest at the end of normal respiration, and hip circumference was measured at the widest circumference around the buttocks using a measuring tape and body mass index (BMI) was calculated as kg/m^2 , as well as waist-to-hip ratios (WHR) were calculated. Blood pressure (mm Hg) was measured twice using a calibrated, mercurial sphygmomanometer, with a 15-minute interval. The

mean of the 2 readings was recorded. Based on BMI and DM status, consenting participants were categorized into 3 groups: Control (14 lean, non- T2DM) subjects with body mass index (BMI) $22.1 \pm 2.4 \text{ kg/m}^2$, range 7.8; overweight ($n=16$) with $28.5 \pm 1.5 \text{ kg/m}^2$, range 3.6 and subjects with early onset of T2DM (BMI: $35.1 \pm 9.0 \text{ kg/m}^2$; range 35.2, $n=34$).

2.4. High oral glucose load

Subjects were asked to return the following day, again in a fasted state, to their respective PCC for intervention. Subjects were given a high-glucose meal similar to oral glucose tolerance test (75 g of sugar dissolved in water). They were instructed to fully consume the liquid provided immediately following baseline blood extraction. Blood samples were drawn via peripheral venous can-nula and a flushing line was added to maintain vein patency since blood was collected serially at baseline (0 hours), after 2 hours and 4 hours. Blood samples were collected and transferred to a non-heparinized tube for centrifugation. Collected serum samples were transferred to pre-labelled plain tubes, placed on ice and delivered to the chair for Biomarkers of Chronic Diseases (CBCD) laboratory in King Saud University, Riyadh, KSA, for storage at -20°C . All participants were given tokens of appreciation for their participation in the study.

2.5. Assessment of biochemical profile

Serum glucose and lipid profile were measured routinely using an autoanalyzer (Konelab, Espoo, Finland). Serum endotoxin was analyzed using a commercially available QCL-1000 LAL End Point.

2.6. Data analysis

All statistical analyses were conducted using SPSS version 22.0 (SPSS, Chicago, IL, USA). All continuous variables were presented as mean \pm standard deviation and were normalized prior to parametric analyses. For comparison between groups, the univariate general linear model (GLM) was used with Bonferroni post-hoc comparisons. Age and BMI were used as covariates. For comparison between pre- and post-intervention, paired *T*-test was used for normally distributed variables and Wilcoxon Signed Ranks test for endotoxin. Bivariate linear regression analysis was used to determine associations between endotoxin variables of interest. Significance was set at $P < .05$.

3. Results

3.1. Baseline characteristics

Table 1 shows the baseline characteristics of the 3 groups. The T2DM subjects had significantly higher cholesterol level ($5.1 \pm 1.2 \text{ mmol/L}$), which was significantly higher from the cholesterol level of control ($3.8 \pm 0.4 \text{ mmol/L}$) subjects. The concentration of LDL cholesterol was also significantly higher in the T2DM subjects ($3.4 \pm 1.1 \text{ mmol/L}$) as compared to Overweight ($2.6 \pm 0.8 \text{ mmol/L}$, P value $< .01$) and control ($2.2 \pm 0.3 \text{ mmol/L}$, P value $< .01$) subjects. However, the concentration of HDL-cholesterol was significantly lower in T2DM ($1.0 \pm 0.2 \text{ mmol/L}$, P value $< .01$) and overweight subjects ($1.2 \pm 0.3 \text{ mmol/L}$, P value $< .01$) as compared to healthy ($1.3 \pm 0.3 \text{ mmol/L}$) subjects.

As expected, subjects with T2DM had significantly higher serum glucose levels ($7.3 \pm 1.8 \text{ mmol/L}$) than the overweight+

Table 1
Clinical characteristics of subjects according to the groups.

Parameters	Control	Overweight	T2DM	P value
N	14	16	34	
Age (years)	23.2±8.4	33.7±7.6 ^A	40.5±5.9 ^{AB}	<.001
BMI (kg/m ²)	22.1±2.4	28.9±1.3 ^A	35.1±9.0 ^{AB}	<.001
Glucose (mmol/L)*	4.4±0.3	4.5±0.5	7.3±1.8 ^{AB}	<.001
Total Cholesterol (mmol/L)	3.8±0.4	4.3±1.0	5.1±1.2 ^A	<.001
Triglycerides (mmol/L)*	0.7±0.2	1.1±0.7	1.6±0.6 ^{AB}	<.001
HDL-Cholesterol (mmol/L)	1.3±0.3	1.2±0.3	1.0±0.2 ^{AB}	<.001
LDL-Cholesterol (mmol/L)	2.2±0.3	2.6±0.8	3.4±1.1 ^{AB}	<.001
Endotoxin (mmol/L)*	2.0±0.5	3.2±1.1 ^A	2.7±1.2	.046

Data presented in terms of Mean±SD.

* denotes non-normal variables, and Superscript A and B indicate a significant difference at 0.05 level from Normal and Overweight respectively.

(4.5±0.5 mmol/L, $P < .01$) and control subjects (4.4±0.3 mmol/L, $P < .001$). The subjects with T2DM also had significantly higher serum triglycerides (1.6±0.6 mmol/L). Furthermore, endotoxin levels of the overweight subjects (3.2±1.1) were significantly higher than control subjects (2.0±0.5). However, the endotoxin level of T2DM subjects (2.7±1.2) was not significantly different from either control or overweight subjects.

3.2. Effects of high glucose load

Table 2 shows the changes in glucose, lipid and endotoxin profiles of all groups after adjusting for age and metformin. The glucose response to a glucose load was expected to increase in all groups after 2 hours with a concomitant decrease after 4 hours. Among the lipids, there were no significant changes observed in either the control or overweight groups. In the

Table 2
Metabolic changes pre- and post-prandial high-glucose load.

Group	0 – Hour	2 – Hour	4 – Hour
Total Cholesterol (mmol/L)			
Control	3.8±0.4	3.9±0.6	3.8±0.5
Overweight	4.3±1.0	4.3±1.0	4.4±1.1
T2DM	5.1±1.2	4.9±1.1	5.1±1.2
Glucose (mmol/L)*			
Control	4.4±0.3	6.0±1.4	4.7±1.5
Overweight	4.5±0.5	5.8±2.1	4.6±2.9
T2DM	7.3±1.8	13.4±4.5 ^A	7.6±4.3 ^B
High-density lipoprotein cholesterol (mmol/L)			
Control	1.3±0.3	1.3±0.3	1.3±0.2
Overweight	1.2±0.3	1.2±0.3	1.2±0.3
T2DM	0.97±0.18	0.95±0.18	1.01±0.21 ^{AB}
Low-density lipoprotein cholesterol (mmol/L)			
Control	2.2±0.3	2.2±0.3	2.2±0.3
Overweight	2.6±0.8	2.6±0.8	2.7±0.9
T2DM	3.4±1.1	3.2±0.9	3.3±1.0
Triglycerides (mmol/L)*			
Control	0.7±0.2	0.7±0.4	0.7±0.3
Overweight	1.1±0.7	1.1±0.7	1.1±0.7
T2DM	1.57±0.60	1.49±0.61	1.64±0.71 ^B
Endotoxin (EU/ml)*			
Control	2.0±0.5	2.1±0.4	1.6±0.7
Overweight	3.2±1.1	4.0±1.0 ^A	4.2±1.4 ^A
T2DM	2.7±1.2	2.5±1.3	2.9±1.6

Data presented in terms of Mean±SD, Superscript A and B indicate a significant difference from 0-hours and 2-hours.

* denotes non-normal variables; P value significant at <.05.

T2DM group, no significant differences in total cholesterol and LDL-cholesterol levels were observed. However, HDL-cholesterol significantly increased in the T2DM group at 4 hours as compared to baseline ($P < .05$) and 2-hours levels ($P < .05$). Furthermore, triglycerides concentration also increased significantly at 4-hours as compared to 2-hours level ($P < .05$). Finally, endotoxin levels significantly increased in the overweight group at 2-hours and 4-hours as compared to baseline endotoxin after adjusting for age and use of metformin ($P < .05$). No significant changes in endotoxin levels were observed in both the control and T2DM group.

3.3. Associations of endotoxins over measured variables over time

Table 3 shows the bivariate associations of endotoxin to glucose and lipids over time. Endotoxin was negatively associated with LDL-cholesterol only in the overall group and this was observed at 2-hour ($R = -0.37$; $P < .05$) (Fig. 1). The rest of the associations were non-significant.

4. Discussion

The main finding in the present study is that circulating levels of endotoxin are elevated after a high glucose meal among overweight non-T2DM subjects and these unfavorable changes were not observed among non-overweight, non-T2DM as well as T2DM subjects. The significant increase of endotoxin levels among overweight subjects suggests exacerbation of an already dysfunctional gut among these subjects. Physiologically, endogenous insulin released into the circulation as a response to acute hyperglycemia.^[25] During this process and in the presence of a healthy gut such as the case of controls in the present study, no aberrant changes will be observed in endotoxin levels. In the case of overweight subjects, however, the significant increase in endotoxin levels following a high glucose load implicates the presence of insulin-resistance and probably systemic low-grade chronic inflammation within this cohort, making the leaky gut even leakier. Chronic exposure to endotoxin may increase T2DM risk since higher levels of endotoxin, in general, is associated with insulin resistance.^[11] However, only a few studies have evaluated the influence of nutrients and/or specific food types on endotoxemia.^[26]

In our study, endotoxin levels following a high-glucose meal showed a dramatic increase only in the overweight and not in the T2DM group. Among several possibilities, is the lack of any reduction in endotoxin levels in the T2DM group may be due to the differences in the subjects dietary make-up. The main effect, however, might be the medications that this group has been taking which include metformin. Metformin has been shown to drastically improve the gut microbiota composition among T2DM.^[5,27,28] This could probably explain why endotoxin at baseline for the T2DM group is much lower than the overweight group and why the response to a glucose load was almost as similar to controls. The significant increase in endotoxin observed in the overweight group can be explained by an already existing insulin-resistant state in the subjects, as evidenced by the modest but significantly higher baseline endotoxin levels compared to other groups.

Finally, in the present study, we observed a significant inverse association between endotoxin and post-prandial LDL-cholesterol. While no significant differences were observed between

Table 3
Bivariate associations between lipids, glucose and endotoxin.

Time (Hour)	Total Cholesterol (mmol/L)	Glucose (mmol/L)*	HDL-Cholesterol (mmol/L)	LDL-Cholesterol (mmol/L)	Triglycerides (mmol/L)*
Overall (N=64)					
0 – Hour	0.04	0.17	0.26	–0.12	0.12
2 – Hour	–0.24	–0.25	0.21	–0.37†	0.10
4 – Hour	–0.09	0.21	–0.11	–0.16	0.27
Control (N=14)					
0 – Hour	0.43	0.49	0.42	–0.27	0.40
2 – Hour	0.20	0.37	0.47	–0.43	0.37
4 – Hour	–0.33	0.07	–0.29	–0.36	0.19
Overweight (N=16)					
0 – Hour	0.13	0.46	0.48	0.06	–0.46
2 – Hour	–0.21	0.43	0.25	–0.61	0.43
4 – Hour	–0.23	0.37	–0.05	–0.05	0.28
T2DM (N=34)					
0 – Hour	0.04	0.36	0.15	–0.04	0.23
2 – Hour	–0.19	–0.03	0.11	–0.28	0.17
4 – Hour	–0.16	0.37	–0.05	–0.27	0.36

Data presented as Spearman correlation coefficient.

* denotes non-normal variables.

† denotes P value $<.05$.

LDL-cholesterol over time in all groups, most studies indicate LDL-cholesterol levels to be lower postprandial as compared to fasting levels, at least in people with T2DM or with insulin resistance.^[29] This could explain why increasing endotoxin levels, particularly in the overweight and T2DM groups, translated to lower LDL-cholesterol levels over-all. The lack of associations between endotoxin and other cardiometabolic parameters in the present study does not supersede our previous findings on the strong relationship between endotoxin and these measures after high fat intake,^[30] and the lack of associations may be due to the sample size issues.

The authors acknowledge some limitations and these include the small sample size and other factors not documented in the

study such as lifestyle and mode of nutrition, which can significantly modify the gut microbiome. Further and longer prospective investigations using the full components of metabolic syndrome,^[31] with a larger number of subjects and inclusion of more inflammatory markers may provide clear insights into how glucose-induced endotoxemia affect the over-all metabolic picture of non-T2DM overweight subjects. Lastly, Arab women are more prone to develop obesity secondary to religious and cultural norms prevalent in the region, making them at higher risk for T2DM and therefore the study focused on this population. Whether the same altered endotoxin levels apply to men of varying glycemic levels need to be investigated separately. The study is nevertheless the first to explore the effects of high glucose




Figure 1. Correlation between log LDL-cholesterol (mmol/L) and Log Endotoxin (EU/ml) at 2-hours in overall patients.

load in endotoxin levels among Arab ethnic women with varying levels of glycemia. This has clinical implications, especially in the region where the concept of gut microbiota and endotoxin in particular, and how it affects obesity-related diseases, remain to be explored.

5. Conclusions

In conclusion, an acute high glucose load promotes endotoxemia among overweight subjects and exacerbates dysbiosis. This effect is not observed among T2DM patients on metformin as well as non-obese, non-T2DM subjects.

Acknowledgments

The authors thank the volunteers and research team from the different primary care centers for the recruitment of subjects.

Author contributions

“Conceptualization, D.A and P.T.; methodology, D.A. and M. M.E; formal analysis, S.D.H. investigation, D.A., M.G.A.A., M. M.E; S.S.; K.W.; resources, N.M.A.; data curation, K.W.; writing—original draft preparation, M.G.A.A.; writing—review and editing, S.S., P.T; supervision, N.M.A.; project administration, D. A; funding acquisition, N.M.A. All authors have read and agreed to the published version of the manuscript.

Conceptualization: Dara Al-Disi, Philip McTernan.

Formal analysis: Syed Danish Hussain.

Funding acquisition: Nasser Al-Daghri.

Investigation: Dara Al-Disi, Shaun Sabico, Kaiser Wani, Mona M. Elshafie.

Methodology: Mohammed Ghouse Ahmed Ansari, Shaun Sabico, Kaiser Wani, Mona M. Elshafie.

Supervision: Philip McTernan, Nasser Al-Daghri.

Writing – original draft: Mohammed Ghouse Ahmed Ansari.

Writing – review & editing: Dara Al-Disi, Shaun Sabico, Kaiser Wani, Syed Danish Hussain, Mona M. Elshafie, Philip McTernan, Nasser Al-Daghri.

References

- Cho NH, Shaw JE, Karuranga S, et al. IDF diabetes atlas: global estimates of diabetes prevalence for 2017 and projections for 2045. *Diabetes Res Clin Pract* 2018;138:271–81.
- Blüher M. Obesity: global epidemiology and pathogenesis. *Nat Rev Endocrinol* 2019;15:288–98.
- Ng M, Fleming T, Robinson M, et al. Global, regional, and national prevalence of overweight and obesity in children and adults during 1980–2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet* 2014;384:766–81.
- Jiang X, Fan X, Wu R, et al. The effect of care intervention for obese patients with type II diabetes. *Medicine (Baltimore)* 2017;96:e7524–17524.
- Bauer PV, Duca FA, Waise TMZ, et al. Metformin alters upper small intestinal microbiota that impact a Glucose-SGLT1-Sensing glucoregulatory pathway. *Cell Metab* 2018;27:101–17. e105.
- Al-Daghri NM, Al-Attas OS, Alkharfy KM, et al. Association of VDR-gene variants with factors related to the metabolic syndrome, type 2 diabetes and vitamin D deficiency. *Gene* 2014;542:129–33.
- Li C, Li X, Han H, et al. Effect of probiotics on metabolic profiles in type 2 diabetes mellitus: A meta-analysis of randomized, controlled trials. *Medicine (Baltimore)* 2016;95:e4088–14088.
- Sato J, Kanazawa A, Watada H. Type 2 Diabetes and Bacteremia. *Ann Nutr Metab* 2017;71(Suppl 1):17–22.
- Makki K, Deehan EC, Walter J, et al. The impact of dietary fiber on gut microbiota in host health and disease. *Cell Host Microbe* 2018;23:705–15.
- Cândido FG, Valente FX, Grzeskowiak ŁM, et al. Impact of dietary fat on gut microbiota and low-grade systemic inflammation: mechanisms and clinical implications on obesity. *Int J Food Sci Nutr* 2018;69:125–43.
- Cani PD, Amar J, Iglesias MA, et al. Metabolic endotoxemia initiates obesity and insulin resistance. *Diabetes* 2007;56:1761–72.
- Al-Attas OS, Al-Daghri NM, Al-Rubeaan K, et al. Changes in endotoxin levels in T2DM subjects on anti-diabetic therapies. *Cardiovasc Diabetol* 2009;8:20–120.
- Schachter J, Martel J, Lin C-S, et al. Effects of obesity on depression: a role for inflammation and the gut microbiota. *Brain Behav Immun* 2018;69:1–8.
- Gangarapu V, Yıldız K, Ince AT, et al. Role of gut microbiota: obesity and NAFLD. *Turk J Gastroenterol* 2014;25:133–40.
- Jovanovich A, Isakova T, Stubbs J. Microbiome and cardiovascular disease in CKD. *Clin J Am Soc Nephrol* 2018;13:1598–604.
- Brandsma E, Kloosterhuis NJ, Koster M, et al. A proinflammatory gut microbiota increases systemic inflammation and accelerates atherosclerosis. *Circ Res* 2019;124:94–100.
- Fukui H. Gut microbiota and host reaction in liver diseases. *Microorganisms* 2015;3:759–91.
- Sabico S, Al-Mashharawi A, Al-Daghri NM, et al. Effects of a 6-month multi-strain probiotics supplementation in endotoxemic, inflammatory and cardiometabolic status of T2DM patients: a randomized, double-blind, placebo-controlled trial. *Clin Nutr* 2019;38:1561–9.
- He J, Zhang F, Han Y. Effect of probiotics on lipid profiles and blood pressure in patients with type 2 diabetes: a meta-analysis of RCTs. *Medicine (Baltimore)* 2017;96:e9166–19166.
- Araújo JR, Tomas J, Brenner C, et al. Impact of high-fat diet on the intestinal microbiota and small intestinal physiology before and after the onset of obesity. *Biochimie* 2017;141:97–106.
- Probiotics Market by Application (Functional Food & Beverages [Dairy Products, Non-dairy Beverages, Infant Formula, Cereals], Dietary Supplements, Feed), Ingredient (Bacteria, Yeast), Form (Dry, Liquid), End User, and Region – Global Forecast to 2023. Available online: <https://www.marketsandmarkets.com/Market-Reports/probiotic-market-advanced-technologies-and-global-market-69.html> (accessed on January 31, 2020)
- Gentile CL, Weir TL. The gut microbiota at the intersection of diet and human health. *Science* 2018;362:776–80.
- Laugerette F, Alligier M, Bastard J-P, et al. Overfeeding increases postprandial endotoxemia in men: inflammatory outcome may depend on LPS transporters LBP and sCD14. *Mol Nutr Food Res* 2014;58:1513–8.
- Al-Disi DA, Al-Daghri NM, Khan N, et al. Postprandial effect of a high-fat meal on endotoxemia in Arab women with and without insulin-resistance-related diseases. *Nutrients* 2015;7:6375–89.
- Toschi E, Camastra S, Sironi AM, et al. Effect of acute hyperglycemia on insulin secretion in humans. *Diabetes* 2002;51(Suppl 1):S130–3.
- Boroni Moreira AP, de Cássia Gonçalves Alfenas R. The influence of endotoxemia on the molecular mechanisms of insulin resistance. *Nutr Hosp* 2012;27:382–90.
- Kim J, Kwak HJ, Cha J-Y, et al. Metformin suppresses lipopolysaccharide (LPS)-induced inflammatory response in murine macrophages via activating transcription factor-3 (ATF-3) induction. *J Biol Chem* 2014;289:23246–55.
- Lee H, Lee Y, Kim J, et al. Modulation of the gut microbiota by metformin improves metabolic profiles in aged obese mice. *Gut Microbes* 2018;9:155–65.
- Lund SS, Petersen M, Frandsen M, et al. Agreement between fasting and postprandial LDL cholesterol measured with 3 methods in patients with type 2 diabetes mellitus. *Clin Chem* 2011;57:298–308.
- Harte AL, Varma MC, Tripathi G, et al. High fat intake leads to acute postprandial exposure to circulating endotoxin in type 2 diabetic subjects. *Diabetes Care* 2012;35:375–82.
- Fattahi MR, Niknam R, Safarpour A, et al. The prevalence of metabolic syndrome in non-alcoholic fatty liver disease; a population-based study. *Middle East J Dig Dis* 2016;8:131–7.