

Pontes, H.M., Kuss, D.J. & Griffiths, M.D. (2017). Psychometric assessment of Internet Gaming Disorder in neuroimaging studies: A systematic review. In Montag, C. & Reuter, M. (Eds.), *Internet Addiction Neuroscientific Approaches and Therapeutical Implications* (pp.181-208). New York: Springer.

Abstract

Background: Little attention has been paid to research on Internet Gaming Disorder (IGD) using neuroimaging techniques even though this type of research is of key importance for the formal recognition of IGD as an independent disorder. **Aims:** The aim of the present study was to conduct a systematic review of some of the methodological and assessment characteristics in neuroimaging studies that have been published from May 2013 to January 2016 and that have assessed IGD whilst also having used a neuroimaging technique to gather neurobiological evidence of this potential disorder. **Methods:** Systematic electronic searches using strict inclusion and exclusion criteria were conducted on ProQuest (in the following scholarly databases: ProQuest Psychology Journals, PsycARTICLES, PsycINFO, Applied Social Sciences Index and Abstracts, and ERIC) and on MEDLINE to identify potential eligible studies. **Results:** A total of 853 studies were screened and after a careful systematic selection process, fourteen studies were found to meet the inclusion criteria. Based on the findings, it was concluded that research on IGD using neuroimaging techniques is on the rise. Additionally, this systematic review offers ten practical recommendations to authors based on the methodological and assessment shortcomings of extant empirical studies on IGD using neuroimaging techniques. **Conclusions:** It is envisaged that the results of this study will help improve the overall quality of research on IGD using neuroimaging techniques.

1. Introduction

Since the introduction of the concept of **Internet Gaming Disorder** (IGD) as a tentative disorder in the latest (fifth) edition of the *Diagnostic and Statistical Manual of Mental Disorders* (American Psychiatric Association, 2013), research into IGD using neuroimaging techniques has steadily increased. Neuroimaging studies appear to offer several advantages over traditional self-report and clinical research by highlighting specific areas of the brain involved in the development and maintenance processes of addiction (Kuss & Griffiths, 2012a). More recently, the APA (2013) defined IGD as a behavior that refers to “persistent and recurrent engagement in video games, often with other players, leading to clinically significant impairments or distress as indicated by five (or more) of the following nine criteria in a 12-month period: (1) preoccupation with games; (2) withdrawal symptoms when gaming is taken away; (3) tolerance, resulting in the need to spend increasing amounts of time engaged in games; (4) unsuccessful attempts to control participation in games; (5) loss of interest in previous hobbies and entertainment as a result of, and with the exception of, games; (6) continued excessive use of games despite knowledge of psychosocial problems; (7) deceiving family members, therapists, or others regarding the amount of gaming; (8) use of games to escape or relieve negative moods; and (9) jeopardizing or losing a significant relationship, job, or education or career opportunity because of participation in games (p.795)”.

Numerous studies have systematically reported potentially harmful effects games can have on human health because of their potentially addictive features (Eichenbaum, Kattner, Bradford, Gentile, & Green, 2015; Lehenbauer-Baum et al., 2015; Schmitt & Livingston, 2015) and overall detrimental effects (Brunborg, Mentzoni, & Frøyland, 2014; Haghbin, Shaterian, Hosseinzadeh, & Griffiths, 2013; Hull, Williams, & Griffiths, 2013; Kuss & Griffiths, 2012b) both from a psychosocial and neuroscientific standpoint. From a psychosocial viewpoint, such harmful effects

related to addiction to gaming can include decreased levels of exercise and sports (Henchoz et al., 2014), sacrificing work, education, hobbies, socializing, time with partner/family, and sleep (Griffiths, Davies, & Chappell, 2004), impaired decision-making (Yao et al., 2015), poorer psychosomatic health (Wittek et al., 2015), decreased emotional and behavioral functioning (Baer, Saran, & Green, 2012), increased stress (Snodgrass et al., 2014), greater incidence of psychiatric symptoms (Király, Urbán, et al., 2015; Vukosavljevic-Gvozden, Filipovic, & Opacic, 2015), lower expected college engagement and grades in adolescent students (Schmitt & Livingston, 2015), decreased academic performance (Brunborg et al., 2014), lower levels of sociability, self-efficacy and satisfaction with life (Festl, Scharnow, & Quandt, 2013), in addition to other psychiatric disorders and abnormal behaviors (see Griffiths, Király, Pontes, & Demetrovics, 2015; Griffiths, Kuss, & King, 2012; Kuss & Griffiths, 2012b).

On the other hand, from a neuroscientific perspective, a systematic review of neuroimaging studies of Internet and gaming addiction (Kuss & Griffiths, 2012a) found that excessive Internet use and gaming is associated with changes on the behavioral, as well as molecular and neural circuitry levels, providing objective evidence of the biological similarity between Internet and gaming addiction and more traditional substance-related addictions. Previous studies (e.g., Dong, DeVito, Du, & Cui, 2012; Dong & Potenza, 2014; Dong, Shen, Huang, & Du, 2013; Liu et al., 2015) have found that IGD is associated with abnormal activations in frontal, insular, temporal, and parietal cortices when affected individuals perform tasks related to impulse control. Additionally, previous structural studies have found that IGD was associated with structural abnormalities in gray matter, such as decreased lower gray matter density in the bilateral inferior frontal gyrus, left cingulate gyrus, insula, right precuneus, and right hippocampus (Lin, Dong, Wang, & Du, 2015). In addition, IGD has also found to be associated with lower white matter

density in the inferior frontal gyrus, insula, amygdala, and anterior cingulate, brain regions that are involved in decision-making, behavioral inhibition and emotional regulation (Lin, Dong, et al., 2015).

1.1. Objectives

Since IGD may be related to a wide range of health and psychiatric disorders as reported by previous studies, it has become vital to understand and evaluate the potential methodological shortcomings of research on IGD using neuroimaging techniques in order to refine future research. For this reason, and given the ongoing debates surrounding the issue of assessment of IGD and the need for unification in the way assessment of this disorder is carried out (Griffiths, King, & Demetrovics, 2014; Griffiths, van Rooij, et al., 2015; Petry & O'Brien, 2013; Petry et al., 2015; Petry et al., 2014; Pontes & Griffiths, 2015b), the present chapter systematically reviews the extant neurobiological evidence of studies that have adopted commonly used types of neuroimaging techniques to investigate the psychobiology of IGD. More specifically, the main aim of this chapter is to investigate what are (if any) the caveats and potential biases and limitations stemming from an assessment of IGD during participant recruitment in such studies. The secondary objective of this chapter is to ascertain the publication rates of IGD studies using neuroimaging techniques as of May 2013 and also briefly summarize the main findings of these studies. To the best of the authors' knowledge, no previous review has attempted to summarize and critique the existing evidence regarding the assessment of IGD in neuroimaging studies (as opposed to describing neuroimaging evidence for Internet and gaming addiction; Kuss & Griffiths, 2012b). This chapter contributes a critical discussion of current practices in the assessment of IGD and may pave the way for new methodologically robust research.

2. Method

2.1. Eligibility criteria

To be eligible for inclusion in this systematic review, only original studies investigating IGD and its associated neurological correlates were included. Additionally, eligible studies had to: (i) assess IGD or direct effects of gaming on neurological functioning, (ii) be an empirical study, (iii) use neuroimaging techniques, (iii) be published in a scholarly peer-reviewed journal, and (iv) be written in English, Spanish, German, Polish or Portuguese language. Searches were limited to articles published from May 2013 to January 2016, because IGD was officially defined and conceptualized by the APA in May 2013, which followed the publication of psychometric tools using this framework to assess IGD. Studies were excluded from review if they were (i) unpublished dissertation and thesis studies, (ii) single-case reports ($N = 1$), and (iii) review studies.

2.2. Information sources and search

The identification of studies was carried out by performing electronic searches on *ProQuest*, which included the following databases: *Applied Social Sciences Index and Abstracts (ASSIA)*, *ERIC*, *ProQuest Psychology Journals*, *PsycARTICLES*, and *PsycINFO*. An additional independent search was carried out on *MEDLINE* to enhance the accuracy of the results regarding the systematic search of relevant studies. The search strategy adopted to identify relevant papers in the aforementioned databases sought to include and be able to retrieve the most common types of neuroimaging techniques employed in research on IGD (i.e., electroencephalogram [EEG], positron emission tomography [PET], single photon emission computed tomography [SPECT], functional magnetic resonance imaging [fMRI], structural magnetic resonance imaging [sMRI],

diffusion-tensor imaging [DTI]) as reported in a previous systematic review (i.e., Kuss & Griffiths, 2012a). As a result, the following search strategy was used:

(patholog* OR problem* OR addict* OR compulsive OR dependen* OR disorder*) AND (video OR computer OR internet) gam* AND (neuroimaging OR eeg OR pet OR spect or fmri OR smri OR dti)

2.3. Study selection and data collection processes

Following the initial literature searches, each study's title and abstract were screened for eligibility. Full texts of all potentially relevant studies were then retrieved and further examined for eligibility. The flow diagram in Figure 1 details the selection process. Information from the included studies was recorded in an electronic spreadsheet after in-depth analysis. The overall data extracted from the studies reviewed subdivided into two larger overarching groups: (i) methodological characteristics and (ii) instrument characteristics. The information extracted concerning the studies' methodological characteristics included: provenience of the sample recruited (i.e., country), sample size, gender distribution, age range (and mean age), sample characteristics, neuroimaging technique used, study aims, and main findings. The information extracted regarding the features of the instruments used to diagnose IGD included: instrument utilized, item sensitivity, criteria included, time scale, theoretical framework, suitability to assess IGD, alignment with the nine IGD criteria, and missing IGD criteria.

3. Results

3.1. Study selection

After performing the aforementioned electronic searches, a total of 853 studies (*ProQuest* $n = 745$; *MEDLINE* $n = 108$) were initially identified, with the search performed on the *ProQuest* website yielding the following results: *ProQuest Psychology Journals* $n = 524$; *PsycARTICLES* $n = 115$; *PsycINFO* $n = 106$; *Applied Social Sciences Index and Abstracts* $n = 0$; and *ERIC* $n = 0$. All 853 papers had their titles and abstracts screened, resulting in the exclusion of 833 papers that were of no relevance for the present review. Consequently, a total of 20 studies were eligible for further review. Of these, six papers had to be further excluded because they were either duplicated ($n = 2$), did not assess IGD ($n = 1$), or were review papers ($n = 3$). Following this study selection process, 14 studies were eligible for further analysis as they fully endorsed all inclusion criteria (see Figure 1).

[Please insert Figure 1 about here]

3.2. *Publication rate of peer-reviewed neuroimaging studies on IGD*

One of the conditions set by the APA in the DSM-5 was that if IGD is to be recognized as an independent disorder in the future, more studies should be carried out to help ascertain the prevalence rates of IGD across the globe, its clinical course and possible genetic influences, and potential biological factors, based on, for instance, brain imaging data (APA, 2013). For this reason, research on IGD employing neuroimaging techniques is of utmost importance not only because of its methodological capabilities, but also because the apparent weight it may carry towards the formal recognition of IGD as an independent disorder in the future (as neurobiological empirical evidence appears to be given more weight than psychological empirical evidence when

considering the inclusion of behavioral addictions – such as IGD and sex addiction – in the DSM [Griffiths, in press]). In light of this, Figure 2 shows a clear trend towards an increase of published peer-reviewed studies on IGD using neuroimaging techniques. This increase trend is most obvious from the period of 2014 to 2015 because the official diagnosis of IGD only appeared in the scientific literature in May 2013 with the publication of the DSM-5, and also because the systematic searches performed for this review were conducted in January 2016.

[Please insert Figure 2 about here]

3.3. Neuroimaging of IGD

In order to present some of the latest neuroimaging research on IGD, this section briefly summarizes the main findings of the studies that fully met the inclusion criteria of this review as previously outlined. Ding et al. (2014) found that the prefrontal cortex may be involved in the circuit modulating impulsivity, while its impaired function may be related to high impulsivity in adolescents with IGD, which in turn, may contribute to the IGD process. Sun et al. (2014) concluded that diffusional kurtosis imaging (DKI) can detect subtle differences in gray matter microstructure between IGD and healthy individuals, and that DKI can provide sensitive imaging biomarkers for assessing the severity of IGD. In the study conducted by Dieter et al. (2015), the authors found that disordered gamers tend to identify significantly more with their avatar in comparison to non-disordered gamers. The authors also noted that the avatar might increasingly replace gamers' ideal self as gaming addiction progresses. H. Wang et al. (2015) found that individuals with IGD presented with significant gray matter volume reduction in the bilateral anterior cingulate cortex and other brain regions when compared to healthy individuals.

Additionally, IGD was also found to compromise both behavioral and neural structure in adolescents with IGD. Xing et al. (2014) found that the abnormal structural connectivity in the right salience network was associated with impaired executive function in adolescents with IGD, with structural connectivity differences found between IGD adolescents and healthy controls. Yuan et al. (2016) found that differences between individuals with IGD and healthy controls in the striatum volume and frontostriatal circuits resting-state functional connectivity (RSFC) emerged. Additionally, cognitive control deficits detected in IGD were associated with reduced frontostriatal RSFC strength. In addition to these findings, Hahn et al. (2014) concluded that converging evidence for a general reward system deficiency in frequent online gamers exists as frequent players displayed significantly decreased neural activation when anticipating both small and large monetary rewards in the ventral striatum. Luijten, Meerkerk, Franken, van de Wetering, and Schoenmakers (2015) found reduced inhibitory control amongst individuals with IGD but no evidence was found for reduced error processing. Furthermore, attentional control and error processing were mostly intact in IGD individuals. At the molecular level, Tian et al. (2014) found that the D_2 receptor level is significantly associated with glucose metabolism in disordered gamers, indicating that $D_2/5-HT_{2A}$ receptor-mediated dysregulation of the orbitofrontal cortex could underlie a mechanism for loss of control and compulsive behavior in individuals with IGD. Duven, Müller, Beutel, and Wölfling (2015) reported that tolerance effects are present in patients with IGD and noted that the initial orienting toward the gaming reward is suggested to consume more capacity for patients with IGD when compared to healthy controls. Lin, Jia, Zang, and Dong (2015) found that individuals with IGD demonstrated lower low-frequency fluctuation (fALFF) values in superior temporal gyrus and higher fALFF values in the cerebellum. Y. Wang et al. (2015) found that adolescents with IGD exhibited decreased voxel-mirrored homotopic connectivity (VMHC)

between the left and right superior frontal gyrus (orbital part), inferior frontal gyrus (orbital part), middle frontal gyrus and superior frontal gyrus. Son et al. (2015) found that lower absolute beta power can be used as a potential trait marker of IGD. Finally, Yuan et al. (2013) found that adolescents with IGD presented with abnormal amplitude of low frequency fluctuations in comparison to controls.

In summary, with regards to the functional and structural changes found in IGD individuals when compared to healthy controls, these are often observed within specific brain areas in the frontal lobe (i.e., dorsolateral prefrontal cortex, orbitofrontal cortex, prefrontal gyrus, and the middle frontal gyrus), parietal and temporal lobes (i.e., parahippocampal gyrus), basal ganglia, thalamus, insula and the cerebellum. A finding that mirrors those recently report in similar review studies (Lemos, Diniz, Peres, & Sougey, 2014). Furthermore, regardless of the neuroimaging technique utilized, these studies provide preliminary evidence that illustrate converging points and similarities between substance use disorders and IGD, especially during craving processes, a finding that has been previously reported (Smith, Hummer, & Hulvershorn, 2015) and that warrants further investigation as at this point, it may be premature to draw definite conclusions regarding the similarities between IGD and substance use disorders as the definition of IGD currently includes many features present in the definition and diagnostic framework of substance use disorders. Taken together, the main findings reported across all reviewed studies do not provide a complete picture of neuroimaging studies on IGD as other potentially useful findings from similar studies may be missing due to strict inclusion criteria employed in this review.

3.4. Methodological characteristics of studies

In order to assess some of the methodological features of existing neuroimaging studies examining IGD, key variables were analyzed amongst the reviewed studies, such as: geographical location of the sample recruited (i.e., country), sample size, gender distribution, age range and mean, sample characteristics, neuroimaging technique used, study aims, and main findings (see Table 1).

[Please insert Table 1 about here]

Regarding the cultural context of the samples recruited, nine studies out of fourteen recruited Chinese samples (Ding et al., 2014; Lin, Jia, et al., 2015; Sun et al., 2014; Tian et al., 2014; H. Wang et al., 2015; Y. Wang et al., 2015; Xing et al., 2014; Yuan et al., 2013; Yuan et al., 2016), three studies used German samples (Dieter et al., 2015; Duven et al., 2015; Hahn et al., 2014), one study used a Dutch sample (Luijten et al., 2015), and one study used a South Korean sample (Son et al., 2015). As for the sample sizes, studies recruited from a minimum of 26 disordered gamers in one study (Tian et al., 2014) to a maximum 87 disordered gamers in another study (Yuan et al., 2016). The mean number of disordered gamers recruited across all 14 studies was 43 ($SD = 18.3$). In terms of gender distribution across all studies, only one study had an even split between genders (Ding et al., 2014) while six studies recruited male-only samples (Duven et al., 2015; Hahn et al., 2014; Lin, Jia, et al., 2015; Luijten et al., 2015; Son et al., 2015; Tian et al., 2014). Although Table 1 summarizes the age of gamers recruited, little information was available in the reviewed studies regarding the overall age range of all gamers as the studies usually reported this information only for each group of participants (e.g., experimental and control groups). Despite this limitation, the data gathered suggests that most IGD-affected gamers recruited were

in their mid- to late adolescence given the characteristics of the samples recruited across all studies. Almost half of all reviewed studies included gamers that were either addicted to *World of Warcraft* (*WoW*) (i.e., Hahn et al., 2014; Tian et al., 2014; Yuan et al., 2013) or *League of Legends* (*LoL*) (i.e., Xing et al., 2014; Yuan et al., 2016) or included both types of gamers (i.e., participants addicted to *WoW* and participants addicted to *LoL*) (i.e., Dieter et al., 2015) only. The procedure of recruiting participants having problems with only a particular game or specific game genre is questionable because it clearly limits the external validity of the findings reported in such studies as it is unclear to what extent these findings can be replicated across all disordered gamers. Interestingly, all studies provided some form of control group, as well as inclusion, and exclusion criteria in the recruitment of participants, which certainly helped mitigate some of the potential threats to the internal validity of the studies in question.

Finally, as to the neuroimaging techniques used, six studies employed resting-state functional magnetic resonance imaging (rsfMRI) (Hahn et al., 2014; Lin, Jia, et al., 2015; Y. Wang et al., 2015; Xing et al., 2014; Yuan et al., 2013; Yuan et al., 2016), four studies used functional magnetic resonance imaging (fMRI) (Dieter et al., 2015; Ding et al., 2014; Luijten et al., 2015; Sun et al., 2014), one study used positron emission tomography (PET) (Tian et al., 2014), one study used electroencephalography (EEG) (Duven et al., 2015), one study used resting-state electroencephalography (rsEEG) (Son et al., 2015), and one study used voxel-based morphometry (VBM) (H. Wang et al., 2015).

3.5. Methodological characteristics of the instruments used to assess IGD

Table 2 provides an overview regarding the preferred methods for diagnosing IGD among the reviewed studies and the features of these instruments and/or diagnostic approaches. Even

though the main construct under investigation of the studies reviewed was IGD, the diagnosis of this disorder was carried out most of the time using measures that were either (i) designed to assess generalized Internet addiction (IA) or (ii) measures that had their theoretical framework based on the concept of IA or non-validated diagnostic criteria. More specifically, in terms of diagnosis of IGD, five studies (i.e., Ding et al., 2014; Sun et al., 2014; H. Wang et al., 2015; Y. Wang et al., 2015; Yuan et al., 2013) used Young's Diagnostic Questionnaire (YDQ) (Young, 1998b) with Beard and Wolf's (2001) suggested criteria for IA, two studies (i.e., Lin, Jia, et al., 2015; Xing et al., 2014) used the Internet Addiction Test (IAT) (Young, 1998a), one study (i.e., Luijten et al., 2015) used the Videogame Addiction Test (VAT)¹ (Van Rooij et al., 2012), one study (i.e., Tian et al., 2014) used Tao and colleagues' (2010) diagnostic criteria for IA, one study (i.e., Dieter et al., 2015) used the checklist for the Assessment of Internet and Computer Game Addiction (AICA-C) (Wölfling, Beutel, & Müller, 2012), another study (i.e., Duven et al., 2015) used the Scale for the Assessment of Internet and Computer Game Addiction (AICA-S, CSV-S) (Wölfling, Müller, & Beutel, 2011), and one study (i.e., Hahn et al., 2014) adopted a frequency criterion based on time spent gaming (i.e., playing at least four times per week for one hour or more for at least one year). Finally, only two studies (i.e., Son et al., 2015; Yuan et al., 2016) used the DSM-5 criteria for IGD (APA, 2013) to diagnose IGD.

[Please insert Table 2 about here]

¹ Although this test presents with some degree of validity to assess gaming addiction, the theoretical framework of this test was based on Compulsive Internet Use theory derived from the DSM-IV criteria for dependence and obsessive-compulsive disorder and the components model of addiction see van Rooij et al. (2012).

Although most studies relied on only one instrument or criterion to diagnose and assess the severity of IGD, five studies, including the two studies that used the IGD criteria to diagnose IGD, adopted two different instruments to diagnose and assess the severity of IGD (Ding et al., 2014; Son et al., 2015; Sun et al., 2014; Y. Wang et al., 2015; Yuan et al., 2016), with these instruments either being the Chen Internet Addiction Scale (CIAS) (i.e., Ding et al., 2014; Sun et al., 2014; Y. Wang et al., 2015) or the Internet Addiction Test (IAT) (i.e., Son et al., 2015; Yuan et al., 2016).

4. Discussion

The present chapter sought to systematically review and evaluate some of the methodological and instrument characteristics of the most recent IGD studies that have employed commonly used neuroimaging techniques. In regards to the methodological features assessed in the reviewed studies, data were collected and analyzed concerning the studies' methodological characteristics such as: geographical location of the sample recruited (i.e., country), sample size, gender distribution, age range and mean, sample characteristics, neuroimaging technique used, study aims, and main findings. Additionally, the quality of assessment of IGD amongst neuroimaging studies published after the formulation of the nine IGD criteria in the DSM-5 (APA, 2013) was also assessed by examining the following indicators across all reviewed studies: instrument utilized, item sensitivity, criteria covered, time scale, theoretical framework, suitability to assess IGD, alignment with the nine IGD criteria, and missing IGD criteria.

Overall, this chapter provides objective data concerning a trend suggesting an increase in the publication rates of IGD studies using neuroimaging techniques since the publication of the DSM-5 (APA, 2013). As shown in Figure 2, the number of such studies is increasing and this may be beneficial to the process of formal recognition of IGD in future editions of the DSM (and other

diagnostic manuals). This is because the APA suggested neurobiological evidence for IGD is of the utmost importance. However, quantity does not necessarily equate to quality because there are a series of methodological issues that appear to hinder the progress of this type of research. Based on the overall results found in this review, the following recommendations are made to help improve the overall quality of future studies on IGD using neuroimaging techniques. Some of these recommendations apply to other types of IGD research (e.g., self-report surveys, focus group interviews, experiments, etc.), so the application of such recommendations is advised whenever possible as it may help strengthen the overall quality of future research into IGD.

Methodological recommendations

1. Recruitment of gamers (cultural background): Since most of the reviewed studies were carried out in Asian countries, mostly in China, and only few studies were conducted in Western countries, it is recommend that future studies recruit participants from both Western and Eastern regions of the world where online games are played. This procedure is crucial to improving the quality of research on IGD as replicability of findings across all cultures where online gaming is prevalent is important if research is to progress.
2. Recruitment of gamers (gender and age groups): Based on the findings collated in this review, it is also recommended that researchers recruit samples that are balanced in terms of gender and age. It is important to recruit gamers from diversified cultural backgrounds otherwise, findings may (potentially) be gender/culture biased or gender/culture-specific, and therefore, less likely to be replicated.
3. Recruitment of gamers (clinical vs. experimental groups): Although it is common practice to recruit participants seemingly addicted to specific types of games (e.g., *WoW*, *LoL*, etc.)

or game genres (e.g., Massively Multiplayer Online Role Playing Games), as suggested by the findings of this review, this procedure should be avoided as it has been largely adopted in past gaming addiction research from the early 2000s (see Pontes & Griffiths, 2014) and is known for being problematic because IGD can occur in any type of gaming and therefore is not limited to a specific type of game and/or genre. Ultimately, limiting the recruitment of IGD-affected individuals to specific games may lead to biased results as other disordered gamers who play other games and genres may be overlooked.

4. Neuroimaging data (use of eclectic techniques): One noticeable finding in this review was that a large number of studies employed rsfMRI in order to collect imaging data from gamers. Although this may not be problematic in and of itself, authors are advised to adopt an eclectic approach to imaging data collection by using more heterogeneous and other cutting-edge techniques so more information regarding the neurobiological etiology and course of IGD may be acquired. For instance, the combination of fMRI techniques with insights from PET research provides more direct insights to the biochemical mechanism of human behaviors, including IGD (Ko, Liu, & Yen, 2015). The adoption of such approaches is likely to lead to quicker recognition of IGD by the psychiatric community as an independent disorder because greater and more diversified data will be available.
5. The inherent potential of using fMRI (refining the diagnostic features of IGD): Due to the need for differentiating chemical and non-chemical addictions coupled with the need to refine the diagnostic features of IGD, studies using fMRI techniques are welcome due to their inherent potential for investigating the specific mechanisms of addiction, including response to a substance, vulnerability for addiction, characteristics or symptoms of addictive behavior, and consequences of addiction (Fowler, Volkow, Kassed, & Chang,

2007; Ko et al., 2015). For this reason, neurobiological research into IGD should formulate clear hypotheses derived from previous evidence from brain imaging studies as they might help clarify the utility and appropriateness of having specific diagnostic criteria within the diagnostic framework of IGD. Consequently, this type of research may provide robust evidence to either confirm or disconfirm claims that IGD may arise without associated withdrawal symptoms (Kaptsis, King, Delfabbro, & Gradisar, 2016) or that the tolerance criterion of IGD might be a prominent characteristic of passionate gaming rather than a criterion indicative of *bona fide* gaming addiction (Kardefelt-Winther, 2015).

6. Impacts of IGD on cognition (avoiding premature conclusions): According to Ko et al. (2015), any conclusions regarding cognitive function in IGD would still be premature as the overall effect of IGD on cognitive functioning remains controversial. Contrary to most controlled substances that are known to have damaging effects on the brain, a reasonable assumption is that they impair cognitive function. However, it is known that gaming can exercise and enhance many specific cognitive functions (e.g., Stroud & Whitbourne, 2015; Toril, Reales, & Ballesteros, 2014). For this reason, the hypothesis that gaming in and of itself produces a deficit in cognitive function is questionable because perfect performance in gaming requires good cognitive function (Ko et al., 2015). Thus, studies reporting potentially negative impacts of IGD on cognition should take into account the fact that gaming may also produce positive cognitive effects on gamers.

Assessment of IGD recommendations

7. Assessment of IGD (key assessment issues): As shown in Table 2, although all studies reviewed concerned the phenomenon of IGD, the diagnosis of this potential disorder was

largely conducted by employing assessment instruments that were either (i) designed to assess generalized IA or (ii) had their theoretical framework based on the concept of IA or non-validated diagnostic criteria. This is perhaps one of the most troublesome findings of this review because the use of different, non-specific, and/or psychometrically weak tools to assess IGD not only represents a threat to the call for unification in the assessment of IGD (Griffiths, van Rooij, et al., 2015), but also hinders the progress of research in this field as cross-cultural comparisons between studies are virtually impossible to be achieved under such circumstances (Kuss, Griffiths, Karila, & Billieux, 2014). Moreover, researchers in the field are discouraged to adopt such procedures because they severely compromise the overall validity of neuroimaging studies on IGD since the main construct under investigation is not being properly assessed but instead, other related (and yet different) constructs are being assessed (i.e., generalized IA rather than IGD).

8. Assessment of IGD (frequency of gameplay and IGD): Although it is not common practice to assess IGD based on frequency of gameplay (i.e., time spent gaming), there are still studies using the frequency criterion as a way to diagnosing IGD cases. It has now been long establish in the field that excessive gaming does not necessarily equate to addiction (Griffiths, 2010; Kuss, Louws, & Wiers, 2012) and that there is a difference between engaged and addicted gamers (see Charlton & Danforth, 2007; Fuster, Carbonell, Pontes, & Griffiths, 2015; Pontes, Király, Demetrovics, & Griffiths, 2014). Therefore, we encourage authors to assess and diagnose IGD based on newly developed standardized instruments available that were devised using the updated framework to assess IGD (i.e., the APA's nine criteria) and that have been shown to possess acceptable levels of validity and reliability (such as: Király, Slezcka, et al., 2015; Lemmens, Valkenburg, & Gentile,

2015; Pontes & Griffiths, 2015a; Pontes et al., 2014). By adopting such a recommendation, the goal of unification in the assessment of IGD will be more tangible and less difficult to achieve.

9. Assessment of IGD (heterogeneity issues): A systematic review conducted by King and colleagues (2013) assessing a total of 18 problematic gaming instruments across 63 empirical studies, demonstrated that most tools presented similar limitations including: (i) inconsistent coverage of core addiction indicators, (ii) use of varying cut-off scores to indicate clinical status, (iii) lack of a temporal dimension, (iv) untested or inconsistent dimensionality, and (v) inadequate data on predictive validity and inter-rater reliability. Similarly, the findings of the present chapter mirrored those found by King and colleagues (2013). More specifically, the present review found that the included studies used eight different methods for diagnosing IGD, which suggests great disparity and heterogeneity between the preferred methods for assessing IGD in this new era of research. As noted above, there are valid and reliable psychometric tools designed to assess IGD based on the APA's suggested framework and these should be used. Because of the inherent difficulties in obtaining a gold standard for IGD across studies, authors can mitigate the possible biases stemming from the assessment of this disorder by adopting commonly used tools that assess IGD using APA criteria and possess some degree of validity and reliability. This is an important aspect of research into IGD more generally because (as shown in Table 2) most instruments used to assess IGD are based on theoretical frameworks of IA that lack content validity to assess IGD because many important aspects of the IGD construct are not fully covered.

10. Assessment of IGD (diagnosis vs. severity): Findings in this chapter also found that most of the studies reviewed relied on one instrument or criterion to diagnose IGD and another one to assess the severity of IGD. More specifically, five studies, including the two studies that have used the IGD criteria to diagnose IGD, adopted two different instruments to diagnose and assess the severity of IGD (Ding et al., 2014; Son et al., 2015; Sun et al., 2014; Y. Wang et al., 2015; Yuan et al., 2016), with these instruments either being the Chen Internet Addiction Scale (CIAS) (i.e., Ding et al., 2014; Sun et al., 2014; Y. Wang et al., 2015) or the Internet Addiction Test (IAT) (i.e., Son et al., 2015; Yuan et al., 2016). Although it is not entirely clear as to why two different instruments were used to diagnose and assess the severity of IGD, this certainly does not facilitate a sound assessment of IGD as more noise is introduced into such studies with the use of different assessment tools. Additionally, and perhaps more worryingly, assessing the severity of IGD with generalized IA is not the way forward as issues of construct validity may emerge. Hence, it is recommended to use one of the newly standardized psychometric tools to assess the severity of IGD as well as an initial structured interview to diagnose IGD based on the nine IGD criteria if necessary as they appear to have clinical validity (Ko et al., 2014).

Based on the findings raised and highlighted in this chapter (and the recommendations generated upon them), it is hoped that future IGD studies, especially those using neuroimaging techniques, may improve their methodological and instrumentation features as there are several different ways to overcome such shortcomings that most of the extant studies on IGD using neuroimaging techniques present with.

References

- American Psychiatric Association. (2013). *Diagnostic and Statistical Manual of Mental Disorders (5th ed.)*. Arlington, VA: Author.
- Baer, S., Saran, K., & Green, D. A. (2012). Computer/gaming station use in youth: Correlations among use, addiction and functional impairment. *Paediatrics & Child Health, 17*(8), 427-431.
- Beard, K. W., & Wolf, E. M. (2001). Modification in the Proposed Diagnostic Criteria for Internet Addiction. *CyberPsychology & Behavior, 4*(3), 377-383. doi:10.1089/109493101300210286
- Brunborg, G. S., Mentzoni, R. A., & Frøyland, L. R. (2014). Is video gaming, or video game addiction, associated with depression, academic achievement, heavy episodic drinking, or conduct problems? *Journal of Behavioral Addictions, 3*(1), 27-32. doi:10.1556/JBA.3.2014.002
- Charlton, J. P., & Danforth, I. D. W. (2007). Distinguishing addiction and high engagement in the context of online game playing. *Computers in Human Behavior, 23*(3), 1531-1548. doi:10.1016/j.chb.2005.07.002
- Dieter, J., Hill, H., Sell, M., Reinhard, I., Vollstädt-Klein, S., Kiefer, F., . . . Leménager, T. (2015). Avatar's Neurobiological Traces in the Self-Concept of Massively Multiplayer Online Role-Playing Game (MMORPG) Addicts. *Behavioral Neuroscience, 129*(1), 8-17. doi:10.1037/bne0000025
- Ding, W.-N., Sun, J.-H., Sun, Y.-W., Chen, X., Zhou, Y., Zhuang, Z.-G., . . . Du, Y.-S. (2014). Trait impulsivity and impaired prefrontal impulse inhibition function in adolescents with internet gaming addiction revealed by a Go/No-Go fMRI study. *Behavioral and Brain Functions, 10*(1), 20. doi:10.1186/1744-9081-10-20
- Dong, G., DeVito, E. E., Du, X., & Cui, Z. (2012). Impaired inhibitory control in 'internet addiction disorder': A functional magnetic resonance imaging study. *Psychiatry Research: Neuroimaging, 203*(2-3), 153-158. doi:10.1016/j.psychresns.2012.02.001
- Dong, G., & Potenza, M. N. (2014). A cognitive-behavioral model of Internet gaming disorder: Theoretical underpinnings and clinical implications. *Journal of Psychiatric Research, 58*, 7-11. doi:10.1016/j.jpsychires.2014.07.005
- Dong, G., Shen, Y., Huang, J., & Du, X. (2013). Impaired Error-Monitoring Function in People with Internet Addiction Disorder: An Event-Related fMRI Study. *European Addiction Research, 19*(5), 269-275. doi:10.1159/000346783
- Duven, E. C. P., Müller, K. W., Beutel, M. E., & Wölfling, K. (2015). Altered reward processing in pathological computer gamers – ERP-results from a semi-natural Gaming-Design. *Brain and Behavior, 5*(1), 13-23. doi:10.1002/brb3.293
- Eichenbaum, A., Kattner, F., Bradford, D., Gentile, D. A., & Green, C. S. (2015). Role-Playing and Real-Time Strategy Games Associated with Greater Probability of Internet Gaming Disorder. *Cyberpsychology, Behavior, and Social Networking, 18*(8), 480-485. doi:10.1089/cyber.2015.0092
- Festl, R., Scharnow, M., & Quandt, T. (2013). Problematic computer game use among adolescents, younger and older adults. *Addiction, 108*(3), 592-599. doi:10.1111/add.12016
- Fowler, J. S., Volkow, N. D., Kassed, C. A., & Chang, L. (2007). Imaging the Addicted Human Brain. *Science & Practice Perspectives, 3*(2), 4-16. doi:10.1151/sp07324

- Fuster, H., Carbonell, X., Pontes, H. M., & Griffiths, M. D. (2015). Spanish validation of the Internet Gaming Disorder-20 (IGD-20) Test. *Computers in Human Behavior*, *56*, 215-224. doi:10.1016/j.chb.2015.11.050
- Griffiths, M. D. (2010). The role of context in online gaming excess and addiction: Some case study evidence. *International Journal of Mental Health and Addiction*, *8*(1), 119-125. doi:10.1007/s11469-009-9229-x
- Griffiths, M. D. (in press). Compulsive sexual behaviour as a behavioural addiction: The impact of the Internet and other issues. *Addiction*.
- Griffiths, M. D., Davies, M., & Chappell, D. (2004). Demographic Factors and Playing Variables in Online Computer Gaming. *CyberPsychology & Behavior*, *7*(4), 479-487. doi:10.1089/cpb.2004.7.479
- Griffiths, M. D., King, D. L., & Demetrovics, Z. (2014). DSM-5 internet gaming disorder needs a unified approach to assessment. *Neuropsychiatry*, *4*(1), 1-4. doi:10.2217/np.13.82
- Griffiths, M. D., Király, O., Pontes, H. M., & Demetrovics, Z. (2015). An overview of problematic gaming. In E. Aboujaoude & V. Starcevic (Eds.), *Mental Health in the Digital Age: Grave Dangers, Great Promise* (pp. 27-45). Oxford: Oxford University Press.
- Griffiths, M. D., Kuss, D. J., & King, D. (2012). Video Game Addiction: Past, Present and Future. *Current Psychiatry Reviews*, *8*(4), 308-318. doi:10.2174/157340012803520414
- Griffiths, M. D., van Rooij, A. J., Kardefelt-Winther, D., Starcevic, V., Király, O., Pallesen, S., . . . Demetrovics, Z. (2015). Working towards an international consensus on criteria for assessing Internet Gaming Disorder: A critical commentary on Petry et al. (2014). *Addiction*, *111*, 167-178. doi:10.1111/add.13057
- Haghighi, M., Shaterian, F., Hosseinzadeh, D., & Griffiths, M. D. (2013). A brief report on the relationship between self-control, video game addiction and academic achievement in normal and ADHD students. *Journal of Behavioral Addictions*, *2*(4), 239-243. doi:10.1556/JBA.2.2013.4.7
- Hahn, T., Notebaert, K. H., Dresler, T., Kowarsch, L., Reif, A., & Fallgatter, A. J. (2014). Linking online gaming and addictive behavior: Converging evidence for a general reward deficiency in frequent online gamers. *Frontiers in Behavioral Neuroscience*, *8*:385. doi:10.3389/fnbeh.2014.00385
- Henchoz, Y., Studer, J., Deline, S., N'Goran, A. A., Baggio, S., & Gmel, G. (2014). Video gaming disorder and sport and exercise in emerging adulthood: a longitudinal study. *Behavioral Medicine*(just-accepted), 1-7. doi:10.1080/08964289.2014.965127
- Hull, D. C., Williams, G. A., & Griffiths, M. D. (2013). Video game characteristics, happiness and flow as predictors of addiction among video game players: A pilot study. *Journal of Behavioral Addictions*, *2*(3), 145-152. doi:10.1556/JBA.2.2013.005
- Kaptsis, D., King, D. L., Delfabbro, P. H., & Gradisar, M. (2016). Withdrawal symptoms in internet gaming disorder: A systematic review. *Clinical Psychology Review*, *43*, 58-66. doi:10.1016/j.cpr.2015.11.006
- Kardefelt-Winther, D. (2015). Assessing the diagnostic contribution of internet gaming disorder criteria requires improved content, construct and face validity—a response to Rehbein and colleagues (2015). *Addiction*, *110*(8), 1359-1360. doi:10.1111/add.12987
- King, D. L., Haagsma, M. C., Delfabbro, P. H., Gradisar, M., & Griffiths, M. D. (2013). Toward a consensus definition of pathological video-gaming: A systematic review of psychometric assessment tools. *Clinical Psychology Review*, *33*(3), 331-342. doi:10.1016/j.cpr.2013.01.002

- Király, O., Slecza, P., Pontes, H. M., Urbán, R., Griffiths, M. D., & Demetrovics, Z. (2015). Validation of the Ten-Item Internet Gaming Disorder Test (IGDT-10) and evaluation of the nine DSM-5 Internet Gaming Disorder criteria. *Addictive Behaviors*. doi:10.1016/j.addbeh.2015.11.005
- Király, O., Urbán, R., Griffiths, M. D., Ágoston, C., Nagygyörgy, K., Kökönyei, G., & Demetrovics, Z. (2015). The Mediating Effect of Gaming Motivation Between Psychiatric Symptoms and Problematic Online Gaming: An Online Survey. *Journal of Medical Internet Research*, *17*(4), e88. doi:10.2196/jmir.3515
- Ko, C.-H., Liu, G.-C., & Yen, J.-Y. (2015). Functional Imaging of Internet Gaming Disorder. In C. Montag & M. Reuter (Eds.), *Internet Addiction* (pp. 43-63): Springer International Publishing.
- Ko, C.-H., Yen, J.-Y., Chen, S.-H., Wang, P.-W., Chen, C.-S., & Yen, C.-F. (2014). Evaluation of the diagnostic criteria of Internet gaming disorder in the DSM-5 among young adults in Taiwan. *Journal of Psychiatric Research*, *53*(6), 103-110. doi:10.1016/j.jpsychires.2014.02.008
- Kuss, D. J., & Griffiths, M. D. (2012a). Internet and gaming addiction: A systematic literature review of neuroimaging studies. *Brain Sciences*, *2*(3), 347-374. doi:10.3390/brainsci2030347
- Kuss, D. J., & Griffiths, M. D. (2012b). Online gaming addiction in children and adolescents: A review of empirical research. *Journal of Behavioral Addictions*, *1*(1), 3-22. doi:10.1556/JBA.1.2012.1.1
- Kuss, D. J., Griffiths, M. D., Karila, L., & Billieux, J. (2014). Internet Addiction: A Systematic Review of Epidemiological Research for the Last Decade. *Current Pharmaceutical Design*, *20*(25), 4026-4052. doi:10.2174/13816128113199990617
- Kuss, D. J., Louws, J., & Wiers, R. W. (2012). Online Gaming Addiction? Motives Predict Addictive Play Behavior in Massively Multiplayer Online Role-Playing Games. *Cyberpsychology, Behavior, and Social Networking*, *15*(9), 480-485. doi:10.1089/cyber.2012.0034
- Lehenbauer-Baum, M., Klaps, A., Kovacovsky, Z., Witzmann, K., Zahlbruckner, R., & Stetina, B. U. (2015). Addiction and Engagement: An Explorative Study Toward Classification Criteria for Internet Gaming Disorder. *Cyberpsychology, Behavior, and Social Networking*, *18*(6), 343-349. doi:10.1089/cyber.2015.0063
- Lemmens, J. S., Valkenburg, P. M., & Gentile, D. A. (2015). The Internet Gaming Disorder Scale. *Psychological Assessment*, *27*(2), 567-582. doi:10.1037/pas0000062
- Lemos, I. L., Diniz, P. R. B., Peres, J. F. P., & Sougey, E. B. (2014). Neuroimagem na dependência de jogos eletrônicos: uma revisão sistemática. *Jornal Brasileiro de Psiquiatria*, *63*(1), 57-71.
- Lin, X., Dong, G., Wang, Q., & Du, X. (2015). Abnormal gray matter and white matter volume in 'Internet gaming addicts'. *Addictive Behaviors*, *40*, 137-143. doi:10.1016/j.addbeh.2014.09.010
- Lin, X., Jia, X., Zang, Y.-F., & Dong, G. (2015). Frequency-dependent changes in the amplitude of low-frequency fluctuations in Internet gaming disorder. *Frontiers in psychology*, *6*. doi:10.3389/fpsyg.2015.01471
- Liu, J., Li, W., Zhou, S., Zhang, L., Wang, Z., Zhang, Y., . . . Li, L. (2015). Functional characteristics of the brain in college students with internet gaming disorder. *Brain Imaging and Behavior*, 1-8. doi:10.1007/s11682-015-9364-x

- Luijten, M., Meerkerk, G.-J., Franken, I. H. A., van de Wetering, B. J. M., & Schoenmakers, T. M. (2015). An fMRI study of cognitive control in problem gamers. *Psychiatry Research: Neuroimaging*. doi:10.1016/j.psychresns.2015.01.004
- Petry, N. M., & O'Brien, C. P. (2013). Internet Gaming Disorder and the DSM-5. *Addiction, 108*(7), 1186–1187. doi:10.1111/add.12162
- Petry, N. M., Rehbein, F., Gentile, D. A., Lemmens, J. S., Rumpf, H.-J., Mößle, T., . . . O'Brien, C. P. (2015). Griffiths et al.'s comments on the international consensus statement of Internet Gaming Disorder: furthering consensus or hindering progress? *Addiction, 111*, 167-178.
- Petry, N. M., Rehbein, F., Gentile, D. A., Lemmens, J. S., Rumpf, H., Mößle, T., . . . Borges, G. (2014). An international consensus for assessing internet gaming disorder using the new DSM-5 approach. *Addiction, 109*, 1399-1406. doi:10.1111/add.12457
- Pontes, H. M., & Griffiths, M. D. (2014). Assessment of Internet Gaming Disorder in clinical research: Past and present perspectives. *Clinical Research and Regulatory Affairs, 31*(2-4), 35-48. doi:10.3109/10601333.2014.962748
- Pontes, H. M., & Griffiths, M. D. (2015a). Measuring DSM-5 Internet Gaming Disorder: Development and validation of a short psychometric scale. *Computers in Human Behavior, 45*, 137-143. doi:10.1016/j.chb.2014.12.006
- Pontes, H. M., & Griffiths, M. D. (2015b). New Concepts, Old Known Issues: The DSM-5 and Internet Gaming Disorder and its Assessment. In J. Bishop (Ed.), *Psychological and Social Implications Surrounding Internet and Gaming Addiction* (pp. 16-30). Hershey, PA: Information Science Reference.
- Pontes, H. M., Király, O., Demetrovics, Z., & Griffiths, M. D. (2014). The conceptualisation and measurement of DSM-5 Internet Gaming Disorder: The development of the IGD-20 Test. *PLoS ONE, 9*(10), e110137. doi:10.1371/journal.pone.0110137
- Schmitt, Z. L., & Livingston, M. G. (2015). Video Game Addiction and College Performance Among Males: Results from a 1 Year Longitudinal Study. *Cyberpsychology, Behavior, and Social Networking, 18*(1), 25-29. doi:10.1089/cyber.2014.0403
- Smith, K. L., Hummer, T. A., & Hulvershorn, L. A. (2015). Pathological Video Gaming and Its Relationship to Substance Use Disorders. *Current Addiction Reports, 1*-8. doi:10.1007/s40429-015-0075-6
- Snodgrass, J. G., Lacy, M. G., Dengah II, H. J., Eisenhauer, S., Batchelder, G., & Cookson, R. J. (2014). A vacation from your mind: Problematic online gaming is a stress response. *Computers in Human Behavior, 38*, 248-260. doi:10.1016/j.chb.2014.06.004
- Son, K. L., Choi, J. S., Lee, J., Park, S. M., Lim, J. A., Lee, J. Y., . . . Kwon, J. S. (2015). Neurophysiological features of Internet gaming disorder and alcohol use disorder: a resting-state EEG study. *Transl Psychiatry, 5*, e628. doi:10.1038/tp.2015.124
- Stroud, M. J., & Whitbourne, S. K. (2015). Casual Video Games as Training Tools for Attentional Processes in Everyday Life. *Cyberpsychology, Behavior, and Social Networking*. doi:10.1089/cyber.2015.0316
- Sun, Y., Sun, J., Zhou, Y., Ding, W., Chen, X., Zhuang, Z., . . . Du, Y. (2014). Assessment of in vivo microstructure alterations in gray matter using DKI in internet gaming addiction. *Behavioral and Brain Functions, 10*(1), 37. doi:10.1186/1744-9081-10-37
- Tao, R., Huang, X., Wang, J., Zhang, H., Zhang, Y., & Li, M. (2010). Proposed diagnostic criteria for internet addiction. *Addiction, 105*(3), 556-564. doi:10.1111/j.1360-0443.2009.02828.x

- Tian, M., Chen, Q., Zhang, Y., Du, F., Hou, H., Chao, F., & Zhang, H. (2014). PET imaging reveals brain functional changes in internet gaming disorder. *European Journal of Nuclear Medicine and Molecular Imaging*, *41*, 1388-1397. doi:10.1007/s00259-014-2708-8
- Toril, P., Reales, J. M., & Ballesteros, S. (2014). Video game training enhances cognition of older adults: A meta-analytic study. *Psychology and Aging*, *29*(3), 706-716. doi:10.1037/a0037507
- Van Rooij, A. J., Schoenmakers, T. M., van den Eijnden, R. M., Vermulst, A. A., & Van de Mheen, D. (2012). Video Game Addiction Test: validity and psychometric characteristics. *Cyberpsychology, Behavior, and Social Networking*, *15*(9), 507-511. doi:10.1089/cyber.2012.0007
- Vukosavljevic-Gvozden, T., Filipovic, S., & Opacic, G. (2015). The Mediating Role of Symptoms of Psychopathology Between Irrational Beliefs and Internet Gaming Addiction. *Journal of Rational-Emotive & Cognitive-Behavior Therapy*, *33*(4), 387-405. doi:10.1007/s10942-015-0218-7
- Wang, H., Jin, C., Yuan, K., Shakir, T. M., Mao, C., Niu, X., . . . Zhang, M. (2015). The alteration of gray matter volume and cognitive control in adolescents with internet gaming disorder. *Frontiers in Behavioral Neuroscience*, *9*, 64. doi:10.3389/fnbeh.2015.00064
- Wang, Y., Yin, Y., Sun, Y.-W., Zhou, Y., Chen, X., Ding, W.-N., . . . Du, Y.-S. (2015). Decreased Prefrontal Lobe Interhemispheric Functional Connectivity in Adolescents with Internet Gaming Disorder: A Primary Study Using Resting-State fMRI. *PLoS ONE*, *10*(3), e0118733. doi:10.1371/journal.pone.0118733
- Wittek, C. T., Finserås, T. R., Pallesen, S., Mentzoni, R. A., Hanss, D., Griffiths, M. D., & Molde, H. (2015). Prevalence and predictors of video game addiction: A study based on a national representative sample of gamers. *International Journal of Mental Health and Addiction*, *1*-15. doi:10.1007/s11469-015-9592-8
- Wölfling, K., Beutel, M. E., & Müller, K. W. (2012). Construction of a Standardized Clinical Interview to Assess Internet addiction: First Findings Regarding the Usefulness of AICA-C. *Journal of Addiction Research & Therapy*, *S6:003*. doi:10.4172/2155-6105.S6-003
- Wölfling, K., Müller, K. W., & Beutel, M. (2011). [Reliability and validity of the Scale for the Assessment of Pathological Computer-Gaming (CSV-S)]. *Psychotherapie, Psychosomatik, Medizinische Psychologie*, *61*(5), 216-224. doi:10.1055/s-0030-1263145
- Xing, L., Yuan, K., Bi, Y., Yin, J., Cai, C., Feng, D., . . . Yu, D. (2014). Reduced fiber integrity and cognitive control in adolescents with internet gaming disorder. *Brain Research*, *1586*, 109-117. doi:10.1016/j.brainres.2014.08.044
- Yao, Y.-W., Wang, L.-J., Yip, S. W., Chen, P.-R., Li, S., Xu, J., . . . Fang, X.-Y. (2015). Impaired decision-making under risk is associated with gaming-specific inhibition deficits among college students with Internet gaming disorder. *Psychiatry Research*, *229*(1-2), 302-309. doi:10.1016/j.psychres.2015.07.004
- Young, K. S. (1998a). *Caught in the net: how to recognize the signs of Internet addiction - and a winning strategy for recovering*. New York: John Wiley & Sons, Inc.
- Young, K. S. (1998b). Internet Addiction: The Emergence of a New Clinical Disorder. *CyberPsychology & Behavior*, *1*(3), 237-244. doi:10.1089/cpb.1998.1.237
- Yuan, K., Jin, C., Cheng, P., Yang, X., Dong, T., Bi, Y., . . . Liu, J. (2013). Amplitude of Low Frequency Fluctuation Abnormalities in Adolescents with Online Gaming Addiction. *PLoS ONE*, *8*(11), e78708. doi:10.1371/journal.pone.0078708

Yuan, K., Yu, D., Cai, C., Feng, D., Li, Y., Bi, Y., . . . Tian, J. (2016). Frontostriatal circuits, resting state functional connectivity and cognitive control in internet gaming disorder. *Addiction Biology*. doi:10.1111/adb.12348