

1 **Journal of Strength and Conditioning Research**

2 Title:

3 Match Analysis of U9 and U10 English Premier League Academy Soccer Players using a
4 Global Positioning System: Relevance for Talent Identification and Development.

5

6 Running head:

7 Match Analysis of Youth Soccer

8

9 Laboratory where the research was conducted:

10 Institute of Youth Sport, School of Sport, Exercise and Health Sciences, Loughborough
11 University, Leicestershire, LE11 3TU, The United Kingdom.

12

13 Authors:

14 Heita Goto, John G Morris and Mary E Nevill

15

16 Institution and departments:

17 Institute of Youth Sport, School of Sport, Exercise and Health Sciences, Loughborough
18 University, Leicestershire, LE11 3TU, The United Kingdom.

19

20 Mailing address:

21 Heita Goto

22 Institute of Youth Sport, School of Sport, Exercise and Health Sciences, Loughborough
23 University, Leicestershire, LE11 3TU, The United Kingdom

24

25 TEL: +44 (0)7951 480068

26 FAX:

27 h.goto@lboro.ac.uk

28 heitagoto@hotmail.com

29

30 Funding:

31 No funding was received for this study

- 1 Title:
- 2 Match Analysis of U9 and U10 English Premier League Academy Soccer Players using a
- 3 Global Positioning System: Relevance for Talent Identification and Development.
- 4

ACCEPTED

1 **ABSTRACT**

2 The purpose of this study was to examine the match activity profile of U9 and U10 elite
3 soccer players and to establish if there were any differences between players who were
4 subsequently retained or released by their clubs. Such information should prove valuable in
5 the design of training programs for these very young players and in the talent identification
6 and development process. A Global Positioning System was used to analyze 2-4 inter-
7 academy 6-a-side matches of English Premier League Academy players (U9: N = 22 and
8 U10: N = 12) who trained three times a week (4.5 h) . Speed zones were created based on 5
9 and 10 m sprint times and an independent sample t-test was employed for a statistical
10 analysis.

11 Both squads covered ~4000 m in total or ~4700 m·h⁻¹ during a match (NS between squads),
12 with the U10s tending to cover a greater distance at moderate (p = 0.10) and high speeds (p =
13 0.08) than the U9s. Retained group covered a greater distance than released group (retained
14 vs. released: 4478 ± 513 m vs. 4091 ± 462 m, p < 0.05) during a match and covered a greater
15 distance during low speed running in absolute (1226 ± 259 m vs. 1005 ± 221 m, p < 0.05) and
16 relative (1325 ± 235 m·h⁻¹ vs. 1132 ± 210 m·h⁻¹, p < 0.05) terms.

17 Thus, U9 and U10 players cover over 4000 m in match play and those players who are
18 retained by academies cover a greater distance in total and at low speeds (2.1-3.1 m·s⁻¹). This
19 information may support the preparation of squad training programs and the talent
20 identification and development process.

21 **Key Words:** association football, intermittent exercise, team sport.

1 INTRODUCTION

2 In youth soccer, Global Positioning Systems (GPS) have been commonly employed to
3 analyze match running performance and most information is available for players between 11
4 and 16 years of age (6, 7, 11, 12, 15). For example, the U12 to U16 age groups (N = 20-25 in
5 each group) from two English professional soccer clubs covered $\sim 6200 \text{ m}\cdot\text{h}^{-1}$ (U12) to ~ 7000
6 $\text{m}\cdot\text{h}^{-1}$ (U15 and U16) during an 11-a-side match and the distances covered by high intensity
7 running (with speed zones calculated using a 10 m flying sprint speed from the last 10 m of a
8 20 m sprint test) ranged from $\sim 1800 \text{ m}\cdot\text{h}^{-1}$ for the U12 to $\sim 2200 \text{ m}\cdot\text{h}^{-1}$ for the U16 (15). For
9 14 year old national youth players, the total distance covered during a match was $\sim 6000 \text{ m}$
10 and the players covered $\sim 500 \text{ m}$ by walking, $\sim 3000 \text{ m}$ by jogging, $\sim 1650 \text{ m}$ by medium
11 intensity running, $\sim 700 \text{ m}$ by high intensity running and $\sim 250 \text{ m}$ by sprinting with the speed
12 zones based on absolute speeds ranging from $0\text{-}0.4 \text{ km}\cdot\text{h}^{-1}$ for walking to over $18 \text{ km}\cdot\text{h}^{-1}$ for
13 sprinting (11, 12). However, to date, there are no match analysis data on the distance covered
14 for young elite players aged 9 and 10 years of age and no data concerning the distances
15 covered in small sided games for these age groups. Thus the practical problem being
16 examined here is to address the gap in the literature regarding the match distances covered in
17 small-sided games by U9 and U10 age academy football players.

18
19 It has been argued that the players in the developing stages, such as 9 and 10 year olds,
20 should not be considered as miniature adults (29) and hence, training programs should be
21 specifically designed for these young players. Knowing the running distances during
22 competitive matches for elite youth soccer players may support coaches and sports scientists
23 to prepare training programs which are specific to the players' age and ability (9).
24 Furthermore, if there is a difference between the match performance of young players who

1 are later retained or released by academies this information could assist in both the talent
2 identification and development process.

3

4 A recent longitudinal study examined the anthropometric and performance characteristics that
5 contributed to success in attaining full international appearances and/or a professional
6 contract in U14-U16 French National Soccer Academy players. Those players who did gain
7 an international appearance and/or a professional contract were taller and heavier and had a
8 faster 40 m sprint speed, higher jump height and faster flying 10 m sprint speed (measured
9 from the last 10 m of 40 m sprint) compared to players who did not gain a professional
10 contract (21). However, whether or not performance during match play differs between
11 players who at a later date are retained or released by academies has not been previously
12 examined and studies comparing match distances run for players of different standards are
13 only available for senior professional soccer players. Elite professional players have been
14 shown to cover a 5 % longer total distance and 28% greater high intensity running during a
15 match compared to moderate professional players (25). Thus the practical problem addressed
16 in the present study is whether or not match analysis data in very young players can
17 contribute to knowledge regarding which players at a later date may be retained by academies
18 and thus can knowledge regarding match distances covered contribute to the talent
19 identification and development process.

20

21 Therefore, the aims of the present study were: 1) to investigate the match performance of elite
22 U9 and U10 soccer players; and 2) to examine if there was a difference in the match
23 performance between retained and released youth soccer players. Based on the information
24 available from previous research on older players, it was hypothesized that the U9 players
25 would cover a shorter distance during match play than the U10 players and that the retained

- 1 players would cover a longer total distance and a longer distance at high speeds than released
- 2 players during a match play.

ACCEPTED

1 METHODS

2 Experimental approach to the problem

3 As there was no match analysis data available at all for any standard of U9 and U10 players,
4 an elite population was selected for the present study. The data was thought to be of value for
5 clubs attempting to develop players to professional soccer standard and would provide a
6 valuable comparison for later studies examining players competing at a lower level.

7 Therefore, players were recruited from an English Premier League Academy, which
8 represents the highest standard of youth development in England.

9
10 Match running performance of the U9 and U10 players and the same group of players divided
11 into two groups of those who were subsequently retained and released by the academy was
12 analyzed using a 1Hz GPS. The distances measured were categorized into speed zones which
13 were walking, jogging, low speed running, moderate speed running and high speed running.
14 All players participated in a 10 m sprint test with a 5 m split time and the results were used to
15 calculate speed zones for each squad. The distances covered in each speed zone, the total
16 distance covered and the percentage of time in each speed zone during a match were
17 compared between the U9 and U10 squads to test the hypothesis that the U10s in comparison
18 with the U9s and the retained in comparison with the released players would cover a greater
19 total distance and a greater distance at high speeds.

20

21 The 1 Hz GPS has been previously shown to be both valid and reliable in terms of indicating
22 the distance and speed covered by games players. The validity of GPS (1 Hz, SPI elite,
23 GPSport, Australia) has been assessed using male and female adult games players (N = 9)
24 using a circuit (487 m). They completed 14 laps in a trial and the circuit involved moving at

1 different speeds including walking to sprinting and various agility runs (8.5-52.3 m). The
2 actual distance covered by the participants in the trial (6818.0 m) and the total distance
3 measured by the devices only differed by 2.5 m and there was less than a 1% difference in the
4 actual distance covered and the distance measured by the devices during agility runs (23). It
5 is encouraging that the study attempted to replicate the movement patterns observed in field
6 hockey, which has a similar movement pattern to soccer, and found very acceptable validity
7 at the speeds and patterns of movement tested. For the reliability, intra- and inter- receiver
8 coefficient of variation in 1 Hz GPS (SPI elite, GPSport, Australia) for the distance covered
9 at speeds up to $5 \text{ m}\cdot\text{s}^{-1}$ is excellent at $\sim 3\%$ or less in both linear and non-linear movements.
10 The coefficient of variation of 3% or less was also observed in linear movements at $7\text{-}8 \text{ m}\cdot\text{s}^{-1}$.
11 Whereas the intra- and inter receiver coefficient of variation during non-linear movements at
12 $7\text{-}8 \text{ m}\cdot\text{s}^{-1}$ was $\sim 5\%$ and $\sim 6\%$, respectively (14).

14 **Subjects**

15 The participants in the study were U9 (N = 22, age: 9.1 ± 0.2 years, height: 136.2 ± 5.0 cm,
16 body mass: 31.5 ± 3.6 kg, estimated chronological age at PHV: 12.8 ± 0.4 years (N = 19), 5
17 m sprint: 1.15 ± 0.04 s, 10 m sprint: 2.04 ± 0.07 s, Yo-Yo intermittent recovery test (level 1)
18 performance: 1413 ± 245 m) and U10 (N = 12, age: 9.8 ± 0.4 years, height: 143.0 ± 5.7 cm,
19 body mass: 36.6 ± 4.2 kg, estimated chronological age at PHV: 13.3 ± 0.2 years (N = 10), 5
20 m sprint: 1.14 ± 0.04 s, 10 m sprint: 2.03 ± 0.08 s, Yo-Yo intermittent recovery test (level 1)
21 performance: 1427 ± 316 m) outfield players from an English Premier League Academy in
22 the Midlands and they had at least 1 year of experience in soccer training and competition.
23 There were 7 defenders, 3 midfielders, 7 strikers and 5 multi-position players in the U9 squad
24 and 5 defenders, 4 midfielders, 2 strikers and 1 multi-position player in the U10 squad. When
25 the players were separated into retained and released groups, there were 14 players in the

1 retained group (age: 9.3 ± 0.4 years, height: 138.6 ± 4.9 cm, body mass: 32.6 ± 3.9 kg,
2 estimated chronological age at PHV: 12.9 ± 0.4 years, 5 m sprint: 1.14 ± 0.04 s, 10 m sprint:
3 2.02 ± 0.08 s, Yo-Yo intermittent recovery test (level 1) performance: 1431 ± 256 m) and 20
4 players in released group (age: 9.4 ± 0.5 years, height: 138.5 ± 7.0 cm, body mass: 33.7 ± 5.0
5 kg, estimated chronological age at PHV: 13.1 ± 0.4 years (N = 15), 5 m sprint: 1.15 ± 0.04 s,
6 10 m sprint: 2.05 ± 0.07 s, Yo-Yo intermittent recovery test (level 1) performance: $1408 \pm$
7 258 m). The players in the retained group were retained in an English Premier League
8 Academy for at least two more seasons after the season in which the match analysis was
9 completed. Within the released group, four players were released at the end of the season in
10 which the match analysis took place, nine players were released during or at the end of the
11 first season after the match analysis took place and seven players were released during or at
12 the end of the second season after the match analysis took place. The players generally
13 participated in three 1.5 hour technical training sessions (no physical conditioning or strength
14 or power training) and one match per week during the season. Two coaches were in charge of
15 each training session and they held UEFA (Union of European Football Associations) 'A' or
16 'B' coaching licenses. Players were provided with a written and verbal explanation of the
17 study including all tests and measurements to be taken. Each player signed an assent form
18 and completed a health screen questionnaire prior to participation in the study. Players'
19 parent, guardian or care-giver signed the consent form prior to the start of the study. Players
20 were free to withdraw from the study without giving any reasons and without any penalty
21 regarding their academy position and these were explained to them verbally and in a written
22 format. Players were withdrawn from a particular test if they did not have a satisfactory
23 health status. The study was approved by Loughborough University Ethical Committee.
24

1 **Biological maturity**

2 The estimated chronological age at PHV of the players was determined using the
3 chronological age, standing height, sitting height and body mass of each player (24) and the
4 assessment took place in the May of the season 08-09 and 09-10.

5

6 **Sprint test**

7 A “flying” 5 m sprint time was obtained on an indoor new generation synthetic sports turf in
8 the September of the season 08-09 and 09-10. A photoelectric timing gate (Brower timing
9 system, Utah, USA) was placed at 0, 5 and 10 m and, the time was recorded nearest to 0.01 s.
10 The players sprinted from 1 m behind the first timing gate with their preferred foot front. No
11 backward movements were allowed just before initiating the sprint. Each player completed
12 three sprints and the fastest sprint time was selected for the calculation of speed zones.

13

14 **Yo-Yo intermittent recovery test (level 1)**

15 The test was conducted on an indoor new generation synthetic sports turf in the May of the
16 season 08-09 and 09-10. The test consisted of repeated running of 2 x 20 m back and forth
17 between 2 lines. The timing was notified by audio bleeps from a CD player and the time
18 allowed to complete each run was progressively shortened. The participants had a 10 s active
19 recovery period which consisted of 2 x 5 m jogs in between running bouts. Participants were
20 withdrawn from the test when they failed to reach the finish line in time twice during the test
21 (20). The distance covered during the test was recorded and used as the test result. An intra-
22 class correlation of 0.93 and coefficient of variation of 8.1% were reported in 28 adult males
23 (20). Moreover, the coefficient of variation was 13% in seventeen 8-9 year old sedentary
24 children (2).

1 **Match analysis**

2 Soccer matches were analyzed during the season 08-09 and 09-10 using a 1 Hz GPS (SPI
3 Elite, GPSport, Australia). The match analysis took place in various parts of the season
4 (September/October, December/January, February/March and May) to reflect the variation in
5 match performance during the season (30). This system required players to wear a small
6 backpack on their back which contained the device; players wore this equipment throughout
7 the match. The matches were 6-a-side and were played on a flat grass pitch (dimension: 44.8
8 x 26.0 m, penalty area: 9.0 x 18.8 m, area per player: 97.1 m²). The matches were part of the
9 regular series of inter-academy matches between Premier League Academies during a season
10 and they were played at ~11am on Sundays with match duration of 60 or 70 min (15 min x 4
11 or (20 min x 2 + 15 min x 2)). The players consumed ad libitum water, sports drinks and/or
12 small snacks (sports bar, banana etc). Completion of at least a half of the duration of a match
13 in two separate matches (mean \pm SD = 2.9 \pm 0.7 matches: range = 2-4 matches) was the
14 criterion for inclusion in the study. Mean values from matches were calculated for each
15 player. The GPS was accessing a mean of 7.6 \pm 1.3 satellites with a mean horizontal dilution
16 of precision of 1.20 \pm 0.39 throughout all the matches analyzed. The coefficient of variation
17 for match performance (m·h⁻¹), reflecting the variation between matches for each player, was
18 6.7 \pm 4.4% for total distance covered, 6.1 \pm 3.6% for walking distance, 8.0 \pm 6.5% for
19 jogging distance, 13.8 \pm 9.7% for low speed running distance, 18.1 \pm 12.7% for moderate
20 speed running distance and 25.8 \pm 13.1% for high speed running distance (31).

21

22 **Match activities**

23 Sprint speeds over 5 m calculated from “flying” 5 m sprint time were averaged for each
24 squad and five speed zones specific to each squad were calculated based on a mean of
25 average “flying” sprint speed over 5 m from each squad (walking, jogging, low speed running,

1 moderate speed running and high speed running). “Flying” 5 m sprint time from the 10 m
2 sprint test was used to calculate average sprint speed because the average sprint distance of
3 the U15 elite Brazilian players was 8.6 m when stride length was used to estimate the
4 distance covered by sprinting (26). The five speed categories were calculated by dividing the
5 speed zones of $0.0 \text{ m}\cdot\text{s}^{-1}$ to two standard deviations below the squad mean of average “flying”
6 sprint speed over 5 m into five equal categories. Running speeds faster than the fastest speed
7 zone were also included in the high speed running category (table 1). The distances covered
8 by the five locomotor categories were estimated using Team AMS software version 1.2
9 (GPSport, Australia) and they were presented in meters per hour ($\text{m}\cdot\text{h}^{-1}$). Also, mean playing
10 time, total distance covered during a match in absolute terms and in $\text{m}\cdot\text{h}^{-1}$ and, the percentage
11 of time spent in each speed zone during a match were calculated. These speed zones are
12 similar to the zones described in previous studies which reported the match performance of
13 U12-U16 youth soccer players (11, 12, 15).

14 ****Table 1 here****

15

16 **Statistical analyses**

17 A Kolmogorov-Smirnov test was employed to examine whether or not the distribution was
18 normal and homogeneity of variance was examined using Levene’s test. An independent
19 sample t-test was used to compare differences between the U9 and U10 squads and
20 differences between retained and released groups. The effect sizes (d) for these differences
21 were also calculated as $(\text{mean A} - \text{mean B}) / (\text{pooled SD})$. Effect size values of 0.2, 0.5 and
22 above 0.8 were considered to represent a small, moderate and large difference, respectively
23 (31). A Pearson’s product moment correlation was employed to examine the relationships
24 between variables.

25

1 A-priori statistical power calculations were conducted on primary outcome variables (i.e.,
2 total distance covered during a match, high speed running distance). Assuming an
3 independent t-test was the method of analysis, for total distance covered during a match, the
4 a-priori calculation suggested a minimum of 27 subjects (with at least 12 in a group) would
5 be needed to detect differences between 2 groups, assuming a statistical power of 80%, an
6 effect size of 0.9 (15, 25) and an alpha level of 0.05. For high speed running distance, the a-
7 priori calculation suggested that a minimum of 26 subjects (with at least 12 in a group) would
8 be required to detect differences between 2 groups, assuming a statistical power of 80%, an
9 effect size of 1.1 (25) and an alpha level of 0.05 (31).
10
11 The level of statistical significance was set at $p < 0.05$. Results are presented as mean \pm
12 standard deviation (SD) and PASW 18.0 was used for all the statistical analyses.

1 RESULTS

2 Match activities of U9 and U10 squads

3 Playing time

4 Mean playing time during a match was longer for the U9 squad compared to the U10 squad
5 (U9 vs. U10: 55.9 ± 4.3 min vs. 50.9 ± 5.2 min, $p < 0.01$)

6 Distance covered

7 Distances covered during a match are presented in table 2 and distances covered per hour of a
8 match are presented in figure 1. There were no differences in the total distance covered
9 during a match for the U9 and U10 squads (U9 vs. U10: 4356 ± 478 m vs. 4056 ± 541 m, $p =$
10 N.S.). Similarly, when the total distance covered during a match was standardized into meters
11 per hour, there were no differences in total distance covered by the U9 and U10 squads (U9
12 vs. U10: 4675 ± 311 m·h⁻¹ vs. 4788 ± 466 m·h⁻¹, $p =$ N.S.). However, there was a tendency for
13 the U10 squad to cover a longer distance (m·h⁻¹) than the U9 squad during moderate speed
14 running (U9 vs. U10: 496 ± 108 m·h⁻¹ vs. 576 ± 170 m·h⁻¹, $p = 0.10$, $d = 0.62$) and high speed
15 running (U9 vs. U10: 178 ± 56 m·h⁻¹ vs. 219 ± 70 m·h⁻¹, $p = 0.08$, $d = 0.69$).

16

17 ****Table 2 here****

18 ****Figure 1 here****

19

20 Percentage time

21 The U9 squad spent a lower percentage of time jogging compared to the U10 squad (U9 vs.
22 U10: $33.2 \pm 2.6\%$ vs. $36.5 \pm 2.8\%$, $p < 0.01$) and the U9 squad spent a higher percentage of

1 time in low speed running compared to the U10 squad (U9 vs. U10: $15.0 \pm 2.4\%$ vs. $12.6 \pm$
 2 2.7% , $p < 0.05$). No significant differences between the U9 and U10 squads were found in
 3 percentage of time spent in standing and walking (U9 vs. U10: $46.4 \pm 4.3\%$ vs. $44.8 \pm 5.8\%$,
 4 $p = \text{N.S.}$), moderate speed running (U9 vs. U10: $4.1 \pm 0.9\%$ vs. $4.8 \pm 1.4\%$, $p = \text{N.S.}$) and
 5 high speed running (U9 vs. U10: $1.1 \pm 0.3\%$ vs. $1.3 \pm 0.4\%$, $p = \text{N.S.}$).

6

7 ****Figure 2 here****

8

9 **Match activities of retained and released players**

10 **Playing time**

11 There were no differences in mean playing time during a match between retained and
 12 released groups (retained vs. released: 55.4 ± 4.2 min vs. 53.2 ± 5.6 min, $p = \text{N.S.}$).

13 **Distance covered**

14 Distances covered during a match are presented in table 3 and distances covered per hour of a
 15 match are presented in figure 3. Retained players covered a greater distance than released
 16 players (retained vs. released = 4478 ± 513 m vs. 4091 ± 462 m, $p < 0.05$) during a match and
 17 covered a greater distance during low speed running in absolute terms (retained vs. released =
 18 1226 ± 259 m vs. 1005 ± 221 m, $p < 0.05$). When the match running distances were
 19 standardized into meters per hour of a match, the retained players tended to cover a longer
 20 total distance (retained vs. released: 4844 ± 313 $\text{m}\cdot\text{h}^{-1}$ vs. 4624 ± 387 $\text{m}\cdot\text{h}^{-1}$, $p = 0.09$, $d =$
 21 0.63) and showed a longer low speed running distance (retained vs. released: 1325 ± 235 $\text{m}\cdot\text{h}^{-1}$
 22 1 vs. 1132 ± 210 $\text{m}\cdot\text{h}^{-1}$, $p < 0.05$) than the released players. However, no differences in
 23 distances covered by walking (retained vs. released = 1018 ± 75 $\text{m}\cdot\text{h}^{-1}$ vs. 1040 ± 83 $\text{m}\cdot\text{h}^{-1}$, p
 24 = N.S.), jogging (retained vs. released = 1760 ± 111 $\text{m}\cdot\text{h}^{-1}$ vs. 1734 ± 217 $\text{m}\cdot\text{h}^{-1}$, $p = \text{N.S.}$),

1 moderate speed running (retained vs. released = $545 \pm 121 \text{ m}\cdot\text{h}^{-1}$ vs. $510 \pm 147 \text{ m}\cdot\text{h}^{-1}$, $p =$
2 N.S.) and high speed running (retained vs. released = $195 \pm 67 \text{ m}\cdot\text{h}^{-1}$ vs. $191 \pm 63 \text{ m}\cdot\text{h}^{-1}$, $p =$
3 N.S.) were observed.

4

5 ****Table 3 here****

6 ****Figure 3 here****

7

8 **Percentage time**

9 In percentage of time spent, the retained group tended to spend a lower percentage of time
10 standing and walking (retained vs. released: $44.2 \pm 3.6\%$ vs. $47.0 \pm 5.3\%$, $p = 0.10$, $d = 0.61$)
11 and spent a higher percentage of time undertaking low speed running (retained vs. released:
12 $15.4 \pm 2.8\%$ vs. $13.3 \pm 2.4\%$, $p < 0.05$) compared to the released group. However, there were
13 no differences in percentage of time spent in jogging (retained vs. released: $34.5 \pm 1.9\%$ vs.
14 $34.3 \pm 3.7\%$, $p = \text{N.S.}$), moderate speed running (retained vs. released: $4.5 \pm 1.0\%$ vs. $4.3 \pm$
15 1.2% , $p = \text{N.S.}$) and high speed running (retained vs. released: $1.2 \pm 0.4\%$ vs. $1.2 \pm 0.4\%$, $p =$
16 N.S.) between the groups.

17

18 ****Figure 4 here****

19

20 **Match performance and Yo-Yo intermittent recovery test**

21 There was a positive relationships between the Yo-Yo intermittent recovery test performance
22 ($N = 34$) and the total distance covered ($r = 0.36$, $p < 0.05$) and the distance covered at
23 moderate speeds ($r = 0.49$, $p < 0.01$).

1 **DISCUSSION**

2 The aim of this study was, for the first time, to examine the match speeds and distances
3 covered for U9 and U10 English Premier League Academy soccer players and to examine if
4 differences existed between players subsequently retained or released by the academy. The
5 main findings were that the total distance covered during a match was ~4000 m in absolute
6 terms or ~4700 m·h⁻¹ for the U9 and U10 squads. When the players were separated into
7 retained and released players, the retained players covered a significantly longer total
8 distance (by ~400 m) during a match and a significantly greater distance at low speed running
9 (by ~200 m) than the released players. The information concerning match distances covered
10 for the U9 and U10 players and the differences in the distances covered between retained and
11 released players should prove valuable for football and conditioning coaches in the design of
12 training programs and in the talent identification and development process. It will be of value
13 for the players themselves to know that a high work rate may increase the chances of
14 retention within the academy system.

15
16 All of the match analysis data in the present study are novel as the match analysis of U9 and
17 U10 players has not been previously reported upon nor has match analysis data been
18 presented for players who were subsequently retained or released by clubs. The U9 and U10
19 squads from the current study covered 4356 and 4056 m during a match respectively, with no
20 significant differences between the squads. However, the U9 players did have a significantly
21 longer playing time compared to the U10 players as the U9 squad contained less players and
22 thus each player received slightly more playing time. A previous study showed that the U12
23 squads from two English professional soccer clubs covered an average of 5967 m during an
24 11-a-side match (15) and this was around 30% more than the distance covered by the U9 and

1 U10 squads from the current study. When the total distance covered during a match was
2 standardized per hour of a match, the U9 and U10 squads from the current study covered
3 4675 and 4788 $\text{m}\cdot\text{h}^{-1}$, respectively whereas the U12 players from the English professional
4 soccer clubs covered 5978 $\text{m}\cdot\text{h}^{-1}$ (15) which suggests a difference of around 1200 m in
5 distance covered per hour of a match between the U10 and U12 squads from the current and
6 previous studies. This is a large difference between the two age groups (between U10 and
7 U12) as the total distance covered per hour of a match has been previously reported to
8 increase by only ~ 1200 m over 4 year groups (5928 $\text{m}\cdot\text{h}^{-1}$ for U12 to 7122 $\text{m}\cdot\text{h}^{-1}$ for U16) in
9 English professional soccer clubs (15). This large difference in total distance covered during
10 a match between the U10 squad reported here and the U12 squad from the earlier study was
11 possibly due to a mixture of the differences in the number of players on a team and the pitch
12 dimensions (18, 27) as well as due to the enhanced physical performance of the older players
13 in terms of both speed and endurance (19). In the previous study (15), an 11-a-side match was
14 played as opposed to a 6-a-side match in the current study and the pitch dimensions were 77
15 x 60 m (area per player = 210 m^2) in the earlier study and 44.8 x 26.0 m (area per player = 97
16 m^2) in the present study, respectively. It has been shown that the number of players does not
17 influence the total distance covered during a match when the area per player was the same
18 (18). However, a match with a greater area per player has been shown to provide a higher
19 relative heart rate, blood lactate concentration and perceived exertion regardless of player
20 number (27). Hence, the large differences in total distances covered between the U9 and U10
21 players from the current study and the U12 players from the previous study (15) may be due
22 at least in part to a difference in the area per player.

23

24 Another potential reason for the large difference in total distance covered between the U9 and
25 U10 players from the current study and the U12 players from the previous study (15) is the

1 use of a GPS with a different manufacturer and frequency between the studies (current: 1 Hz
2 SPI Elite, GPSport, Australia vs. previous: 5 Hz MinimaxX, Catapult Innovations, Australia).
3 Previously, when match analysis data from a 1 Hz and 5 Hz GPS worn during the same
4 match was compared, the 5 Hz GPS estimated a 12% longer total distance covered during a
5 match than the 1 Hz GPS (28). However, this difference in the sampling frequencies of GPS
6 between studies only accounts for a fraction of the difference in the distances run between the
7 U9/U10 players in this study and U12 players in other studies. Thus it seems likely there is a
8 real physiological difference in performance between the U9/U10 players and the U12
9 players in other studies, possibly due to improvements in endurance fitness with age, as
10 supported by the slightly higher (NS) Yo-Yo test performance of the U10 in comparison with
11 U9 boys in the present study. However, 4000 m already represents a substantial distance
12 covered for these very young boys with a short stride length because of their height and
13 coaches may consider whether or not a small supplementary training program in addition to
14 technical work might be included for this age group, at least as a preparation for the fitness
15 work that is to follow in older age group squads.

16

17 During a match, the U9 and U10 squads covered 166 and 186 m by high speed running and
18 462 and 485 m by moderate speed running, respectively. When these distances were
19 expressed relative to an hour of a match, the U10 squad showed a trend to cover a longer
20 distance by high speed running and moderate speed running compared to the U9 squad even
21 though the speed zones for high speed running and moderate speed running were slightly
22 faster for the U10 squad. This may suggest that there is a development in the ability to
23 complete high intensity running during a match between the U9 and U10 squads in this
24 English Premier League Academy club. This is possibly because of a development in speed,
25 agility, jump height and endurance with age (19) and such findings suggest that each age

1 group requires a specific training program. Moreover, if a coach moves an U9 age player into
2 the U10 squad, the player needs to be able to cope with the match demands of the U10 squad.

3

4 The U10 squad jogged more than the U9s (distance covered and percentage of match time)
5 whereas the U9s spent more time in low speed running (faster than jogging) than the U10s.

6 This is possibly because the U10 squad was required to spend a longer time performing lower
7 intensity exercise to recover compared to the U9 squad as the U10 squad tended to cover a
8 longer distance than the U9 squad during moderate speed running and high speed running.

9 The U9 squad spent more time in low speed running possibly because they tended to cover
10 less distance in higher intensity exercises (moderate speed running and high speed running).

11 Moreover, the U9 squad walked a significantly longer distance than the U10 squad during a
12 match. However, this was due to the U9 players having a significantly longer playing time
13 compared to the U10 squad as, when the distance walked was standardized into per hour of a
14 match, the difference no longer existed.

15

16 The percentage of time spent in moderate speed running during a match was 4.1 and 4.8 %
17 for the U9 and U10 squads, respectively. For high speed running, the percentage of time
18 spent during a match was 1.1 and 1.3 % for the U9 and U10 squads, respectively. Using
19 similar speed zones (based on senior player sprint performance), the English FA Premier
20 (Senior) League soccer players spent 6.4 and 2.6% of total match time in moderate speed
21 running and high speed running, respectively (4). These findings imply that the U9 and U10
22 elite soccer players may spend a lower proportion of match time in high intensity running
23 than top elite professional soccer players and this is possibly because of differences in the
24 area per player (1, 5, 27). Hence, for the U9 and U10 players to achieve a similar proportion
25 of time in each speed zone in comparison to the elite senior players, the pitch dimensions

1 could be increased. However, as a result of the potential enhancement in the development of
2 technical ability due to gaining more touches on the ball and because of the possible
3 undeveloped understanding of technical elements by young boys, 6-a-side matches may be
4 preferred by the coaches of U9 and U10 players (8, 10).

5

6 When the players were separated into retained and released players, the retained players
7 covered a significantly longer distance than the released players during a match and covered a
8 greater distance at low speeds (distance covered in absolute and relative terms and percentage
9 of match time). In senior professional players, total distance covered during a match has been
10 shown to distinguish the standard of soccer players (25). Moreover, 16 years old soccer
11 players who competed at international level had a higher maximal oxygen uptake than same-
12 age players from local and school teams (29) and endurance fitness has been shown to relate
13 to total distance covered during a match in 14 year old soccer players (11, 12) and in the
14 present study for the U9 and U10 players. The slightly higher (NS) Yo-Yo intermittent
15 recovery test results of the retained players suggest that possibly the retained players had a
16 higher peak oxygen uptake or enhanced endurance fitness in comparison with the released
17 players. Such a difference in endurance fitness would provide more frequent involvement
18 with the ball (17) in retained players which possibly offers more chances to improve technical
19 ability in the retained players compared to the released players. Hence, the provision of
20 endurance training to improve endurance ability may enhance the match running distance of
21 players increasing involvement in the game, enhancing opportunities to improve technical
22 ability (17) and thus improving the chances of being retained in the academy. Also it is
23 possible that the greater distances covered by the retained players reflects an attitude of mind
24 or enhanced reading of the game which enables them to be more fully engaged in match play.

25

1 However, no significant differences were found in high intensity running (high speed running
2 and moderate speed running) distances between retained and released players. This is an
3 interesting finding as high intensity running distance has been reported to distinguish the
4 standard of senior soccer players (25). This may be because the players from the current
5 study spent a lower proportion of match time in high intensity running compared to the senior
6 professional soccer players (4) possibly due to a difference in the area per player (27)
7 between the current and previous (4) studies or because the anaerobic energy supplying
8 pathways will not be fully developed in boys of this age (13, 16, 22).

9
10 In summary, the total distance covered during a match was approximately 4000 m for the U9
11 and U10 elite soccer players and only 600-700 m were covered by high intensity running
12 which was achieved during 5-6% of match playing time. The U10 squad tended to cover a
13 longer distance at moderate and high speeds compared to the U9 squad. This suggests that
14 there is a possible development in the ability to perform high intensity match running activity
15 from the U9 to the U10 squads. The players who were retained in the English Premier League
16 Academy covered a greater total distance and covered a greater distance at low speeds
17 compared to the players who were released. Thus, knowledge concerning the distances run in
18 young players may prove valuable in the talent identification and development process and
19 coaches could consider whether or not an introductory fitness training program targeting
20 match distance run might be appropriate in this age group for all or some players.

1 PRACTICAL APPLICATION

2 Despite their young age, the U9 and U10 boys in a professional football academy covered
3 over 4000 m in a match and the U10 boys tended to cover a greater distance at high and
4 moderate speeds than the U9 boys. Thus, coaches and sports scientists may be best advised to
5 create specific training programs for each age group. In addition, if an U9 player is moved up
6 into the U10 squad, coaches and sports scientists need to consider if that player can cope with
7 the physical demands of the older age group.

8
9 The U9 and U10 boys who were later retained by their academy covered a greater total match
10 running distance and a greater low speed running distance than those players that were later
11 released. Thus, the distances and speeds covered during match play for these age groups
12 could form an important part of the talent identification and development process and
13 coaches and sports scientists could consider recording match distances run occasionally
14 through the season. In addition, an introductory modest fitness program may enhance match
15 distances covered for players of these ages increasing their engagement with game and thus
16 enhancing the chances of improving their match-play. For players with low endurance fitness
17 enhancing this aspect of their performance may increase their chances of retention in an
18 academy.

1 REFERENCES

- 2 1. Abrantes, CI, Nunes, MI, Maçãs, VM, Leite, NM, and Sampaio JE. Effects of the
3 number of players and game type constraints on heart rate, rating of perceived exertion,
4 and technical actions of small-sided soccer games. *J Strength Cond Res* 26: 976-981,
5 2012.
- 6 2. Ahler, T, Bendiksen, M, Krusturup, P, and Wedderkopp N. Aerobic fitness testing in 6- to
7 9-year-old children: reliability and validity of a modified Yo-Yo IR1 test and the
8 Andersen test. *Eur J Appl Physiol* 112: 871-876, 2012.
- 9 3. Bangsbo, J, Marcello, FI, and Krusturup, P. The Yo-Yo Intermittent Recovery Test: A
10 Useful Tool for Evaluation of Physical Performance in Intermittent Sports. *Sports*
11 *Medicine*, 38: 37-51, 2008.
- 12 4. Bradley, PS, Sheldon, W, Wooster, B, Olsen, P, Boanas, P, and Krusturup, P. High-
13 intensity running in English FA Premier League soccer matches. *J Sports Sci* 27: 159-
14 168, 2009.
- 15 5. Brandes, M, Heitmann, A, and Müller, LJ. Physical responses of different small-sided
16 game formats in elite youth soccer players. *J Strength Cond Res* 26: 1353-1360, 2012.
- 17 6. Buchheit, M, Mendez-Villanueva, A, Simpson, BM, and Bourdon, PC. Match running
18 performance and fitness in youth soccer. *Int J Sports Med* 31: 818-825, 2010.
- 19 7. Buchheit, M, Mendez-villanueva, A, Simpson, BM, and Bourdon, PC. Repeated-sprint
20 sequences during youth soccer matches. *Int J Sports Med* 31: 709-716, 2010.
- 21 8. Capranica, L, A. Tessitore, LG, and Figura, F. Heart rate and match analysis in pre-
22 pubescent soccer players. *J. Sports Sci* 19: 379-384, 2001.

- 1 9. Carling, C, Bloomfield, J, Nelsen, L, and Reilly, YT. The role of motion analysis in elite
2 soccer: contemporary performance measurement techniques and work rate data. *Sports*
3 *Med* 38: 839-862, 2008.
- 4 10. Castagna, C, D'Ottavio, S, and Abt, G. Activity profile of young soccer players during
5 actual match play. *J Strength Cond Res* 17: 775-780, 2003.
- 6 11. Castagna, C, Impellizzeri, F, Cecchini, E, Rampinini, E, and Barbero-Alvarez, JC.
7 Effects of intermittent-endurance fitness on match performance in young male soccer
8 players. *J Strength Cond Res* 23: 1954-1959, 2009.
- 9 12. Castagna, C, Manzi, V, Impellizzeri, F, Weston, M, and Barbero-Alvarez, JC.
10 Relationship between endurance field tests and match performance in young soccer
11 players. *J Strength Cond Res* 24: 3227-3233, 2010.
- 12 13. Eriksson, O, and Saltin, B. Muscle metabolism during exercise in boys aged 11 to 16
13 years compared to adults. *Acta Paediatrca Belgica* 28: 257-265, 1974.
- 14 14. Gray, AJ, Jenkins, D, Andrews, MH, Taaffe, DR, and Glover, ML. Validity and
15 reliability of GPS for measuring distance travelled in field-based team sports. *J Sports*
16 *Sci* 28: 1319-1325, 2010.
- 17 15. Harley, JA, Barnes, CA, Portas, M, Lovell, R, Barrett, S, Paul, D, and Weston, M.
18 Motion analysis of match-play in elite U12 to U16 age-group soccer players. *J Sports Sci*
19 28: 1391-1397, 2010.
- 20 16. Haralambie, G. Enzyme activities in skeletal muscle of 13-15 years old adolescents.
21 *Bulletin Europeen de Physiopathologie Respiratoire* 18: 65-74, 1982.
- 22 17. Helgerud, J, Engen, LC, Wisloff, U, and Hoff, J. Aerobic endurance training improves
23 soccer performance. *Med Sci Sport Exerc* 33: 1925-1931, 2001.

- 1 18. Hill-Haas, SV, Dawson, BT, Coutts, AJ, and Rowsell, GJ. Physiological responses and
2 time-motion characteristics of various small-sided soccer games in youth players. *J*
3 *Sports Sci* 27: 1-8, 2009.
- 4 19. Hulse, MA, Morris, JG, and Nevill, ME. A field-test battery for elite, young soccer
5 players. *Int J Sports Med (In Press)*, 2012.
- 6 20. Krstrup, P, Mohr, M, Amstrup, T, Rysgaard, T, Johansen, J, Steensberg, A, Pedersen,
7 PK, and Bangsbo, J. The Yo-Yo Intermittent Recovery Test: Physiological response,
8 reliability, and validity. *Med Sci Sport Exerc* 35: 697-705, 2003.
- 9 21. Le Gall, F, Carling, C, Williams, M, and Reilly, T. Anthropometric and fitness
10 characteristics of international, professional and amateur male graduate soccer players
11 from an elite youth academy. *J Sci Med Sport* 13: 90-95, 2010.
- 12 22. Lexell, J, Sjöström, M, Nordlund, AS, and Taylor, CC. Growth and development of
13 human muscle: a quantitative morphological study of whole vastus lateralis from
14 childhood to adult age. *Muscle Nerve* 15: 404-409, 1992.
- 15 23. MacLeod, H, Morris, JG, Nevill, A, and Sunderland, C. The validity of a non-differential
16 global positioning system for assessing player movement patterns in field hockey. *J*
17 *Sports Sci* 27: 2, 121-128, 2009.
- 18 24. Mirwald, RL, Baxter-Jones, AD, Bailey, DA, and Beunen, GP. An assessment of
19 maturity from anthropometric measurements. *Med Sci Sport Exerc* 34: 689-694, 2002.
- 20 25. Mohr, M, Krstrup, P, and Bangsbo, J. Match performance of high standard soccer
21 players with special reference to development of fatigue. *J Sports Sci* 21: 519-528, 2003.
- 22 26. Pereira Da Silva, N, Kirkendall, DT, and De Barros Neto, TL. Movement patterns in elite
23 Brazilian youth soccer. *J Sports Med Phys Fitness* 47: 270-275, 2007.

- 1 27. Rampinini, E, Impellizzeri, FM, Castagna, C, Abt, G, Chamari, K, Sassi, A, and Marcora,
2 SM. Factors influencing physiological responses to small-sided soccer games. *J Sports*
3 *Sci* 25: 659-666, 2007.
- 4 28. Randers, MB, Mujika, I, Hewitt, A, Santisteban, J, Bischoff, R, Solano, R, Zubilaga, A,
5 Peltola, E, Krustup, P, and Mohr, M. Application of four different football match analysis
6 systems: A comparative study. *J Sports Sci* 28: 171-182, 2010.
- 7 29. Reilly, T, Williams, AM, Nevill, A, and Franks, A. A multidisciplinary approach to talent
8 identification in soccer. *J Sports Sci* 18: 695-702, 2000.
- 9 30. Silva, JR, Magalhães, J, Ascensão, A, Seabra, AF, and Rebelo AN. Training status and
10 match activity of professional soccer players throughout a season. *J Strength Cond Res*
11 27: 20-30, 2013.
- 12 31. Vincent, WJ. *Statistics in Kinesiology* (3rd ed). Champaign, IL: Human Kinetics, 2005.
13

ACCEPTED

1 **Acknowledgement**

- 2 We would like to thank the academy staff for their support and the players for taking part in
3 the study. This study was supported by Loughborough University.

ACCEPTED

1 **FIGURE LEGEND**

2

3 **Figure 1.** Mean and standard deviation ($\text{m}\cdot\text{h}^{-1}$) of total distance covered and distance covered
4 by walking, jogging, low speed running, moderate speed running and high speed running in
5 an hour of a match in the U9 and U10 squads. *significantly different at $p < 0.05$. ^atended to
6 be different ($p = 0.08$). ^btended to be different ($p = 0.10$).

7

8

9 **Figure 2.** Mean and standard deviation (%) of percentage of time spent on standing and
10 walking, jogging, low speed running, moderate speed running and high speed running in a
11 match in the U9 and U10 squads. *significantly different at $p < 0.05$.

12

13

14 **Figure 3.** Mean and standard deviation ($\text{m}\cdot\text{h}^{-1}$) of total distance covered and distance covered
15 by walking, jogging, low speed running, moderate speed running and high speed running in
16 an hour of a match in the retained and released groups. *significantly different at $p < 0.05$.
17 ^atended to be different ($p = 0.09$).

18

19

20 **Figure 4.** Mean and standard deviation (%) of percentage of time spent on standing and
21 walking, jogging, low speed running, moderate speed running and high speed running in a
22 match in the retained and released groups. *significantly different at $p < 0.05$. ^atended to be
23 different ($p = 0.10$).

1 **TABLES**2 **Table 1.** The speed zones for match analyses of the U9 and U10 squads presented in m·s⁻¹.

	U9	U10
Standing and walking	0.0 - 1.0	0.0 - 1.0
Jogging	1.1 - 2.0	1.1 - 2.1
Low speed running	2.1 - 3.1	2.2 - 3.1
Moderate speed running	3.2 - 4.1	3.2 - 4.2
High speed running	> 4.1	> 4.2

3

4

5

6

7

8

9

10

11

12

13

14

15

16

17

18

19

20

21

22

23

24

25

26

27

1 **Table 2.** Mean and standard deviation (m) of total distance covered and distance covered by
 2 walking, jogging, low speed running, moderate speed running and high speed running in a
 3 match from the U9 and U10 squads.

		U9	U10
Total (m)	Mean	4356	4056
	SD	478	541
Walking (m)	Mean	966*	865
	SD	89	131
Jogging (m)	Mean	1560	1594
	SD	207	229
Low speed running (m)	Mean	1189*	927
	SD	239	208
Moderate speed running (m)	Mean	462	485
	SD	109	139
High speed running (m)	Mean	166	186
	SD	52	60

4 *significantly different to the U10 squad at $p < 0.05$.

5

6

7

1

2

3 **Table 3.** Mean and standard deviation (m) of distance covered by walking, jogging, low
 4 speed running, moderate speed running and high speed running in a match by retained and
 5 released players.

		Retained	Released
Total (m)	Mean	4478*	4091
	SD	513	462
Walking (m)	Mean	938	925
	SD	82	135
Jogging (m)	Mean	1627	1534
	SD	190	223
Low speed running (m)	Mean	1226*	1005
	SD	259	221
Moderate speed running (m)	Mean	505	446
	SD	125	110
High speed running (m)	Mean	180	167
	SD	66	51

6 *significantly different to released group at $p < 0.05$.

7

8

Figure 1




Figure 2




Figure 3




Figure 4

