

SCIENTIFIC REPORTS



OPEN

Escherichia coli of sequence type 3835 carrying *bla*_{NDM-1}, *bla*_{CTX-M-15}, *bla*_{CMY-42} and *bla*_{SHV-12}

Received: 04 March 2015

Accepted: 22 June 2015

Published: 21 July 2015

Yu Feng^{1,2,*}, Ping Yang^{1,2,*}, Yi Xie³, Xiaohui Wang^{1,2}, Alan McNally⁵ & Zhiyong Zong^{1,2,4}

New Delhi metallo- β -lactamase (NDM) represents a serious challenge for treatment and public health. A carbapenem-resistant *Escherichia coli* clinical strain WCHEC13-8 was subjected to antimicrobial susceptibility tests, whole genome sequencing and conjugation experiments. It was resistant to imipenem (MIC, >256 μ g/ml) and meropenem (MIC, 128 μ g/ml) and belonged to ST3835. *bla*_{NDM-1} was the only carbapenemase gene detected. Strain WCHEC13-8 also had a plasmid-borne AmpC gene (*bla*_{CMY-42}) and two extended-spectrum β -lactamase genes (*bla*_{CTX-M-15} and *bla*_{SHV-12}). *bla*_{NDM-1} and *bla*_{SHV-12} were carried by a 54-kb IncX3 self-transmissible plasmid, which is identical to plasmid pNDM-HF727 from *Enterobacter cloacae*. *bla*_{CMY-42} was carried by a 64-kb IncI1 plasmid and *bla*_{CTX-M-15} was located on a 141-kb plasmid with multiple F replicons (replicon type: F36:A4:B1). *bla*_{CMY-42} was in a complicated context and the mobilisation of *bla*_{CMY-42} was due to the transposition of *ISEcp1* by misidentifying its right-end boundary. Genetic context of *bla*_{NDM-1} in strain WCHEC13-8 was closely related to those on IncX3 plasmids in various *Enterobacteriaceae* species in China. In conclusion, a multidrug-resistant ST3835 *E. coli* clinical strain carrying *bla*_{NDM-1}, *bla*_{CTX-M-15}, *bla*_{CMY-42} and *bla*_{SHV-12} was identified. IncX3 plasmids may be making a significant contribution to the dissemination of *bla*_{NDM} among *Enterobacteriaceae* in China.

Carbapenems have long served as reliable and potent agents against Gram-negative bacilli. However, a variety of enzymes produced by bacteria are able to hydrolyse carbapenems, termed carbapenemases. New Delhi metallo- β -lactamase (NDM) is a type of carbapenemase and confers resistance to all β -lactams except monobactams¹. NDM was first found in a *Klebsiella pneumoniae* isolate in 2008² and since then bacterial isolates producing NDM have been identified worldwide, representing a serious challenge for treatment, infection control and public health. In addition to India, China is a common place from which NDM-producing isolates are identified. In China, NDM was initially seen in *Acinetobacter* spp³, but NDM-producing *Enterobacteriaceae* have been increasingly reported⁴. Among carbapenem-resistant *Enterobacteriaceae* in China, *K. pneumoniae* is the most common species with 10% of 12,121 *K. pneumoniae* clinical isolates collected from 16 major teaching hospitals in 2013 resistant to imipenem⁵. In contrast, carbapenem resistance is relatively uncommon in *Escherichia coli* in China and only 1% of 16,794 *E. coli* clinical isolates from the 16 hospitals in 2013 were resistant to imipenem⁵. During routine clinical duties, a carbapenem-resistant *E. coli* clinical isolate was encountered. This strain was then characterised and is reported here.

¹Center of Infectious Diseases, West China Hospital, Sichuan University, Chengdu, China. ²Division of Infectious Diseases, State Key Laboratory of Biotherapy, Chengdu, China. ³Laboratory of Clinical Microbiology, Department of Laboratory Medicine, West China Hospital, Sichuan University, Chengdu, China. ⁴Department of Infection Control, West China Hospital, Sichuan University, Chengdu, China. ⁵Pathogen Research Group, Nottingham Trent University, Nottingham, UK. *These authors contributed equally to this work. Correspondence and requests for materials should be addressed to Z.Z. (email: zongzhiy@scu.edu.cn)

Results and Discussion

Clinical strain WCHEC13-8 was recovered on November 2012. This strain was identified as *E. coli* and was resistant to imipenem (Minimum inhibitory concentration [MICs], >256 µg/ml), meropenem (MIC, 128 µg/ml), ceftazidime (MIC, >256 µg/ml), ciprofloxacin (MIC, 256 µg/ml) but was intermediate to amikacin (MIC, 32 µg/ml) determined by the microdilution method. It was also resistant to ampicillin-sulbactam, aztreonam, cefazolin, cefepime, cefotaxime, ceftiofloxacin, gentamicin, piperacillin-tazobactam, tobramycin and trimethoprim-sulphamethoxazole determined by Vitek II. *bla*_{NDM-1} was the only carbapenemase-encoding gene that was detected and sequencing the complete coding sequence of the gene revealed that it was *bla*_{NDM-1}.

Strain WCHEC13-8 was subjected to whole genome sequencing on the Illumina HiSeq platform. A total of 7,149,960 reads and 643,496,400 bases were obtained from the genome sequencing with a 49.8% GC content. Reads were assembled to 268 contigs that were ≥500 bp in length (N50 metric, 118,912 bp) and contained 5,145,308 bp nucleotides. To identify whether strain WCHEC13-8 had any new genes or not, its genome sequence was compared to other *E. coli* genome sequences including 83 complete and 2,159 draft genomes available in GenBank. Despite comparing with such a large number of *E. coli* genomes, the strain still had three unique genes that were located on contigs belonging to the chromosome. Two of the three genes were clustered together and resided in the centre of a 149-kb large contig, which was part of the chromosome as it shows significant identity (74% coverage and 97% identity) with the chromosome of *E. coli* strain E2348/69 (GenBank accession number FM18568) identified using the BLAST program. The two novel genes encoded hypothetical proteins with 420 and 346 amino acids, respectively. Unfortunately, the function of the two hypothetical proteins were not able to be predicted using both Protein BLAST and InterProScan programs. The remaining unique gene was 2,976 bp on a small 3,322 bp contig and contained both an endonuclease/exonuclease/phosphatase and a reverse transcriptase domain identified using the InterProScan program.

Strain WCHEC13-8 belonged to the phylogenetic group A and a new Sequence type (ST), ST3835 (*adk-fumC-gyrB-icd-mdh-purA-recA* allele numbers, 10-4-4-411-8-13-73). Multiple ST3835 isolates have been found in Korea in 2013, which is later than strain WCHEC13-8 (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>). ST3835 has a single allele different from ST1284 (10-4-4-8-8-13-73) and ST3835 *E. coli* has been recovered from a human in Spain and from a dog in Germany (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>). ST3835 belongs to the ST10 complex. Besides ST3835, STs belonging to the ST10 complex (ST40, ST167, ST205, ST744 and ST1237), ST88 complex (ST224 and ST410), ST101 complex (ST101) or ST361 complex (ST361) have been found to carry *bla*_{NDM-1} in China^{4,6-9}. The diversity of clonal background of *E. coli* carrying *bla*_{NDM-1} suggests that the dissemination of *bla*_{NDM-1} in China was unlikely to be mainly mediated by a particular strain.

In addition to *bla*_{NDM-1}, strain WCHEC13-8 had other resistance genes including *bla*_{CTX-M-15} (an extended-spectrum β-lactamase [ESBL] which is globally distributed), *bla*_{SHV-12} (an ESBL gene), *bla*_{CMY-42} (a plasmid-borne AmpC cephalosporinase gene), *bla*_{OXA-1} (a non-ESBL oxacillinase gene), *bla*_{ampC} (a chromosome-based AmpC gene), *aac(6′)-Ib-cr* (encoding an aminoglycoside acetyltransferase with low-level activity against fluoroquinolones), *aac(3)-II* (encoding an aminoglycoside acetyltransferase), *mph* (encoding macrolide 2′-phosphotransferase I) and *tetB* (conferring resistance to tetracycline). Of note, CMY-42, the AmpC enzyme encoded by *bla*_{CMY-42}, differs from CMY-2 by a single amino acid substitution, i.e. Ser for Val at Ambler's position 211¹⁰. The co-existence of *bla*_{NDM-1}, *bla*_{CMY-42}, *bla*_{CTX-M-15} and *bla*_{SHV-12}, all of which encode a β-lactamase with the ability to hydrolyse broad-spectrum cephalosporins, in a single isolate has not been reported before. The co-presence of *bla*_{NDM-1}, *bla*_{CMY-42} and *bla*_{CTX-M-15} has been found in an *E. coli* of ST101 from a two-year-old child who had recently travelled to India¹¹, while two *E. coli* isolates of ST167 in China have been found carrying *bla*_{NDM-1}, *bla*_{CMY-42} and *bla*_{CTX-M-14}⁶. In addition, *bla*_{CMY-42} has also been found in two *E. coli* isolates from a river in India (GenBank accession numbers KJ661335 and KJ661336) and one from a surgical wound at a university hospital in Germany¹⁰, but STs of the three *E. coli* isolates have not been reported. Of note, a report from Egypt also claimed that *bla*_{CMY-42} has been detected in 8 *E. coli* isolates¹², but the gene detected was 99% similar to *bla*_{CMY-42} and the amino acid sequence of CMY was not available for analysis. Therefore, it remains unclear whether these Egyptian isolates carried *bla*_{CMY-42} or another variant.

Strain WCHEC13-8 had three plasmids, which were approximately 50, 60 and 140 kb in size, as revealed by S1 nuclease pulse-field gel electrophoresis (S1-PFGE). The three plasmids were completely circularised using PCR and Sanger sequencing. *bla*_{NDM-1} and *bla*_{SHV-12} were carried by a 54-kb plasmid, assigned pNDM1_EC8 here, of the IncX3 group. *bla*_{CMY-42} was located on a 64-kb IncI1 plasmid, assigned pCMY42_EC8 here, while *bla*_{CTX-M-15} was carried by a 141-kb IncF plasmid, assigned pCTXM15_EC8.

pCTXM15_EC8 contained multiple F replicons including one FIA, one FIB and two FII replicons. Based on the replicon sequence typing (RST) for IncF plasmids¹³, the replicon types of pCTXM15_EC8 were F36:A4:B1 (FII 36, FIA 4 and FIB 1) with both FII replicons belonging to the same type. The F36:A4:B1 type has also been seen in plasmid p6409-151.583 kb (GenBank accession number CP010372), which was recovered from an *E. coli* strain in Colombia. However, p6409-151.583 kb has an additional F31 type FII replicon and does not carry *bla*_{CTX-M}. Furthermore, the link between pCTXM15_EC8 and p6409-151.583 kb could not be established and the origin of pCTXM15_EC8 remains unknown.

pCMY42_EC8 harboured four of the five loci used for the IncI1 plasmid MLST scheme¹⁴. The four loci and their allele numbers were *repI1* (IncI1 replicon) 4, *ardA* (alleviation of restriction of DNA) 5,

trbA (a gene essential for conjugative transfer) 15 and *pilL* (a pili gene) 3. The *sogS* (suppressor of *dnaG* mutation) locus was absent from pCMY42_EC8 and therefore no ST could be assigned. Nonetheless, pCMY42_EC8 was closely related to ST55 (*rep11-ardA-trbA-sogS-pilL* alleles 4-5-15-11-3) and ST141 (4-5-15-4-3). Both STs comprise Inc11 plasmids carrying *bla*_{CMY-42} found in Taiwan (pubmlst.org/plasmid).

The genetic context of *bla*_{CMY-42} has not been described before. On pCMY42_EC8, *bla*_{CMY-42} is adjacent to the insertion sequence *ISEcp1*, which was truncated by the insertion of *IS1*, at upstream and a gene encoding outer membrane lipoprotein at downstream (Panel A, Fig. 1). The remaining part of *ISEcp1* was found 24.7-kb further upstream of *bla*_{CMY-42} and was also truncated by *IS1*. After carefully examining the flanking sequences of *ISEcp1* and *IS1*, it became evident that the complex genetic context of *bla*_{CMY-42} has been formed by the transposition of *ISEcp1* and the insertion of two copies of *IS1* followed by homologous recombination between the two *IS1* (Panel B, Fig. 1) as explained as below. First, both of the two *IS1* were flanked by a remnant of *ISEcp1* and a part of the *traB* gene, which is involved in plasmid conjugation. The two remnants of *ISEcp1* form a complete *ISEcp1* plus a 9-bp repeat, which abuts the two *IS1* and is characteristic of the insertion of *IS1*. Similarly, the two parts of *traB* make the complete *traB* plus the characteristic 9-bp repeat. It is therefore clear that both *ISEcp1* and *traB* had been interrupted by the insertion of *IS1* and the subsequent homologous recombination between the two copies of *IS1* could result in the inversion of the intervening region (Panel B, Fig. 1). Second, it has been proven that a single copy of *ISEcp1* is able to mobilise its downstream genetic components by using alternative sequences as the right-hand inverted repeat (IRR)¹⁵. By comparing genetic contexts of *bla*_{CMY-2} and its variants available in GenBank using BLAST, the same 4 kb element comprising *ISEcp1*, *bla*_{CMY}, *blc* (encoding an outer membrane lipoprotein) and *sugE* (encoding a quaternary ammonium compound-resistance protein) has been found on different plasmids but with varied abutting sequences. This element is bounded by a 14bp sequence (AACCAGAAAGTCGA) at one end, which shows some similarity with the left-hand inverted repeat (IRL) of *ISEcp1* (Panel B, Fig. 1) and might have served as the alternative IRR (IRR2) for *ISEcp1* to mobilise *bla*_{CMY} to different locations. Indeed, when we put the two remnants of *ISEcp1* back together, we found that the *finQ* gene (the transcriptional inhibitor of plasmid transfer) has been interrupted by the *ISEcp1-bla*_{CMY-42}-IRR2 region with the presence of 5-bp direct target repeats (DR), which characterises the insertion of *ISEcp1* (Panel B, Fig. 1). This confirms that the mobilisation of *bla*_{CMY-42} has been mediated by *ISEcp1* by misidentifying its IRR.

In strain WCHEC13-8, *bla*_{NDM-1} was able to be transferred to *E. coli* strain J53, suggesting that pNDM1_EC8 was a self-transmissible plasmid. None of the other resistance genes listed above except *bla*_{SHV-12} were co-transferred with *bla*_{NDM-1}. PCR-based replicon typing (PBRT) failed to assign pNDM1_EC8 to a replicon type as all replicon typing PCR reactions were negative. pNDM1_EC8 was an IncX3 plasmid and IncX3 has not been integrated into the PBRT scheme. pNDM1_EC8 is identical to pNDM-HF727 (GenBank accession number KF976405), an IncX3 plasmid carrying *bla*_{NDM-1} from an *Enterobacter cloacae* strain in China⁹.

IncX3 plasmids are narrow-host-range plasmids of *Enterobacteriaceae*¹⁶. A few IncX3 plasmids carrying ESBL genes (*bla*_{SHV-11} or *bla*_{SHV-12}) and/or carbapenemase genes (*bla*_{KPC-2}, *bla*_{NDM-1} or *bla*_{NDM-5}) from various species of *Enterobacteriaceae* in several countries have been completely sequenced^{9,17-21}. In particular, IncX3 plasmids carrying *bla*_{NDM-1} and *bla*_{SHV-12} have been identified in various species of the *Enterobacteriaceae* at multiple locations in China. The completely-sequenced IncX3 plasmids except pKPI3d are almost identical in their backbone sequences with less than 9 single nucleotide variations between any two plasmids (Fig. 2). The remarkable similarity between IncX3 plasmids suggests that a common IncX3 plasmid has spread to different continents, acquired the carbapenemase gene that is prevalent locally, e.g. *bla*_{KPC-2} in USA and *bla*_{NDM-1} in China, and then were transformed as a vehicle mediating the interspecies spread of carbapenemase genes in local settings.

IncX3 plasmids carrying *bla*_{NDM-1} and *bla*_{SHV-12} have also been found in strains causing an outbreak in China²². IncX3 plasmids carrying *bla*_{NDM-1} and *bla*_{SHV-12} have also been found among the *Enterobacteriaceae* in the United Arab Emirates (UAE). Of note, one of the UAE strain was recovered two years earlier than the first Chinese strain that harboured an IncX3 plasmid carrying *bla*_{NDM-1} and *bla*_{SHV-12} was isolated. It remains unclear where the exact origin of emergence of IncX3 plasmids carrying *bla*_{NDM-1} and *bla*_{SHV-12} is. More studies are warranted to characterise the prevalence of IncX3 plasmids in the *Enterobacteriaceae* and to understand the emergence and spread of IncX3 plasmids²³.

The genetic context of *bla*_{NDM-1} on pNDM1_EC8 is identical to those on pNDM-HF727 and several other IncX3 plasmids from China. In such a context, *bla*_{NDM-1} is adjacent to an IS5-interrupted IS3000-truncated IS_{Aba125} at upstream and is linked with *bla*_{SHV-12} at downstream (Fig. 3). *bla*_{SHV-12} is flanked by IS26, which could form a composite transposon with the potential to mobilise *bla*_{SHV-12} into different locations.

Conclusions

An ST3835 *E. coli* clinical isolate carrying multiple genes, e.g. *bla*_{NDM-1}, *bla*_{CTX-M-15}, *bla*_{CMY-42} and *bla*_{SHV-12}, which are able to confer resistance to broad-spectrum cephalosporins, was identified in China. Carbapenemase gene *bla*_{NDM-1} was carried by a self-transmissible IncX3 plasmid in this strain. IncX3 plasmids may have served as a common vehicle mediating the dissemination of *bla*_{NDM} among the *Enterobacteriaceae*. The recently-identified plasmid-born AmpC gene *bla*_{CMY-42} was in a complex genetic




Figure 1. Genetic context of *bla*_{CMY-42} and its formation on pCMY42_EC8. Panel A, The context of *bla*_{CMY-42} and shapes with a forked tail represent truncated genes or insertion sequences. The broken line represents a 27.4 kb region between *traC* and *finQ*, which is not scaled. The putative alternative IRR (IRR2) of *ISEcp1* is depicted as a red pole and the alignment of the IRL of *ISEcp1* and the IRR2 is shown. *ISEcp1* and *traB* are both interrupted into two parts. The 9-bp nucleotide repeats belonging to *ISEcp1* are depicted in blue and those of *traB* are in red, while the 5-bp nucleotide repeats belonging to *finQ* is shown in bold. Panel B, The proposed scheme for the formation of the context of *bla*_{CMY-42}. The *ISEcp1*-*bla*_{CMY-42}-IRR2 region (structure 1) is inserted into the *finQ* gene of the *traA*-*trbA* region (structure 2) on an IncII plasmid and then generates the structure 3 with the 5-bp characteristic DR (GATAA). Two copies of *IS1* (structure 4) are then inserted into *traB* and *ISEcp1*, respectively, generating the structure 5 with the 9-bp DR (CCGTCAATA for *traB* and CGTTTGAAG for *ISEcp1*), which is the characteristic of *IS1* insertion. The subsequent homologous recombination between the two copies of *IS1* mediates the inversion (shown by two broken arrows) of the intervening region (grey part) and then forms the context of *bla*_{CMY-42} on pCMY42_EC8 (structure 6).




Figure 2. Phylogenetic trees of IncX3 plasmids. The trees are generated using MEGA based on the complete sequence of IncX3 plasmids. Panel A, a tree of IncX3 plasmids including pKPI3d, which carries *bla*_{KPC-2} in a *K. pneumoniae* strain from Brazil (GenBank accession number CP003997). pKPI3d is distinct from other IncX3 plasmids in sequence. Panel B, a higher resolution tree of IncX3 plasmids excluding pKPI3d. The carbapenemase and ESBL genes, the host species, the locations of recovery and the GenBank accession number of each plasmid are shown. For host species, CF refers to *Citrobacter freundii*, EC to *E. coli*, ECL to *E. cloacae*, KP to *K. pneumoniae* and RP to *Raoultella planticola*. The plasmid carrying *bla*_{NDM-1} identified in this study is identical to pNDM-HF727, which is underlined. The GenBank accession number of pNDM5_0215¹⁷ is not available.

context, which was formed by the transposition of *ISEcp1* and the insertion of two copies of *IS1* followed by homologous recombination between the two *IS1*.

Material and Methods

Strain and *in vitro* susceptibility. Strain WCHC13-8 was a clinical isolate recovered in West China Hospital of Sichuan University, Chengdu, China. Species identification and *in vitro* susceptibility test were performed using the Vitek II automated system (bioMerieux, Lyon, France). MICs of imipenem, meropenem, amikacin, ceftazidime and ciprofloxacin were also determined using the microdilution broth method followed recommendations of the Clinical Laboratory Standards Institute²⁴.

Detection of carbapenemase-encoding genes. The strain was screened for acquired carbapenemase-encoding genes *bla*_{GES}, *bla*_{KPC}, *bla*_{IMP}, *bla*_{IMI}, *bla*_{NDM}, *bla*_{OXA-48} and *bla*_{VIM} using PCR as described previously²⁵⁻²⁸. The complete coding sequence of *bla*_{NDM} was amplified with an additional pair of primers (NDM-up/NDM-dw)²⁵ and the amplicon was sequenced in both directions using an ABI 3730xl DNA Analyzer (Applied Biosystems, Foster City, CA, USA) at the Beijing Genomics Institute (Beijing, China).




Figure 3. Genetic context of bla_{NDM-1} in *E. coli* strain WCHEC13-8. ISAbal25 was interrupted by the insertion of IS5. Genes shown from the left in order are *mpr* (encoding a zinc metalloproteinase), bla_{NDM-1} , *ble* (mediating bleomycin resistance), *trpF* (encoding a phosphoribosylanthranilate isomerase), *dsbC* (encoding an oxidoreductase), *cutA1* (encoding an ion tolerant protein), *groL* (encoding a chaperonin subunit), *ygbI* (a putative dehydrogenase gene), a truncated *ygbJ* (a putative DEOR-type transcriptional regulator gene), bla_{SHV-12} and a truncated *umuD* (encoding a mutagenesis protein).

Genome sequencing and analysis. Genomic DNA of strain WCHEC13-8 was prepared using QIAamp DNA Mini Kit (Qiagen, Hilden, Germany) and was subjected to whole genome sequencing with a ca. $140\times$ coverage using the HiSeq 2500 Sequencer (Illumina, San Diego, CA, USA) following the manufacturer's protocol at the Beijing Genomics Institute. Reads were assembled to contigs using the Spades program²⁹. The Prokka program³⁰ was employed for annotating the genomic sequence. To identify whether this strain had unique genes, its assembled sequence was compared to the complete chromosome sequences of all of the 83 *E. coli* strains with their complete genome available in GenBank using the Gegenees program³¹. Potential unique genes were called using a stringent threshold of a 0.9 or higher score generated by Gegenees and then were confirmed or rejected using the BLAST program (<http://blast.ncbi.nlm.nih.gov>) against the 2,159 draft genomes of *E. coli* (accessed on December 18, 2014) to test whether they were absent from the *E. coli* draft genome database or not. Function of the products encoded by unique genes was predicted using both the Protein BLAST and InterProScan (<http://www.ebi.ac.uk/interpro/>) programs. The plasmids carrying bla_{NDM-1} , bla_{CMY-42} or $bla_{CTX-M-15}$ were completely circularised in sequence with intervals between contigs being filled by PCR with primers designed based on available sequences and Sanger sequencing.

Strain typing. Phylogenetic group (A, B1, B2 and D) typing was performed as described previously³². Strain WCHEC13-8 was assigned to a ST using the assembled genomic sequence to query the seven alleles of the multi-locus sequence typing scheme for *E. coli* (<http://mlst.warwick.ac.uk/mlst/dbs/Ecoli>)³³. Clonal complexes were assigned according to the founder ST identified using eBURST.

Mating experiments and plasmid typing. Conjugation experiment was carried out in broth using azide-resistant *E. coli* strain J53 as the recipient. Transconjugants were selected on plates containing 4 $\mu\text{g/ml}$ meropenem plus 150 $\mu\text{g/ml}$ sodium azide. The presence of bla_{NDM} in transconjugants was confirmed using PCR and ERIC-PCR was used for further distinguishing transconjugants from the donor strain. Plasmid DNA that was prepared from the transconjugant using alkaline lysis was subjected to PCR-based replicon typing (PBRT)³⁴. Sequences of IncX3 plasmids were retrieved from GenBank and aligned using MEGA (version 6). Maximum likelihood phylogenetic tree was constructed using MEGA using backbone sequences. IncI1 plasmid was assigned to ST and the replicon types of the IncF plasmids were assigned after querying their sequence against the plasmid MLST databases (<http://pubmlst.org/plasmid/>). Plasmids of the same type as pCTXM15_EC8 were identified using concatenated sequence of the IncF replicon alleles of pCTXM15_EC8 for BLAST.

S1-PFGE. S1-PFGE was performed to determine the number and size of plasmids carried by strain WCHEC13-8 as described previously³⁵. Briefly, agarose plugs containing whole-cell DNA of strain WCHEC13-8 were treated with 8 U of S1 nuclease (Fermentas, Thermo Scientific; Waltham, MA, US) and the reaction was stopped by adding 0.5 M EDTA (pH 8). PFGE was conducted with a 1% SeaKem Gold agarose gel (Lonza, Basal, Switzerland) using a CHEF DRII system (Bio-Rad, Hercules, CA, US) at 14°C, with a 6-V/cm current and run times of 12 h at switch time of 5 to 40 s followed by 8 h at switch time of 3 to 8 s. MidRange I PFG Marker (NEB, Ipswich, MA, US) was used for size estimation.

Nucleotide sequence accession number. Reads and the Whole Genome Shotgun project of WCHEC13-8 genomic sequence have been deposited into DDBJ/EMBL/GenBank under accession SRR2012643 and LCWG00000000, respectively. The sequence of pCMY42_EC8 and pCTXM15_EC8 has been deposited into DDBJ/EMBL/GenBank under accession numbers KP789019 and KP789020, respectively.

References

- Kumarasamy K. K. *et al.* Emergence of a new antibiotic resistance mechanism in India, Pakistan, and the UK: a molecular, biological, and epidemiological study. *Lancet Infect Dis* **10**, 597–602 (2010).
- Yong D. *et al.* Characterization of a new metallo- β -lactamase gene, bla_{NDM-1} , and a novel erythromycin esterase gene carried on a unique genetic structure in *Klebsiella pneumoniae* sequence type 14 from India. *Antimicrob Agents Chemother* **53**, 5046–5054 (2009).

3. Fu Y., Du X., Ji J., Chen Y., Jiang Y. & Yu Y. Epidemiological characteristics and genetic structure of *bla*_{NDM-1} in non-*baumannii* *Acinetobacter* spp. in China. *J Antimicrob Chemother* **67**, 2114–2122 (2012).
4. Zhou G. *et al.* NDM-1-producing strains, family *Enterobacteriaceae*, in hospital, Beijing, China. *Emerg Infect Dis* **20**, 340–342 (2014).
5. Hu F. *et al.* CHINET 2013 surveillance of bacterial resistance in China. *Chin J Infect Chemother* **14**, 369–378 (2014).
6. Zhang X. *et al.* First identification of coexistence of *bla*_{NDM-1} and *bla*_{CMY-42} among *Escherichia coli* ST167 clinical isolates. *BMC Microbiol* **13**, 282 (2013).
7. Liu Z. *et al.* Identification and characterization of the first *Escherichia coli* strain carrying NDM-1 gene in China. *PLoS One* **8**, e66666 (2013).
8. Qin S. *et al.* High incidence and endemic spread of NDM-1-positive *Enterobacteriaceae* in Henan Province, China. *Antimicrob Agents Chemother* **58**, 4275–4282 (2014).
9. Ho P., Li Z., Lo W., Cheung Y., Lin C. & Sham P. Identification and characterization of a novel incompatibility group X3 plasmid carrying *bla*_{NDM-1} in *Enterobacteriaceae* isolates with epidemiological links to multiple geographical areas in China. *Emerging Microbes and Infections* **1**, e39 (2012).
10. Hentschke M., Kotsakis S. D., Wolters M., Heisig P., Miriagou V. & Aepfelbacher M. CMY-42, a novel plasmid-mediated CMY-2 variant AmpC β -lactamase. *Microb Drug Resist* **17**, 165–169 (2011).
11. Pannaraj P. S., Bard J. D., Cerini C. & Weissman S. J. Pediatric Carbapenem-Resistant *Enterobacteriaceae* in Los Angeles, California, a High-Prevalence Region in the United States. *Pediatr Infect Dis J* **34**, 11–16 (2014).
12. Helmy M. M. & Wasfi R. Phenotypic and molecular characterization of plasmid mediated AmpC β -lactamases among *Escherichia coli*, *Klebsiella* spp., and *Proteus mirabilis* isolated from urinary tract infections in Egyptian hospitals. *Biomed Res Int* **2014**, 171548 (2014).
13. Villa L., Garcia-Fernandez A., Fortini D. & Carattoli A. Replicon sequence typing of IncF plasmids carrying virulence and resistance determinants. *J Antimicrob Chemother* **65**, 2518–2529 (2010).
14. Garcia-Fernandez A. *et al.* Multilocus sequence typing of IncI1 plasmids carrying extended-spectrum β -lactamases in *Escherichia coli* and *Salmonella* of human and animal origin. *J Antimicrob Chemother* **61**, 1229–1233 (2008).
15. Cattoir V., Nordmann P., Silva-Sanchez J., Espinal P. & Poirel L. ISEcp1-mediated transposition of *qnrB*-like gene in *Escherichia coli*. *Antimicrob Agents Chemother* **52**, 2929–2932 (2008).
16. Johnson T. J. *et al.* Expansion of the IncX plasmid family for improved identification and typing of novel plasmids in drug-resistant *Enterobacteriaceae*. *Plasmid* **68**, 43–50 (2012).
17. Yang P., Xie Y., Feng P., Zong Z. *bla*_{NDM-5} carried by an IncX3 plasmid in *Escherichia coli* of the sequence type 167. *Antimicrob Agents Chemother* **58**, 7548–7552 (2014).
18. Garcia-Fernandez A. *et al.* *Klebsiella pneumoniae* ST258 producing KPC-3 identified in Italy carries novel plasmids and OmpK36/OmpK35 porin variants. *Antimicrob Agents Chemother* **56**, 2143–2145 (2012).
19. Ramos P. I. *et al.* Comparative analysis of the complete genome of KPC-2-producing *Klebsiella pneumoniae* Kp13 reveals remarkable genome plasticity and a wide repertoire of virulence and resistance mechanisms. *BMC Genomics* **15**, 54 (2014).
20. Ho P. L. *et al.* Molecular characterization of an atypical IncX3 plasmid pKPC-NY79 carrying *bla*_{KPC-2} in a *Klebsiella pneumoniae*. *Curr Microbiol* **67**, 493–498 (2013).
21. Kassis-Chikhani N. *et al.* Complete nucleotide sequence of the first KPC-2- and SHV-12-encoding IncX plasmid, pKpS90, from *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* **57**, 618–620 (2013).
22. Wang X. *et al.* An outbreak of a nosocomial NDM-1-producing *Klebsiella pneumoniae* ST147 at a teaching hospital in mainland China. *Microb Drug Resist* **20**, 144–149 (2014).
23. Sonnevend A. *et al.* Emergence and spread of NDM-1 producer *Enterobacteriaceae* with contribution of IncX3 plasmids in the United Arab Emirates. *J Med Microbiol* **62**, 1044–1050 (2013).
24. CLSI. Performance Standards for Antimicrobial Susceptibility Testing; Twenty-Third Informational Supplement. M100-S23. (ed[^](eds). Clinical and Laboratory Standards Institute (2013).
25. Zong Z. & Zhang X. *bla*_{NDM-1}-carrying *Acinetobacter johnsonii* detected in hospital sewage. *J Antimicrob Chemother* **68**, 1007–1010 (2013).
26. Mendes R. E. *et al.* Rapid detection and identification of metallo- β -lactamase-encoding genes by multiplex real-time PCR assay and melt curve analysis. *J Clin Microbiol* **45**, 544–547 (2007).
27. Poirel L., Le Thomas I., Naas T., Karim A. & Nordmann P. Biochemical sequence analyses of GES-1, a novel class A extended-spectrum β -lactamase, and the class 1 integron In52 from *Klebsiella pneumoniae*. *Antimicrob Agents Chemother* **44**, 622–632 (2000).
28. Bradford P. A. *et al.* Emergence of carbapenem-resistant *Klebsiella* species possessing the class A carbapenem-hydrolyzing KPC-2 and inhibitor-resistant TEM-30 β -lactamases in New York City. *Clin Infect Dis* **39**, 55–60 (2004).
29. Bankevich A. *et al.* SPAdes: a new genome assembly algorithm and its applications to single-cell sequencing. *J Comput Biol* **19**, 455–477 (2012).
30. Seemann T. Prokka: rapid prokaryotic genome annotation. *Bioinformatics* **30**, 2068–2069 (2014).
31. Agren J., Sundstrom A., Hafstrom T. & Segerman B. Gegenees: fragmented alignment of multiple genomes for determining phylogenomic distances and genetic signatures unique for specified target groups. *PLoS One* **7**, e39107 (2012).
32. Clermont O., Bonacorsi S. & Bingen E. Rapid and simple determination of the *Escherichia coli* phylogenetic group. *Appl Environ Microbiol* **66**, 4555–4558 (2000).
33. Wirth T. *et al.* Sex and virulence in *Escherichia coli*: an evolutionary perspective. *Mol Microbiol* **60**, 1136–1151 (2006).
34. Carattoli A., Bertini A., Villa L., Falbo V., Hopkins K. L. & Threlfall E. J. Identification of plasmids by PCR-based replicon typing. *J Microbiol Methods* **63**, 219–228 (2005).
35. Partridge S. R., Zong Z. & Iredell J. R. Recombination in IS26 and Tn2 in the evolution of multiresistance regions carrying *bla*_{CTX-M-15} on conjugative IncF plasmids from *Escherichia coli*. *Antimicrob Agents Chemother* **55**, 4971–4978 (2011).

Acknowledgements

This work was supported by a grant from the National Natural Science Foundation of China (project no. 81222025), the New Century Excellent Talents Program, Ministry of Education, China (project no. NCET-13-0399) and a grant from Sichuan Bureau of Science, China (project no. 2013JQ0042) to ZZ and Royal Society/NSFC international collaboration award IE121459 (Royal Society)/81311130143 (NSFC) to AM and ZZ.

Author Contributions

Concept and design: Z.Z.; Acquisition of data: Y.F., P.Y., X.W. and X.Y.; Analysis/Interpretation of data: A.M. and Z.Z.; Draft of manuscript: Z.Z.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Feng, Y. *et al.* *Escherichia coli* of sequence type 3835 carrying *bla*_{NDM-1}, *bla*_{CTX-M-15}, *bla*_{CMY-42} and *bla*_{SHV-12}. *Sci. Rep.* **5**, 12275; doi: 10.1038/srep12275 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>