

Diabetes-induced microvascular complications at the level of the spinal cord; a contributing factor in diabetic neuropathic pain

^{**a,c}N. Ved, ^{**a}M.E. Da Vitoria Lobo, ^bS.M. Bestall, ^aC.L. Vidueira, ^bN. Beazley-Long, ^dK. Ballmer-Hofer, ^e M. Hirashima, ^{a,g}D.O. Bates, ^bL.F Donaldson and ^{a,f}R.P. Hulse*

^aCancer Biology, Division of Cancer and Stem Cells, School of Medicine University of Nottingham, Nottingham NG7 2UH, ^bArthritis Research UK Pain Centre and School of Life Sciences, The Medical School QMC, University of Nottingham, Nottingham NG7 2UH. ^cInstitute of Ophthalmology, 11-43 Bath St, London EC1V 9EL, ^dPaul Scherer Institute, Villingen, Switzerland, ^e Division of Vascular Biology, Kobe University. ^f School of Science and Technology, Nottingham Trent University, Nottingham, NG11 8NS. ^g **Centre of Membrane and Protein and Receptors (COMPARE), University of Birmingham and University of Nottingham, Midlands, UK.**

*Corresponding author

**These authors contributed equally to this work

Corresponding Author Contact Details

Dr RP Hulse

Richard.Hulse@nottingham.ac.uk

School of Science and Technology

NHB 165

Nottingham Trent University

Clifton Lane

Nottingham

NG11 8NS

Richard.Hulse@ntu.ac.uk

Tel: 01158486368

This is an Accepted Article that has been peer-reviewed and approved for publication in the The Journal of Physiology, but has yet to undergo copy-editing and proof correction. Please cite this article as an 'Accepted Article'; [doi: 10.1113/JP275067](https://doi.org/10.1113/JP275067).

This article is protected by copyright. All rights reserved.

Category: Original Article

Running title: Diabetes-induced microvascular degeneration and neuropathic pain

Keywords: pain, diabetes, endothelial

Key Point Summary

1. Diabetes is thought to induce neuropathic pain through activation of dorsal horn sensory neurons in the spinal cord.
2. Here we explore the impact of hyperglycaemia on the blood supply supporting the spinal cord and chronic pain development.
3. In streptozotocin-induced diabetic rats, neuropathic pain is accompanied by a decline in microvascular integrity in the dorsal horn. Hyperglycaemia-induced degeneration of the endothelium in the dorsal horn was associated with a loss in VEGF-A_{165b} expression. VEGF-A_{165b} treatment prevented diabetic neuropathic pain and degeneration of the endothelium in the spinal cord.
4. Using an endothelial specific VEGFR2 knockout transgenic mouse model, the loss of endothelial VEGFR2 signalling led to a decline in vascular integrity in the dorsal horn and the development of hyperalgesia in VEGFR2 knockout mice.
5. This highlights that vascular degeneration in the spinal cord could be a previously unidentified factor in the development of diabetic neuropathic pain.

Abstract

Abnormalities of neurovascular interactions within the central nervous system of diabetic patients is associated with the onset of many neurological disease states. However, to date, the link between the neurovascular network within the spinal cord and regulation of nociception has not been investigated despite neuropathic pain being common in diabetes. We hypothesised that hyperglycaemia-induced endothelial degeneration in the spinal cord, due to suppression of VEGF-A/VEGFR2 signalling, induces diabetic neuropathic pain. Nociceptive pain behaviour was investigated in a chemically induced model of type 1 diabetes (streptozotocin induced, insulin supplemented; either vehicle or VEGF-A_{165b}

treated) and an inducible endothelial knockdown of VEGFR2 (tamoxifen induced). Diabetic animals developed mechanical allodynia and heat hyperalgesia. This was associated with a reduction in the number of blood vessels and reduction in Evans blue extravasation in the lumbar spinal cord of diabetic animals versus age-matched controls. Endothelial markers occludin, CD31 and VE-cadherin were downregulated in the spinal cord of the diabetic group versus controls, as well as a concurrent reduction of VEGF-A₁₆₅b expression. In diabetic animals, VEGF-A₁₆₅b treatment (biweekly intraperitoneal, 20ng/g) restored normal Evans blue extravasation and prevented vascular degeneration, diabetes-induced central neuron activation and neuropathic pain. Inducible knockdown of VEGFR2 (tamoxifen treated *Tie2CreER^{T2}-vegfr2^{fl/fl}* mice) led to a reduction in blood vessel network volume in the lumbar spinal cord and development of heat hyperalgesia. These findings indicate that hyperglycaemia leads to a reduction in the VEGF-A/VEGFR2 signalling cascade resulting in endothelial dysfunction in the spinal cord, which could be an undiscovered contributing factor to diabetic neuropathic pain.

Abbreviations

VEGF-A = vascular endothelial growth factor-A

VEGFR2 = vascular endothelial growth factor receptor 2

STZ = streptozotocin

i.p. = intra-peritoneal

Introduction

Diabetes mellitus leads to an array of health complications that can cause significant morbidity. In people with diabetes, neuropathic pain is common (Tesfaye *et al.*, 2013), characterised by enhanced responses to noxious (painful) stimuli (hyperalgesia) as well as to innocuous stimuli (allodynia). These alterations in pain perception are due to maladaptive changes in the sensory neuronal circuitry. The plasticity of the nociceptive neuronal systems, both peripheral (Reichling & Levine, 2009) and central (Latremoliere A, 2009), means that they can respond to disease and/or treatment such as in diabetes (Chen & Levine, 2001) (Morgado *et al.*, 2010; Tan *et al.*, 2012). These responses lead to neuronal sensitisation, and in diabetes, chronic pain development. The peripheral sensory nerves are well known to be affected by hyperglycaemia, including degeneration of intra-epidermal nerve fibre innervation patterns (Hulse *et al.*, 2015) and hyper-excitability (Chen & Levine, 2003). However, pain

management in people with diabetes often only provides partial pain relief (Tesfaye *et al.*, 2011; Tesfaye *et al.*, 2013). There are now focussed efforts to investigate how changes in nociceptive processing in the central nervous system, in particular the spinal cord, are altered in diabetic neuropathic pain (Biessels *et al.*, 2014; Tesfaye *et al.*, 2016). Studies have identified that in diabetic rodents, sensory neurons within the spinal cord elicit exaggerated responses to sensory stimulation (Morgado *et al.*, 2010; Tan *et al.*, 2012). Despite this evidence for the involvement of spinal cord changes in the pathogenesis of diabetic neuropathic pain, there are few investigations into those mechanisms that may underlie the development of central sensitisation in the spinal cord in diabetes (Tan *et al.*, 2012; Lee-Kubli CA, 2013).

An important component of the nervous system is the supporting blood vessel network. A compromised vascular system is integral to the development of multiple neurological diseases (e.g. stroke, Alzheimer's disease) (Tiehuis *et al.*, 2008; Vandal *et al.*, 2014; Winkler *et al.*, 2015). Hyperglycaemia induces extensive vascular remodelling in the nervous system (Taylor *et al.*, 2015; Hardigan *et al.*, 2016) as well as direct glucose toxicity on sensory neurons (Chowdhury *et al.*, 2014; Hulse *et al.*, 2015), with both of these contributing to neurological complications including increased susceptibility of people with diabetes to cognitive decline, stroke, and peripheral ischaemic neuropathies (motor, sensory and autonomic) (Said, 2007; Hardigan *et al.*, 2016). Vascular endothelial growth factor-A (VEGF-A) is strongly implicated in diabetic vascular disease, including driving aberrant vessel growth and increased permeability in diabetic retinopathy (Cai & Boulton, 2002). This is highly correlated with increased VEGF-A expression and is therefore a prime target for diabetic retinopathy treatment (Gupta *et al.*, 2013). The VEGF-A gene gives rise to a variety of VEGF-A splice variants, differing in length and terminal sequence, leading to contrasting functions (Harper & Bates, 2008). The archetypal proangiogenic isoform is VEGF-A_{165a} and is typically associated with vascular remodelling such as in diabetic retinopathy and cancer (Perrin *et al.*, 2005). The VEGF-A_{165b} isoform is predominantly found in normal tissues, areas with reduced angiogenesis (Pritchard-Jones *et al.*, 2007), and pathologies where angiogenesis is impaired (e.g. systemic sclerosis and peripheral arterial disease) (Manetti *et al.*, 2011; Kikuchi *et al.*, 2014; Ngo *et al.*, 2014). It is known to compete with VEGF-A_{165a} for VEGFR2 binding (Bates *et al.*, 2002; Cébe Suarez S, 2006) and can independently act through VEGFR2, resulting in cytoprotective effects (Beazley-Long *et al.*, 2013). For example, in diabetic nephropathy VEGF-A_{165b} treatment leads to a rescue in endothelial cell

survival and return to normal kidney function (Oltean *et al.*, 2014), and we have previously found that indicators of altered blood vascular integrity, enhanced in the peripheral nervous system and the dorsal root ganglia, can be reversed by VEGF-A_{165b}, resulting in amelioration of pain behaviours in rats (Hulse *et al.*, 2015).

Enhanced sensitivity in central nociceptive networks (spinal cord (Tan *et al.*, 2012; Lee-Kubli CA, 2013), brain (Silva M, 2013)) have long been attributed to neuroplastic changes. Despite extensive evidence that the cerebral vasculature is altered in diabetic rodents, there has been limited investigation into the neurovascular interactions in the spinal cord particularly with reference to nociception (Costigan *et al.*, 2009; Beggs *et al.*, 2010). Here we hypothesised that a decline (endothelial cell loss) in the vascular system within the spinal cord could contribute to the onset of diabetic neuropathic pain. Using an *in vivo* rat model of type 1 diabetes, and an inducible VEGF receptor-2 (VEGFR2) knockdown transgenic mouse, neuro-vascular disruption in the spinal cord was associated with changes in neuropathic pain. Administration of the VEGF-A_{165b} isoform protected the endothelial component of the CNS, and prevented diabetic neuropathic pain.

Methods

Ethical Approval, Animals used and Diabetes induction

24 male (21 Evans blue and 3 S1 tissue collection) and 42 female (24 Evans blue and 18 S1 tissue collection) Sprague Dawley rats (~250g) were used in this study. Experiments were carried out in accord with the institution's animal welfare committee (UoN), and conform to the principles and regulations as described in the Editorial by Grundy (Grundy, 2015). Procedures were carried out in accordance with the UK Home Office Animals (Scientific Procedures) Act 1986 and EU Directive 2010/63/EU after review by the local Animal Welfare and Ethics Review Board (UoN). Diabetes was induced by intraperitoneal (i.p.) injection of streptozotocin (STZ) (50mg/kg)(Hulse *et al.*, 2015). Animals had *ad libitum* access to standard chow and were housed in groups (n>2) under 12:12h light:dark conditions. In male rats, experimental groups (1 week, no insulin supplementation) were naïve (sham injected n=9) and STZ treated (n=12). At the end of the study animal weight and blood glucose (>15mmol) were naïve=339±8.26g and 6.8±0.85 mmol; diabetics=314±11.86g and 30.12±0.95 mmol respectively.

In female rats (8 week experiments), animals were treated with insulin using one third of an insulin pellet (Linshin, Canada) implanted under isoflurane anaesthesia (2-3% in O₂) (Calcutt, 2004). Experimental treatments were biweekly recombinant human (rh)VEGF-A_{165b} (20ng/g body weight, i.p. twice weekly from week 1) or saline (vehicle; i.p.). This VEGF-A_{165b} regime has previously been used (Beazley-Long *et al.*, 2013; Hulse *et al.*, 2014)). Blood glucose and weight was measured in all animals at the end of the study. Blood glucose: naïve=8.08±0.72, diabetic+vehicle=29.27±1.35, diabetic+VEGF-A_{165b}=30.81±0.78. Animal body weight: naïve=324.1g±8.5, diabetic+vehicle=284.2±7.3g, diabetic+VEGF-A_{165b}=289.1±5.4g.

75 transgenic mice were used in this study (C57.bl6, 25-30g; both genders). Tie2CreER^{T2} mice [Tg(Tek-cre/ER^{T2})1Arnd, European Mutant Mouse Archive] were crossed with *vegfr2*^{fl/fl} (generated/used as previously described (Albuquerque *et al.*, 2009; Sison *et al.*, 2010; Beazley-Long *et al.*, 2018)). All mice used were *vegfr2*^{fl/fl} and either Tie2CreER^{T2} positive (n=27) or Tie2CreER^{T2} negative (n=28) and dosed once daily by i.p. with 1mg tamoxifen or vehicle (10% ethanol in sunflower oil) for 5 consecutive days.

Nociceptive Behaviour

Nociceptive behavioural experiments were carried out as previously described (Drake *et al.*, 2014; Hulse *et al.*, 2016). on 8 week diabetic animals and age matched sham controls (naïve). Mechanical withdrawal thresholds were measured using von Frey (vF) monofilaments (Hulse *et al.*, 2015) or a mechanical pincher (Drake *et al.*, 2014; Hirschberg *et al.*, 2017). A range of von Frey hairs were applied to the hind paw plantar surface (a maximum of five seconds or until paw withdrawal). A total of five vF applications were applied per weighted hair and force response curves were generated and withdrawal values were calculated as the weight at which withdrawal frequency = 50%. A mechanical pincher (equipped with strain gauges and calibrated to force (g)) (Drake *et al.*, 2014) was applied to the hind paw until the animal withdrew to determine mechanical hyperalgesia. Raw data were acquired through a Neurolog power unit and a bridge amp module (Digitimer), with digital acquisition via CED micro1401v3 and Spike2 v7 software (Cambridge Electronic Design UK). Withdrawal to heat was determined using the Hargreaves test (Hargreaves *et al.*, 1988). The experimenter was blinded to treatment.

Evans blue extravasation

In vivo vascular perfusion was evaluated using Evans blue dye, as previously described (Xu *et al.*, 2001). Animals (age matched sham controls (naïve), week 1 and week 8 diabetic rats) were terminally anesthetized (ketamine medetomidine i.v. 50 mg/kg) and infused (via the external jugular vein) with Evans blue dye i.v. (Sigma-Aldrich, 45 mg/kg) at 120mmHg pressure. Two minutes post-infusion, 0.2mL arterial blood was withdrawn, followed by subsequent 0.1mL withdrawals every 15 minutes for 2 hours. After 2 hours 0.2mL blood was withdrawn followed by cardiac perfusion of 50mL saline at 120mmHg. Lumbar spinal cord and a single brain hemisphere were excised (whole spinal cord including ventral and dorsal horn) and weighed (wet weight). Tissue was dried at 70°C overnight and weighed (dry weight). Dried tissue was incubated in 0.15mL formamide (Sigma Aldrich) at 70°C overnight. Blood samples were centrifuged (12,000rpm, 45 minutes, 4°C), and the supernatant from tissue and blood samples were analysed at 620nm. Evans blue extravasation was calculated as (solute flux (µg/min/g) = Evans blue mass (µg)/ tissue dry weight (g) divided by time (120min).

Immunofluorescence analyses

Animals from all experimental groups (normal, diabetes, diabetes+VEGF-A_{165b}; 8 weeks) were terminally anaesthetised (sodium pentobarbital 60mg/kg i.p.) and transcardially perfused, with phosphate buffered saline (PBS) followed by 4% paraformaldehyde in PBS (PFA; pH7.4). Tissue was prepared as previously described (Hulse *et al.*, 2016). Spinal cords (40µm thickness) were incubated in primary antibodies (see below) in blocking solution (5% bovine serum albumin, 10% fetal calf serum), overnight at 4°C (IB₄ 72 hours). Primary antibodies/markers and dilutions used were: biotin conjugated isolectin B₄ (IB₄;1 in 100, Sigma-Aldrich); rat anti-CD31 (MEC13.3;1 in 10, Santa Cruz), mouse anti-CD31 (1 in 100, Abcam), rabbit anti-GFAP (1 in 500, Abcam), rabbit anti-fos (1 in 100, Santa Cruz,); mouse anti-NeuN (1 in 200; Millipore), rabbit anti-VEGFR2 (1 in 200, 55B11, Cell Signalling), rabbit anti-cleaved caspase 3 (1 in 500, Cell Signalling) and anti-rabbit biotinylated IgG (1 in 500, Jackson Laboratories). Secondary antibodies were incubated in PBS + 0.2% Triton X-100, which were Alexa Fluor 488-conjugated chicken anti-mouse, Alexa Fluor 555-conjugated donkey anti-rabbit and streptavidin-conjugated Alexa Fluor-555 (1 in 500, all Invitrogen, UK. Confocal imaging of the dorsal horn of the lumbar spinal cord of all groups was performed on a Leica TCS SPE confocal microscope.

Spinal cord Endothelial Cell Culture

Whole lumbar spinal cords were dissected from adult male Sprague Dawley rats and dissociated (0.125% collagenase). Endothelial cells were extracted and cultured in endothelial media (M199 media, 60µg/ml endothelial cell growth supplement and 50ug/ml Heparin). Endothelial cells were plated (1% gelatin coated) onto either 6 well plates (for protein extraction) or 96 well plates (for cell viability studies). For cell viability assays, when 80% confluent endothelial cells were incubated for 24 hours in either 5mM glucose (normal), 50mM mannitol (osmotic control) or 50mM glucose (high glucose) \pm 2.5nM VEGF-A_{165b} or vehicle. Cell death was determined with neutral red (Sigma-Aldrich).

Flow Cytometry

Spleens were isolated from CTL and eCKO transgenic mouse experimental groups and placed in media (containing RPMI1640, Penicillin/streptomycin, 10% FBS, 1%l-glutamine, 0.1% sodium pyruvate). Tissue was mechanically dissociated through a 40µm cell strainer and washed through with media. Cell suspension was then centrifuged at 1500rpm for 3mins. Supernatant was removed and cells were resuspended in 2 ml red cell lysis medium (Sigma)., then left for 30 seconds. Following this 10ml of media was added and the cells were centrifuged as previously. Cells were then fixed in 4% paraformaldehyde for 15minutes at room temperature and then subsequently washed three times. Cells were permeabilised with 0.4% Triton X-100 in PBS for 15 minute at room temperature. Cells were blocked in 1% FBS in PBS, mixed gently and incubated for 30 min at room temperature. Antibodies (CD11b-APC (1 in 100; Biolegend) and F4/80-PE (1 in 100; Biolegend)) were added, left overnight in the fridge, and samples analysed on MoFlo analyser.

qPCR Method

Total RNA was extracted from whole lumbar spinal cord tissue isolated from CTL and eCKO transgenic colony experimental groups using TRIzol reagent (Invitrogen). cDNA synthesis was carried using PrimeScriptTMRT reagent kit (TaKaRa,RR037A) with a starting amount of 1µg of RNA. The resulting cDNA was used for Q-PCR using a LightCycler 480 SYBR Green I Mastermix (Roche 04707516001) following the manufacturer's instructions. CD31 and VE-cadherin primers were synthesised by Eurofins. Beta actin and VEGFR2 primers were

synthesised by Sigma-Aldrich. Beta actin was used as a reference gene. The primer sequences were as follows (5'-3');

β -actin F – 5' ATTGCCAATGAGCGGTTC-3'

β -actin R – 5' GGATGCCACAGGACTCCA-3'

CD31 F - 5' GAAATGCTCTCGAAGCCCAG-3'

CD31 R - 5' ACCTCGAGAGTCTGGAAGTC-3'

VE- cadherin F - 5' TCCCTGGACTATGAAGTCAT-3'

VE- cadherin R - 5' GAAGACAGGGGGCTCATCCA-3'

VEGFR2 F - 5' GGATCTGAAAAGACGCTTGG-3'

VEGFR2 R – 5' TGCTCCAAGGTCAGGAAGTC-3'

Western Blotting

Protein was extracted from spinal cord endothelial cells and human umbilical vein endothelial cells as well as spinal cord tissue as previously described (Vencappa *et al.*, 2015; Hulse *et al.*, 2016). Animals from all experimental groups (CTL and eCKO transgenic mice; normal, diabetes, diabetes+VEGF-A_{165b}; 8 weeks) were terminally anaesthetized with sodium pentobarbital (i.p. 60 mg/kg, Sigma-Aldrich). Lumbar spinal cords were extracted, frozen immediately and stored at -80°C until sample processing. 100µg endothelial cell lysate and 80µg of each spinal cord lysates were loaded in a 4%-20% precast Mini-Protean TGX gel (Biorad), separated by SDS-PAGE electrophoresis and transferred using Trans-blot turbo transfer system (Bio-Rad). The membrane was incubated in 5% milk powder in tris-buffered saline (TBS)-Tween 0.1% (TBST) for 30 minutes at room temperature. Primary antibodies, mouse anti-CD31 (2µg/ml, Abcam; AB24590), rabbit anti-occludin (5µg/ml, Invitrogen; 71-1500), mouse anti-VE cadherin (5µg/ml, BD Biosciences; 550548), rabbit anti-VEGFR2 (1 in 200, 55B11, Cell Signalling), rabbit anti-Pan VEGF-A (A20, 1µg/ml, Santa Cruz; sc-152), mouse anti-VEGF-A_{165b} (2µg/ml, Abcam; ab-14994), and rabbit anti-Actin (1:100 Santa Cruz) antibodies were diluted in blocking solution and incubated overnight at 4°C. Secondary antibodies (Licor donkey anti-rabbit and anti-mouse antibodies 1:10000) in TBST-0.1% 1% BSA and visualised on the Licor Odessey.

Statistical Analysis

All data are represented as mean±SEM unless stated and experimenter was blinded where appropriate. Data were acquired/quantified using Microsoft Excel 2010, Graphpad Prism 6

and Imaris (Bitplane) Spike2 v7 software (CED) was used to digitally acquire mechanical withdrawal thresholds from the pincher and for offline analysis. Immunofluorescence was quantified by obtaining 10 random non-sequential sections (Z stacks) per animal and a mean value calculated per animal. Dorsal horn spinal cords (Lamina I-V) were imaged with the confocal as described above, and vessels were identified through CD31 and IB₄ immunoreactivity. IB₄ signal fluorescence was vascular in the majority of the spinal cord with C-fibre projections also staining in the peripheral laminae (I-II), but vessels could also be clearly delineated here by CD31 staining as well as morphology. Stained images were rendered on Imaris 8.11 imaging software. This allows for automated quantification of vessel diameter and blood vessel volume. Neuron number (cleaved caspase-3 and C-Fos quantification) was determined as per laminae of the dorsal horn and therefore expressed according to lamina I-V as previously characterised (Hsieh *et al.*, 2015). Western blot densitometry was quantified using ImageJ (<https://imagej.nih.gov/ij/>) gel quantification plugin. Paw mechanical withdrawal thresholds, the number of blood vessels in the dorsal horn of the spinal cord and spinal cord Evans blue extravasation were analysed using a Mann-Whitney test. Evans blue, IB₄ positive blood vessel volume and length, mechanical and heat nociceptive behaviour and western blot densitometry quantification were analysed using a Kruskal Wallis and appropriate post-hoc tests.

Results

Diabetes resulted in increased mechanical hyperalgesia - a reduction in withdrawal threshold to a noxious stimulus (hindpaw pinch) when compared to both before diabetes ($p < 0.05$) and to vehicle/age matched animals at week 8 post STZ injection (naïve; Fig. 1A, $*p < 0.05$). This was accompanied by a reduction in the number of blood vessels in the dorsal horn of the lumbar region of the spinal cord (Fig. 1B-D, $*p < 0.05$) as well as a reduction in microvessel diameter (labelled with CD31 and IB₄) (Fig. 1E-G);. The VEGF-A family is a key regulator of angiogenic processes. Pan-VEGF-A expression was unaltered in the lumbar spinal cord of diabetic rats when compared to age/gender matched control animals (Fig. 1H&I). However, VEGF-A_{xxx}b expression was significantly reduced in the lumbar spinal cord of diabetic animals (Fig. 1H, I, $*p < 0.05$).

As hyperglycaemia affected spinal cord microvasculature and reduced VEGF-A₁₆₅b expression, we investigated the direct cytoprotective actions of VEGF-A₁₆₅b upon cultured spinal cord endothelial cells. Isolated spinal cord endothelial cells showed increased cell

death when cultured in high (50mM) versus low glucose (5mM) conditions (Fig. 2A). There was no effect of 50mM mannitol, an osmotic control. VEGF-A₁₆₅b treatment prevented high glucose-induced endothelial cell death (Fig. 2A). Consequently, the cytoprotective actions of VEGF-A₁₆₅b on the spinal cord endothelium were investigated *in vivo*. The vascular network was well defined in naïve animals (Fig. 2B), was reduced in number in diabetic+vehicle treated animals (Fig. 2C) but this was prevented by VEGF-A₁₆₅b (Fig. 2D). Quantification indicated a reduction in endothelium (reduced total vascular volume) within the dorsal horn of the lumbar region of the spinal cord, which was rescued by VEGF-A₁₆₅b (Fig. 2E). There was also an overall reduction in dorsal horn vessel diameter, again prevented by VEGF-A₁₆₅b (Fig. 2F). There was a non-significant increase in vessel diameter in the diabetic+VEGF-A₁₆₅b treated group when compared with the naïve group (Fig. 2F). Analysis of the frequency distribution of the vessels by size indicated that the reduction in size in the diabetics was due to a reduced number of the “larger microvessels” (8-12µm) rather than a reduction in the number of small (<8µm vessels). (Fig. 2G). To determine whether markers of endothelial integrity/activation (including junctional markers) were altered in the lumbar spinal cord in diabetes, lumbar spinal cord protein samples were subjected to immunoblotting for VE-cadherin (Fig. 3A upper band), CD31 (Fig. 3A lower band), and occludin (Fig. 3B upper band; lower band actin). There was a marked reduction in junctional and adhesion molecules in diabetic+vehicle treated rats, with non-significant reductions in VE-cadherin (Fig. 3C) and significant reductions in CD31 (Fig. 3D) and occludin (Fig. 3E), all of which were prevented by VEGF-A₁₆₅b treatment.

To investigate the possible functional changes in the microvasculature within the spinal cord in diabetes, Evans blue extravasation was measured (Xu *et al.*, 2001). Evans blue vascular leakage is dependent on blood flow, surface area, hydrostatic pressure and vascular permeability. In an acute diabetic rodent model (1 week) there was a pronounced decrease in Evans blue extravasation within the lumbar region of the spinal cord compared with that of the control cohort (Fig. 4A). Diabetic groups were systemically (i.p.) treated with either saline or VEGF-A₁₆₅b (20ng/g) using a longer term diabetic model (8 weeks). In longer duration diabetes (8 weeks) there was also a significant reduction in Evans blue extravasation compared with the control naïve group (Fig. 4B), which was prevented by VEGF-A₁₆₅b (Fig. 4B). Brains were also extracted from the long term (8 week) study (Fig. 4C) though there was no difference in solute flux between groups.

Mechanical hypersensitivity and heat hyperalgesia developed only in the diabetic+vehicle animals. VEGF-A_{165b} treatment (timepoint of administration shown via arrow) not only prevented the spinal cord vascular degeneration (consistent with previous experiments (Hulse *et al.*, 2015; Ved *et al.*, 2017)) but also prevented diabetic neuropathic pain behaviours (mechanical allodynia Fig. 5A and heat hyperalgesia Fig. 5B). Within the spinal cord, not only was the vasculature disturbed in the diabetic+vehicle group, but sensory neurons (NeuN) in the dorsal horn also expressed increased cleaved caspase-3 (CC3), an indicator of neuronal damage, when compared with naïve age matched animals (Fig. 5C). There was increased CC3 immunoreactivity in the superficial lamina (I & II) of the dorsal horn of diabetic animals (Fig. 5D). This was blocked by VEGF-A_{165b} treatment (Fig. 5E&F-D). Sensory neurons within the spinal cord, once activated, express the immediate early gene *c-fos* (Hunt SP, 1987). *C-fos* is a marker of neuronal activation in chronic pain states (Kalynovska *et al.*, 2017; Khasabov *et al.*, 2017) and is an indicator of spinal neuronal activation in diabetic neuropathic pain (Morgado *et al.*, 2010). There was an increase in *c-fos* expression in neurons (NeuN co-labelled) within the lumbar dorsal horn in the spinal cord in diabetic rats (Fig. 6A&B) compared with naïve animals. Systemic treatment with VEGF-A_{165b} led to an attenuation of the diabetes induced *c-fos* expression (Fig. 6A&B). All lamina of the dorsal horn demonstrated increased *c-fos* expression in neurons in the diabetic+vehicle group compared with the naïve and diabetic+VEGF-A_{165b} groups (Fig. 6C).

The involvement of VEGF-A/VEGFR2 signalling on spinal cord endothelial cell function and survival and the relationship to the generation of behavioural hypersensitivity was determined by using an endothelial cell specific inducible VEGFR2 knockdown *in vivo*. Systemic tamoxifen treatment led to a reduction in VEGFR2 protein expression in endothelial cells from *vegfr2^{fl/fl} Tie2CreER^{T2}*-positive mice, i.e. VEGFR2 endothelial cell knockout (VEGFR2^{ECKO}) compared with control mouse endothelial cells (CTL = *vegfr2^{fl/fl} Tie2CreER^{T2}*-negative + tamoxifen) isolated from lung (Fig. 7A-B), as well as a reduction in VE-cadherin expression (Fig. 7C-D). In the spinal cord there was a reduction in VEGFR2 (Fig. 7E) as well as VE-cadherin (Fig. 7F) and CD31 (Fig. 7G) expression in the VEGFR2^{ECKO} animals compared with CTL animals. In addition, VE-cadherin protein expression from the spinal cord of VEGFR2^{ECKO} animals was reduced when compared with the CTL animals (Fig. 7H, I).

When compared with CTL mice (Fig. 8A-C), the VEGFR2^{ECKO} (Fig. 8A; low power, 8B high power) had a significant decline in vascular integrity within the lumbar spinal cord 8 days after the final drug injection. The endothelium demonstrated a reduced endothelial volume (Fig. 8C) as well as showing a reduced diameter of spinal cord microvessels in the VEGFR2^{ECKO} mice treated with tamoxifen when compared with CTL mice (and VEGFR2^{fl/fl} with Cre but not given tamoxifen “vehicle”) (Fig. 8D). There was an increase number of smaller vessels in the VEGFR2^{ECKO} mice when compared to control mice (CTL and vehicle) (Fig. 8E).

In VEGFR2^{ECKO} and CTL mice there was no difference in nociceptive behaviour prior to tamoxifen administration (Fig. 8F). The *vegfr2*^{fl/fl} Tie2CreER^{T2} mice treated with vehicle (“vehicle”) and CTL mice treated with either vehicle or tamoxifen demonstrated no change in nociceptive behavioural responses to heat (Fig. 8F). However, the VEGFR2^{ECKO} mice demonstrated a pronounced heat hypersensitivity compared with all other groups following tamoxifen injection (Fig. 6I). To exclude the involvement of a subset of haemopoietic cells that express Tie2 and VEGFR2 we determined the impact of VEGFR2 knockdown in this cell population. There was no change in the number of F4/80 (macrophage marker) or CD11b marker of leukocytes including monocytes, neutrophils, natural killer cells, granulocytes and macrophages) positive cell types isolated from the spleen in either the CTL mice treated with tamoxifen or VEGFR2^{ECKO} mice (Fig. 9).

Discussion

We show here that the microvasculature in the spinal cord was disrupted in a rodent model of diabetic neuropathic pain, demonstrated by a reduction in the volume of the blood vessel network in the spinal cord. These findings are associated with neuropathic pain development and spinal neuronal activation. There was a concurrent reduction in the spinal cord expression of the VEGF-A_{165b} isoform, but no overall change in total VEGF-A expression. In the inducible VEGFR2^{ECKO} mice, the vasculature in the lumbar spinal cord was also reduced and was accompanied by the development of hyperalgesia. Critically, reversing the diabetes-induced vascular degeneration using systemic VEGF-A_{165b} treatment also resulted in reversal of chronic pain.

Neuropathic pain is common in diabetic patients and is typically associated with hyperglycaemia impacting upon the peripheral vasculature through alterations in blood flow and solute leakage (Poduslo & Curran, 1992; Tesfaye *et al.*, 2013; Hulse *et al.*, 2015). The peripheral sensory nerves are compromised in these instances through a degeneration of nerve terminals (loss of intra epidermal nerve fibre innervation) in the skin (Narayanaswamy *et al.*, 2012), atrophy of the nerve trunk (shrinkage of nerve fibre diameter) (Hulse *et al.*, 2015), reduction in nerve conduction velocity (Ali *et al.*, 2014) and excitation of C nociceptor fibres (Chen & Levine, 2001). However, despite these widely diagnosed symptoms, current treatments have poor success rates and low long-term success. Recent advances in understanding diabetic pain have highlighted that neuropathic pain is associated with activation and alteration of nociceptive processing in the central nervous system. The spinal cord (Morgado *et al.*, 2010) and brain (Silva M, 2013) are hyperactivated in rodent models of diabetic neuropathic pain, with increased activity of spinal wide dynamic range neurons (Pertovaara *et al.*, 2001; Morgado *et al.*, 2010; Tan *et al.*, 2012). Microvascular degeneration is disrupted in the brain of diabetic rodents and linked to e.g. cognitive impairment (Taylor *et al.*, 2015; Hardigan *et al.*, 2016), however the impact of hyperglycaemia on the *spinal cord* microvasculature has not been reported previously. The results we present here support the conclusion that there is a widespread vasculopathy in the spinal cord, which is associated with diabetes and nociceptive processing. The reduced spinal vasculature associated with hyperglycaemia could be in part due to an increase in endothelial cell death. This compromises the function of the spinal cord microcirculation – to provide delivery of solutes as well as appropriate endothelial cell signalling – **evidenced by altered neuronal activation (c-fos staining) In all five lamina of the dorsal horn, and caspase staining in laminae 1-2** . Whether the effect on sensory neuronal function within the spinal cord is impaired as a result of decreased nutrient/oxygen delivery or through disturbance of an endothelial-neuronal-glia signalling event remains to be determined.

Multiple diabetic complications arise due to vascular pathology whereby vessel or cell function is impaired (e.g. retinopathy, nephropathy) (Gupta *et al.*, 2013; Oltean *et al.*, 2014). We have previously shown that enhanced extravasation in diabetes does occur in the peripheral nervous system, including in the DRG, and this is mediated in part through VEGF-A, as treatment with VEGF-A₁₆₅b – which can inhibit endogenous VEGF-A₁₆₅b signalling through VEGFR2, and prevent TRPA1 and TRPV1 activation in cultured DRG cells *ex vivo*, blocked the enhanced extravasation and the associated pain behaviours. In this case we

describe a reduction in solute flux in the CNS, which combined with a loss of vessels, can only be attributed to a reduction in blood flow. Evans blue extravasation is often taken as an indication of increased vascular permeability, and if blood is still flowing to the vessels where the vascular permeability increases, this is often consistent with the data. However, extravasation requires a provision of solute in the blood, and therefore is highly dependent on blood flow. A *reduction* in Evans blue extravasation in the spinal cord, where the permeability is already very low **compared with most tissues (e.g. heart, skin, DRG(Hulse *et al.*, 2015))**), can only be reasonably interpreted as an indication of reduced surface area for exchange, or reduced capillary pressure. The loss of endothelial cell staining, reduction of CD31 combined with reduced Evans blue extravasation indicates that diabetic animals have lost functional blood vessels from the spinal cord. This is an interesting comparison with most other vascular beds (including eyes (Ved *et al.*, 2017) and the peripheral nerves (Hulse *et al.*, 2015)), where diabetes results in increased solute flux, probably through increased vascular permeability (Poduslo & Curran, 1992). In peripheral tissues such as in the sensory nerve and the DRG, increased blood-nerve barrier breakdown could arise due to direct glucose toxicity upon the vessels in the epineurium. However, sensory nerves are activated following exposure to high blood glucose (Chen & Levine, 2001; Chen & Levine, 2003). Such activity would drive peripheral extravasation through release of CGRP or activation of TRPA1; which is activated in sensory nerves of diabetic rodents (Koivisto *et al.*, 2012; Hulse *et al.*, 2015). These systems are not expected to be in play in the spinal cord therefore alternative mechanisms must be in action.

In this study we find that alterations in VEGF-A/VEGFR2 **activation, as evidenced both by inhibition of VEGFR2 by VEGF-A_{165b} (Bates *et al.*, 2002; Cébe Suarez S, 2006) and VEGFR2 knockout**, is culpable in the decline in microvascular function. The VEGF-A family consists of two families of alternative spliced isoforms termed VEGF-A_{xxx}a and VEGF-A_{xxx}b (xxx denote amino acid number). These isoforms differ solely due to exon 8 splicing giving rise to differing C terminus sequences that critically alter isoform function (Harper & Bates, 2008). It has been shown that both bind to VEGFR2 with equal affinity and both possess cytoprotective actions, however VEGF-A_{xxx}a is pro-angiogenic and VEGF-A_{xxx}b is able to inhibit VEGF-A_{xxx}a mediated angiogenesis (Woolard *et al.*, 2004). Therefore VEGF-A_{165b} (acting via VEGFR2) may act as a vascular protective agent under normal conditions, preventing loss of blood vessels. Loss of VEGF-A_{165b} in diabetic spinal cord removes such protection, resulting in vascular damage; reducing both vascular integrity and function. This

concept, that loss of endogenous endothelial cell VEGFR2 mediated cytoprotection contributes to nociceptive processing, **is consistent with** the results from the endothelial specific VEGFR2 KO mice. These mice demonstrated loss of spinal cord endothelium and associated development of hyperalgesia, consistent with this concept that endogenous maintenance of VEGFR2 activity is cytoprotective to the spinal cord endothelium, **albeit that we have not directly measured spinal cord neuronal nociceptive processing**. One interesting difference between the diabetic and VEGFR2^{ECKO} animals was that whereas the diabetic rats had a reduction in larger microvessels and no change in smaller vessels, the VEGFR2^{ECKO} animals had increased number of small but reduced numbers of larger vessels. This suggests that in diabetes the loss of vessels was due to either selective loss of larger microvessels (8-12µm), or a combination of loss of all vessels and a reduction in size of the larger vessels (hypotrophy or atrophy, or vasoconstriction). In contrast the VEGFR2^{ECKO} mice had the latter effect. The functional implications would still be ischemia, but may have other subtle differences in nutrient or cell delivery. While the understanding of spinal nociceptive neuronal processing in diabetes is progressing, the contribution of microvascular alteration has never been investigated in this context. Furthermore, it has been shown that peripheral sensory neurons become sensitised upon reduced perfusion (So *et al.*, 2016), leading us to speculate that the changes in the spinal cord vascular network, as a result of hyperglycaemia that we report, could alter the microenvironment of the spinal cord sensory neurons and thus alter their level of **activation**. This would be anticipated to contribute to changes in pain perception and underlie neuropathic pain development.

The VEGFR2 knockdown was restricted to Tie2-positive cell types, which encompass the endothelial and haemopoietic cell populations. The Tie2 promoter driven transgenic models are widely used to investigate endothelial function (Makino *et al.*, 2014; Moyes *et al.*, 2014), however it has been reported that Tie2 is expressed on CD11b/CD45 positive cells. (De Palma *et al.*, 2005; Tang *et al.*, 2010). In this study there was no impact of VEGFR2 knockout on haematopoietic cell populations. An impact of this knockout system on circulatory macrophages and consequent endothelium would be expected to be minimal as there is a very small number of, if any, Tie2-positive, VEGFR2 positive macrophages (Okubo *et al.*, 2016).

People with diabetes are susceptible to neurological disease as a result of the microvascular dysfunction, impacting on motor, autonomic and cognitive systems, and increased

susceptibility to stroke and dementia (Kissela *et al.*, 2005; Tiehuis *et al.*, 2008). The VEGF-A angiogenic family has a pivotal role in managing vascular development and function (Bates, 2010), however additional evidence now also supports a fundamental role for this family in neuronal activity and survival (Verheyen *et al.*, 2012; Beazley-Long *et al.*, 2013). As a consequence, VEGF-A treatment has been trialled for peripheral diabetic neuropathy (Schratzberger *et al.*, 2001; Ropper *et al.*, 2009). Experimental hyperglycaemia leads to a reduction in VEGF-A and VEGFR2 expression in the brain; VEGF-A treatment can protect both endothelium and neuronal circuits (Taylor *et al.*, 2015). Dysfunction of the VEGF-A signalling pathway within the hippocampus impacts upon the microvasculature and leads to impaired spatial memory (Reeson *et al.*, 2015; Taylor *et al.*, 2015). A decline in vascular support and diminished vascular response in diabetes and stroke models therefore clearly affects neuronal function. However, VEGF-A_{165a} also increases vascular permeability and stimulates abnormal angiogenesis, which is detrimental in the CNS, for instance in diabetic retinopathy (Perrin *et al.*, 2005). The VEGF-A_{xxx}b isoforms, however, do not stimulate angiogenesis or increase solute flux in diabetes, in fact they can reverse it in rodent models of diabetic retinopathy (Ved *et al.*, 2017). Despite these opposing profiles both families are cytoprotective, for endothelial cells and neurons (Beazley-Long *et al.*, 2013; Oltean *et al.*, 2014; Hulse *et al.*, 2015; Vencappa *et al.*, 2015). For example, reduced VEGF-A_{165b} expression in diabetic patients is associated with reduced kidney function (Oltean *et al.*, 2014). Furthermore, treatment of diabetic mice with VEGF-A_{165b} prevented endothelial dysfunction in the kidney of these animals (Oltean *et al.*, 2014). Here we show that a reduction in VEGF-A_{165b} in the lumbar spinal cord of diabetic animals was associated with neuropathic pain and a degeneration of the spinal vasculature. Systemically reintroducing VEGF-A_{165b} prevented vascular degeneration, spinal cord neuron activation and pain. VEGF-A_{165b} was reduced in diabetic rats, despite reports that VEGF-A_{165b} is associated with inhibition of vessel growth (Woolard *et al.*, 2004). Thus VEGF-A_{165b} should be considered both an anti-angiogenic, and an endothelial survival factor –or a homeostatic counterpart to its vascular remodelling isoform, VEGF-A_{165a}. It must be noted that the actions of VEGF-A_{165b} and VEGFR2 on pain may not be restricted to vascular protection but also to effects on the spinal cord neurons, either indirectly, as VEGF-A_{165b} has been shown to inhibit peripheral sensory neuron excitability and activation of peripheral nociceptors induce c-fos expression in the dorsal horn (Hulse *et al.*, 2014), or directly through inhibiting VEGF-A_{165a} actions on spinal cord circuitry (Hulse *et al.*, 2016). **Accompanying this, neutralising endogenous VEGF-A_{165b} as well as pharmacological blockade of VEGFR2 signalling at the**

level of the spinal cord led to the development of pain (Hulse *et al.*, 2016). In addition, reductions in VEGF-A_{165b} expression and no change in total VEGF-A expression would highlight a plausible increase in alternative isoforms such as VEGF-A_{165a}, an event associated with pro-nociception and chronic pain(Hulse *et al.*, 2014; Hulse *et al.*, 2016) which cannot be ruled out in this instance.

Furthermore, some consideration does need to be made when using diabetic models such as in the instance of using STZ. STZ is a reliable rodent model of type 1 diabetic neuropathy displaying comparable pain behaviour, nerve electrophysiological parameters and nerve histology, used widely throughout the field of diabetes research. However, it is an experimental model of islet cell ablation by a toxic agent that may have other effects, so despite its widespread use in many experimental models of diabetic complications (nephropathy, neuropathy, retinopathy and others) it is still critical that these findings are reproduced in type II diabetic models, and other models of type I diabetes before being applied to people with diabetes.

These findings lead us to speculate that diabetes induced alterations of the somatosensory systems, for instance by affecting the vasculature supporting somatosensory processing at the level of the spinal cord, could be a key concept in regulating neuropathic pain and treatment of such complications (for example spinal cord vasculopathy) could provide a key target in treating diabetic neuropathic pain.

In summary, we report for the first time a significant vascular degeneration in the spinal cord of diabetic rats, which is associated with a loss of spinal VEGF-A_{165b}. This was accompanied by spinal neuron activation, indicative of altered function of the spinal cord neurons, and also by enhanced nociceptive pain behaviour. Administration of VEGF-A_{165b} alleviated both the spinal cord vascular degeneration and pain neuropathic pain. This work is complementary to previous work in the nervous system demonstrating a neuroprotective action of VEGF-A (Reeson *et al.*, 2015; Taylor *et al.*, 2015), but herein we demonstrate central effects in the spinal cord with potential contribution to the control of chronic pain development, rather than actions purely at peripheral sites. These findings provide additional avenues for to further understanding diabetic neuropathic pain and the possible mechanisms that underlie sensitisation of nociceptive pathways.

Acknowledgements

NV, RPH, MD, NBL, CV and SMB performed the animal work, the immunofluorescence, cell culture and immunoblotting in the Tumour and Vascular Biology Laboratories at the University of Nottingham. KBH generated the recombinant human VEGF-A_{165b} protein. RPH, NV, NBL, CV, MH, KBH, MD, SMB, LFD & DOB contributed to the conception or design of the work in addition to acquisition, analysis or interpretation of data for the work. RPH, NV, NBL, CV, MH, KBH, MD, SMB, LFD & DOB drafted the work or revised it critically for important intellectual content. RPH drafted the manuscript with contributions from all authors. All authors approved the final version of the manuscript, agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved all persons designated as authors qualify for authorship, and all those who qualify for authorship are listed. We also thank the Flow Cytometry Facility (UoN) and Dr David Onions.

Funding

This work was supported by Diabetes UK (11/0004192 to LFD, and 10/0004152 to DOB), MRC grant MR/K013157/1 to DOB, The Richard Bright VEGF Research Trust (UK Registered Charity 1095785), the European Foundation for the Study of Diabetes Microvascular Programme supported by Novartis to RPH and the Rosetree Trust (A1360) to RPH.

Conflict of Interest

LFD and DOB are co-inventors on patents protecting VEGF-A_{165b} and alternative RNA splicing control for therapeutic application in a number of different conditions. LFD and DOB are founder equity holders in, and DOB is a director of Exonate Ltd, a company with a focus on development of alternative RNA splicing control for therapeutic application in a number of different conditions, including diabetic complications. The authors have no other conflicts of interest to declare.

References

- Albuquerque RJ, Hayashi T, Cho WG, Kleinman ME, Dridi S, Takeda A, Baffi JZ, Yamada K, Kaneko H, Green MG, Chappell J, Wilting J, Weich HA, Yamagami S, Amano S, Mizuki N, Alexander JS, Peterson ML, Brekken RA, Hirashima M, Capoor S, Usui T, Ambati BK & Ambati J. (2009). Alternatively spliced vascular endothelial growth factor receptor-2 is an essential endogenous inhibitor of lymphatic vessel growth. *Nature medicine* **15**, 1023-1030.
- Ali S, Driscoll HE, Newton VL & Gardiner NJ. (2014). Matrix metalloproteinase-2 is downregulated in sciatic nerve by streptozotocin induced diabetes and/or treatment with minocycline: Implications for nerve regeneration. *Exp Neurol* **261**, 654-665.
- Bates D, Cui T, Doughty J, Winkler M, Sugiono M, Shields J, Peat D, Gillatt D & Harper S. (2002). VEGF165b, an inhibitory splice variant of vascular endothelial growth factor, is down-regulated in renal cell carcinoma. *Cancer research* **62**, 4123-4131.
- Bates DO. (2010). Vascular endothelial growth factors and vascular permeability. *Cardiovascular research* **87**, 262-271.
- Beazley-Long N, Ashby WR, Moss CE, Bestall SM, Almahasneh F, Durrant AeM, Benest AV, Blackley Z, Ballmer-Hofer K, Hirashima M, Hulse RP, Bates DO & Donaldson LF. (2018). VEGFR2 promotes central endothelial activation and the spread of pain in inflammatory arthritis. *Brain, Behavior, and Immunity In Press*.
- Beazley-Long N, Hua J, Jehle T, Hulse RP, Dersch R, Lehrling C, Bevan H, Qiu Y, Lagreze WA, Wynick D, Churchill AJ, Kehoe P, Harper SJ, Bates DO & Donaldson LF. (2013). VEGF-A165b Is an Endogenous Neuroprotective Splice Isoform of Vascular Endothelial Growth Factor A in Vivo and in Vitro. *The American journal of pathology* **183**, 918-929.
- Beggs S, Liu XJ, Kwan C & Salter MW. (2010). Peripheral nerve injury and TRPV1-expressing primary afferent C-fibers cause opening of the blood-brain barrier. *Molecular pain* **6**, 74.
- Biessels G, Bril V, Calcutt N, Cameron N, Cotter M, Dobrowsky R, Feldman E, Fernyhough P, Jakobsen J, Malik R, Mizisin A, Oates P, Obrosova I, Pop-Busui R, Russell J, Sima A, Stevens M, Schmidt R, Tesfaye S, Veves A, Vinik A, Wright D, Yagihashi S, Yorek M, Ziegler D & Zochodne D.

- (2014). Phenotyping animal models of diabetic neuropathy: a consensus statement of the diabetic neuropathy study group of the EASD (Neurodiab). *J Peripher Nerv Syst* **19**, 77-87.
- Cai J & Boulton M. (2002). The pathogenesis of diabetic retinopathy: old concepts and new questions. *Eye* **16**, 242-260.
- Calcutt NA. (2004). Modeling diabetic sensory neuropathy in rats. *Methods in molecular medicine* **99**, 55-65.
- Cébe Suarez S PM, Cariolato L, Arn S, Hoffmann U, Bogucki A, Manlius C, Wood J, Ballmer-Hofer K. (2006). A VEGF-A splice variant defective for heparan sulfate and neuropilin-1 binding shows attenuated signaling through VEGFR-2. *Cellular and molecular life sciences : CMLS* **63**, 2067-2077.
- Chen X & Levine JD. (2001). Hyper-responsivity in a subset of c-fiber nociceptors in a rat model of painful diabetic neuropathy in the rat. *Neuroscience* **102**, 185-192.
- Chen X & Levine JD. (2003). Altered temporal pattern of mechanically evoked C-fiber activity in a model of diabetic neuropathy in the rat. *Neuroscience* **121**, 1007-1015.
- Chowdhury SR, Saleh A, Akude E, Smith DR, Morrow D, Tessler L, Calcutt NA & Fernyhough P. (2014). Ciliary neurotrophic factor reverses aberrant mitochondrial bioenergetics through the JAK/STAT pathway in cultured sensory neurons derived from streptozotocin-induced diabetic rodents. *Cellular and molecular neurobiology* **34**, 643-649.
- Costigan M, Moss A, Latremoliere A, Johnston C, Verma-Gandhu M, Herbert T, Barrett L, Brenner G, Vardeh D, Woolf C & Fitzgerald M. (2009). T-cell infiltration and signaling in the adult dorsal spinal cord is a major contributor to neuropathic pain-like hypersensitivity. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **28**, 14415-14422.
- De Palma M, Venneri MA, Galli R, Sergi L, Politi LS, Sampaolesi M & Naldini L. (2005). Tie2 identifies a hematopoietic lineage of proangiogenic monocytes required for tumor vessel formation and a mesenchymal population of pericyte progenitors. *Cancer cell* **8**, 211-226.
- Drake R, Hulse R, Lumb B & Donaldson L. (2014). The degree of acute descending control of spinal nociception in an area of primary hyperalgesia is dependent on the peripheral domain of afferent input. *The Journal of physiology* **592**, 3611-3624.

Grundy D. (2015). Principles and standards for reporting animal experiments in The Journal of Physiology and Experimental Physiology. *J Physiol* **593**, 2547-2549.

Gupta N, Mansoor S, Sharma A, Sapkal A, Sheth J, Falatoonzadeh P, Kuppermann B & Kenney M. (2013). Diabetic retinopathy and VEGF. *The open ophthalmology journal* **7**, 4-10.

Hardigan T, Ward R & Ergul A. (2016). Cerebrovascular complications of diabetes: focus on cognitive dysfunction. *Clin Sci (Lond)* **130**, 1807-1822.

Hargreaves K, Dubner R, Brown F, Flores C & Joris J. (1988). A new and sensitive method for measuring thermal nociception in cutaneous hyperalgesia. *Pain* **32**, 77-88.

Harper S & Bates D. (2008). VEGF-A splicing: the key to anti-angiogenic therapeutics? *Nature reviews Cancer* **8**, 880-887.

Hirschberg S, Li Y, Randall A, Kremer EJ & Pickering AE. (2017). Functional dichotomy in spinal- vs prefrontal-projecting locus coeruleus modules splits descending noradrenergic analgesia from ascending aversion and anxiety in rats. *Elife* **6**.

Hsieh MT, Donaldson LF & Lumb BM. (2015). Differential contributions of A- and C-nociceptors to primary and secondary inflammatory hypersensitivity in the rat. *Pain* **156**, 1074-1083.

Hulse RP, Beazley-Long N, Hua J, Kennedy H, Prager J, Bevan H, Qiu Y, Fernandes ES, Gammons MV, Ballmer-Hofer K, Gittenberger de Groot AC, Churchill AJ, Harper SJ, Brain SD, Bates DO & Donaldson LF. (2014). Regulation of alternative VEGF-A mRNA splicing is a therapeutic target for analgesia. *Neurobiology of disease* **71**, 245-259.

Hulse RP, Beazley-Long N, Ved N, Bestall SM, Riaz H, Singhal P, Ballmer Hofer K, Harper SJ, Bates D & Donaldson LF. (2015). Vascular endothelial growth factor-A165b prevents diabetic neuropathic pain and sensory neuronal degeneration. *Clin Sci* **129**, 741-756.

Hulse RP, Drake RA, Bates DO & Donaldson LF. (2016). The control of alternative splicing by SRSF1 in myelinated afferents contributes to the development of neuropathic pain. *Neurobiology of disease* **96**, 186-200.

Hunt SP PA, Evan G. (1987). Induction of c-fos-like protein in spinal cord neurons following sensory stimulation. *Nature* **328**, 632-634.

Kalynovska N, Adamek P & Palecek J. (2017). TRPV1 receptors contribute to mediate paclitaxel-induced c-Fos expression in spinal cord dorsal horn neurons. *Physiol Res* **66**, 549-552.

Khasabov SG, Malecha P, Noack J, Tabakov J, Giesler GJ & Simone DA. (2017). Hyperalgesia and sensitization of dorsal horn neurons following activation of NK-1 receptors in the rostral ventromedial medulla. *Journal of neurophysiology* **118**, 2727-2744.

Kikuchi R, Nakamura K, MacLauchlan S, Ngo DT, Shimizu I, Fuster JJ, Katanasaka Y, Yoshida S, Qiu Y, Yamaguchi TP, Matsushita T, Murohara T, Gokce N, Bates DO, Hamburg NM & Walsh K. (2014). An antiangiogenic isoform of VEGF-A contributes to impaired vascularization in peripheral artery disease. *Nature medicine* **20**, 1464-1471.

Kissela B, Khoury J, Kleindorfer D, Woo D, Schneider A, Alwell K, Miller R, Ewing I, Moomaw C, Szaflarski J, Gebel J, Shukla R & Broderick J. (2005). Epidemiology of ischemic stroke in patients with diabetes: the greater Cincinnati/Northern Kentucky Stroke Study. *Diabetes care* **28**, 355-359.

Koivisto A, Hukkanen M, Saarnilehto M, Chapman H, Kuokkanen K, Wei H, Viisanen H, Akerman KE, Lindstedt K & Pertovaara A. (2012). Inhibiting TRPA1 ion channel reduces loss of cutaneous nerve fiber function in diabetic animals: sustained activation of the TRPA1 channel contributes to the pathogenesis of peripheral diabetic neuropathy. *Pharmacological research : the official journal of the Italian Pharmacological Society* **65**, 149-158.

Latremoliere A WC. (2009). Central sensitization: a generator of pain hypersensitivity by central neural plasticity. *J Pain* **10**, 895-926.

Lee-Kubli CA CN. (2013). Altered rate-dependent depression of the spinal H-reflex as an indicator of spinal disinhibition in models of neuropathic pain. *Pain* **155**, 250-260.

Makino K, Jinnin M, Aoi J, Kajihara I, Makino T, Fukushima S, Sakai K, Nakayama K, Emoto N, Yanagisawa M & Ihn H. (2014). Knockout of endothelial cell-derived endothelin-1 attenuates skin fibrosis but accelerates cutaneous wound healing. *PLoS One* **9**, e97972.

- Manetti M, Guiducci S, Romano E, Ceccarelli C, Bellando-Randone S, Conforti ML, Ibba-Manneschi L & Matucci-Cerinic M. (2011). Overexpression of VEGF165b, an inhibitory splice variant of vascular endothelial growth factor, leads to insufficient angiogenesis in patients with systemic sclerosis. *Circ Res* **109**, e14-26.
- Morgado C, Terra PP & Tavares I. (2010). Neuronal hyperactivity at the spinal cord and periaqueductal grey during painful diabetic neuropathy: effects of gabapentin. *European journal of pain* **14**, 693-699.
- Moyes AJ, Khambata RS, Villar I, Bubb KJ, Baliga RS, Lumsden NG, Xiao F, Gane PJ, Rebstock AS, Worthington RJ, Simone MI, Mota F, Rivilla F, Vallejo S, Peiro C, Sanchez Ferrer CF, Djordjevic S, Caulfield MJ, MacAllister RJ, Selwood DL, Ahluwalia A & Hobbs AJ. (2014). Endothelial C-type natriuretic peptide maintains vascular homeostasis. *The Journal of clinical investigation* **124**, 4039-4051.
- Narayanaswamy H, Facer P, Misra VP, Timmers M, Byttebier G, Meert T & Anand P. (2012). A longitudinal study of sensory biomarkers of progression in patients with diabetic peripheral neuropathy using skin biopsies. *Journal of clinical neuroscience : official journal of the Neurosurgical Society of Australasia* **19**, 1490-1496.
- Ngo DT, Farb MG, Kikuchi R, Karki S, Tiwari S, Bigornia SJ, Bates DO, LaValley MP, Hamburg NM, Vita JA, Hess DT, Walsh K & Gokce N. (2014). Antiangiogenic actions of vascular endothelial growth factor-A165b, an inhibitory isoform of vascular endothelial growth factor-A, in human obesity. *Circulation* **130**, 1072-1080.
- Okubo M, Kioi M, Nakashima H, Sugiura K, Mitsudo K, Aoki I, Taniguchi H & Tohnai I. (2016). M2-polarized macrophages contribute to neovasculogenesis, leading to relapse of oral cancer following radiation. *Scientific reports* **6**, 27548.
- Oltean S, Qiu Y, Ferguson J, Stevens M, Neal C, Russell A, Kaura A, Arkill K, Harris K, Symonds C, Lacey K, Wijeyaratne L, Gammons M, Wylie E, Hulse R, Alsop C, Cope G, Damodaran G, Betteridge K, Ramnath R, Satchell S, Foster R, Ballmer-Hofer K, Donaldson L, Barratt J, Baelde H, Harper S, Bates D & Salmon A. (2014). Vascular Endothelial Growth Factor-A165b Is Protective and Restores Endothelial Glycocalyx in Diabetic Nephropathy. *J Am Soc Nephrol* **26**, 1889-1904.

- Perrin R, Konopatskaya O, Qiu Y, Harper S, Bates D & Churchill A. (2005). Diabetic retinopathy is associated with a switch in splicing from anti- to pro-angiogenic isoforms of vascular endothelial growth factor. *Diabetologia* **48**.
- Pertovaara A, Wei H, Kalmari J & Ruotsalainen M. (2001). Pain behavior and response properties of spinal dorsal horn neurons following experimental diabetic neuropathy in the rat: modulation by nitecapone, a COMT inhibitor with antioxidant properties. *Exp Neurol* **167**, 425-434.
- Poduslo J & Curran G. (1992). Increased permeability across the blood-nerve barrier of albumin glycated in vitro and in vivo from patients with diabetic polyneuropathy. *Proceedings of the National Academy of Sciences of the United States of America* **89**, 2218-2222.
- Pritchard-Jones RO, Dunn DB, Qiu Y, Varey AH, Orlando A, Rigby H, Harper SJ & Bates DO. (2007). Expression of VEGF(xxx)b, the inhibitory isoforms of VEGF, in malignant melanoma. *British journal of cancer* **97**, 223-230.
- Reeson P, Tennant KA, Gerrow K, Wang J, Weiser Novak S, Thompson K, Lockhart KL, Holmes A, Nahirney PC & Brown CE. (2015). Delayed inhibition of VEGF signaling after stroke attenuates blood-brain barrier breakdown and improves functional recovery in a comorbidity-dependent manner. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **35**, 5128-5143.
- Reichling DB & Levine JD. (2009). Critical role of nociceptor plasticity in chronic pain. *Trends in neurosciences* **32**, 611-618.
- Ropper AH, Gorson KC, Gooch CL, Weinberg DH, Pieczek A, Ware JH, Kershen J, Rogers A, Simovic D, Schratzberger P, Kirchmair R & Losordo D. (2009). Vascular endothelial growth factor gene transfer for diabetic polyneuropathy: a randomized, double-blinded trial. *Ann Neurol* **65**, 386-393.
- Said G. (2007). Diabetic neuropathy--a review. *Nature clinical practice Neurology* **3**, 331-340.
- Schratzberger P, Walter DH, Rittig K, Bahlmann FH, Pola R, Curry C, Silver M, Krainin JG, Weinberg DH, Ropper AH & Isner JM. (2001). Reversal of experimental diabetic neuropathy by VEGF gene transfer. *The Journal of clinical investigation* **107**, 1083-1092.

- Silva M AD, Almeida A, Tavares I, Pinto-Ribeiro F, Morgado C. (2013). Pronociceptive changes in the activity of rostroventromedial medulla (RVM) pain modulatory cells in the streptozotocin-diabetic rat. *Brain research bulletin* **96**, 39-44.
- Sison K, Eremina V, Baelde H, Min W, Hirashima M, Fantus IG & Quaggin SE. (2010). Glomerular structure and function require paracrine, not autocrine, VEGF-VEGFR-2 signaling. *J Am Soc Nephrol* **21**, 1691-1701.
- So K, Tei Y, Zhao M, Miyake T, Hiyama H, Shirakawa H, Imai S, Mori Y, Nakagawa T, Matsubara K & Kaneko S. (2016). Hypoxia-induced sensitisation of TRPA1 in painful dysesthesia evoked by transient hindlimb ischemia/reperfusion in mice. *Scientific reports* **6**, 23261.
- Tan AM, Samad OA, Fischer TZ, Zhao P, Persson AK & Waxman SG. (2012). Maladaptive dendritic spine remodeling contributes to diabetic neuropathic pain. *The Journal of neuroscience : the official journal of the Society for Neuroscience* **32**, 6795-6807.
- Tang Y, Harrington A, Yang X, Friesel RE & Liaw L. (2010). The contribution of the Tie2+ lineage to primitive and definitive hematopoietic cells. *Genesis* **48**, 563-567.
- Taylor S, Trudeau D, Arnold B, Wang J, Gerrow K, Summerfeldt K, Holmes A, Zamani A, Brocardo P & Brown C. (2015). VEGF can protect against blood brain barrier dysfunction, dendritic spine loss and spatial memory impairment in an experimental model of diabetes. *Neurobiology of disease* **78**, 11.
- Tesfaye S, Boulton A & Dickenson A. (2013). Mechanisms and management of diabetic painful distal symmetrical polyneuropathy. *Diabetes care* **36**, 2456-2465.
- Tesfaye S, Selvarajah D, Gandhi R, Greig M, Shillo P, Fang F & Wilkinson ID. (2016). Diabetic peripheral neuropathy may not be as its name suggests: evidence from magnetic resonance imaging. *Pain* **157 Suppl 1**, S72-80.
- Tesfaye S, Vileikyte L, Rayman G, Sindrup S, Perkins B, Baconja M, Vinik A & Boulton A. (2011). Painful diabetic peripheral neuropathy: consensus recommendations on diagnosis, assessment and management. *Diabetes/metabolism research and reviews* **27**, 629-638.

- Tiehuis AM, van der Graaf Y, Visseren FL, Vincken KL, Biessels GJ, Appelman AP, Kappelle LJ, Mali WP & Group SS. (2008). Diabetes increases atrophy and vascular lesions on brain MRI in patients with symptomatic arterial disease. *Stroke; a journal of cerebral circulation* **39**, 1600-1603.
- Vandal M, White PJ, Tremblay C, St-Amour I, Chevrier G, Emond V, Lefrancois D, Virgili J, Planel E, Giguere Y, Marette A & Calon F. (2014). Insulin reverses the high-fat diet-induced increase in brain Abeta and improves memory in an animal model of Alzheimer disease. *Diabetes* **63**, 4291-4301.
- Ved N, Hulse RP, Bestall SM, Donaldson LF, Bainbridge JW & Bates DO. (2017). Vascular endothelial growth factor-A165b ameliorates outer-retinal barrier and vascular dysfunction in the diabetic retina. *Clin Sci (Lond)* **131**, 1225-1243.
- Vencappa S, Donaldson LF & Hulse RP. (2015). Cisplatin induced sensory neuropathy is prevented by vascular endothelial growth factor-A. *American journal of translational research* **7**, 1032-1044.
- Verheyen A, Peeraer E, Nuydens R, Dhondt J, Poesen K, Pintelon I, Daniels A, Timmermans JP, Meert T, Carmeliet P & Lambrechts D. (2012). Systemic anti-vascular endothelial growth factor therapies induce a painful sensory neuropathy. *Brain : a journal of neurology* **135**, 2629-2641.
- Winkler EA, Nishida Y, Sagare AP, Rege SV, Bell RD, Perlmutter D, Sengillo JD, Hillman S, Kong P, Nelson AR, Sullivan JS, Zhao Z, Meiselman HJ, Wenby RB, Soto J, Abel ED, Makshanoff J, Zuniga E, De Vivo DC & Zlokovic BV. (2015). GLUT1 reductions exacerbate Alzheimer's disease vasculo-neuronal dysfunction and degeneration. *Nature neuroscience* **18**, 521-530.
- Woolard J, Wang WY, Bevan HS, Qiu Y, Morbidelli L, Pritchard-Jones RO, Cui TG, Sugiono M, Waite E, Perrin R, Foster R, Digby-Bell J, Shields JD, Whittles CE, Mushens RE, Gillatt DA, Ziche M, Harper SJ & Bates DO. (2004). VEGF165b, an inhibitory vascular endothelial growth factor splice variant: mechanism of action, in vivo effect on angiogenesis and endogenous protein expression. *Cancer research* **64**, 7822-7835.
- Xu Q, Qaum T & Adamis AP. (2001). Sensitive blood-retinal barrier breakdown quantitation using Evans blue. *Invest Ophthalmol Vis Sci* **42**, 789-794.

Figure 1 – Diabetes-induced neuropathic pain is associated with a reduction in spinal cord vasculature and a decrease in VEGF-A_{165b} expression. [A] Diabetes resulted in a reduction in mechanical withdrawal threshold measured by pincher, when compared with naïve age matched animals (**p*<0.05, n=5/group). [B] Blood vessels identified (IB₄) in the deeper laminar layers of the spinal cord (layers III-VI) of the naïve animal [C] with a decline in vascular staining in the diabetic animal. [D] There was a reduction in blood vessel (CD31/IB₄+ve) number (**p*<0.05, n=4/group) [E= naïve, F = Diabetic; reduced diameter (**p*<0.05, n=4/group) as well as diameter [G] in the lumbar spinal cord in diabetic animals compared with naïve controls. [H] Immunoblot of panVEGF, VEGF-A_{165b} and actin in lysates from spinal cord of normal and diabetic animals. [I]. Densitometry analysis demonstrates no change in pan-VEGF-A expression and a decrease in VEGF-A_{165b} expression in diabetic lumbar spinal cord versus naïve animals. (**p*<0.05, n=5/group) (B&C Scale bar= 40µm, E&F Scale bar= 20µm).




Figure 2 – Diabetes induced vascular impairment in the spinal cord. [A] Isolated spinal cord endothelial cells demonstrated increased cell death in 50mM glucose when compared with 50mM mannitol and 5mM glucose (* $p < 0.05$, *** $p < 0.001$). [A] VEGF- A_{165b} treatment prevented high glucose-induced endothelial cell death (** $p < 0.01$). [B] IB₄ stained vasculature in the spinal cord of naïve age matched controls was compared with that in [C] diabetic+vehicle (Arrow heads = vessels smaller than 6 μ m) and [D] diabetic +VEGF- A_{165b} . [E] There was a significant reduction in total volume of the microvasculature in the spinal cord of the diabetic+vehicle group in addition to [F] a reduction in vessel diameter compared with naïve controls (** $p < 0.01$, *** $p < 0.001$, $n = 4$ /group). [E-F] VEGF- A_{165b} treatment prevented the diabetes-induced vascular degeneration in the lumbar spinal cord (** $p < 0.01$, $n = 4$ /group). [G] VEGF- A_{165b} treatment also prevents the diabetes induced decrease in larger and intermediate microvessels. (B-D Scale bar = 25 μ m).




Figure 3 – Diabetes induced degeneration of the endothelium. Immunoblots using dual colour far red imaging for endothelial markers ([A] VE-Cadherin, [A] CD31 and [B] occludin and [B] actin) demonstrated [C] a non-significant reduction in VE-cadherin expression and significant reductions in [D] CD31 and [E] occludin in the diabetic+vehicle group compared with naïve and diabetic+VEGF-A₁₆₅b animals (*p<0.05,n=5).




Figure 4 – Reduced vascular functionality in the spinal cord of diabetic rats. There was a significant reduction in Evans blue solute flux in the lumbar spinal cord of diabetic animals after [A] 1 week (naïve;n=9, diabetes;n= 12, *** $p<0.001$) and [B] 8 weeks (* $p<0.05$,n=4/5 /group). VEGF- A_{165b} treatment prevented the diabetes-induced reduction in solute flux within the lumbar spinal cord at 8 weeks (* $p<0.05$,n=4/5 /group). [C] There was no change in solute flux in the brain of any treatment group (naïve vs diabetic+vehicle = $p=0.52$; diabetic+vehicle vs diabetic +VEGF- A_{165b} = $p=>0.99$; n=4/5 /group).




Figure 5 – Diabetes induced dysfunction of microvasculature in the spinal cord and neuropathic pain is reversed by VEGF-A₁₆₅b. [A] Diabetic+vehicle animals demonstrated a decrease in mechanical withdrawal threshold (vF hairs) and [B] reduced withdrawal latency to heat compared with both naïve and diabetic+VEGF-A₁₆₅b treated groups (** p<0.01, *** p<0.001 naïve vs diabetic+vehicle; # p<0.001 diabetic +VEGF-A₁₆₅b vs diabetic+vehicle, n=5 per group). **Arrow highlight onset of VEGF-A₁₆₅b treatment**, [C] Cleaved caspase-3 (red=CC3) and sensory neuron (green=NeuN) staining in the spinal cord (Scale bar=40µm). [D] There was an increase in cleaved caspase-3 expression in sensory neurons in the superficial lamina (I&II) of the dorsal horn of the spinal cord in the diabetic + vehicle groups compared with naïve age matched controls and VEGF-A₁₆₅b treated diabetic animals. [E] There is an increased number of CC3 positive dorsal horn sensory neurons (arrows in E) in the dorsal horn of the spinal cord in the diabetic + vehicle groups compared with naïve age matched controls and VEGF-A₁₆₅b treated diabetic animals. This is graphically represented in [F] (*p<0.05, n=4).




Figure 6 – Diabetes induced hyperactivity in sensory neurons of the dorsal horn in the spinal cord. [A-B] Immunoreactivity of a marker of central sensitisation, c-fos (red; NeuN green), was increased in sensory neurons in the dorsal horn of the spinal cord in diabetic+vehicle animals versus naïve animals. This was reduced by VEGF-A₁₆₅b treatment. [C] There was an increase of c-fos expression in sensory neurons in all lamina of the dorsal horn (I-V) in the diabetic+vehicle group when compared to naïve and diabetic+VEGF-A₁₆₅b (*p<0.05, **p<0.001, ***p<0.001; n=4).




Figure 7 – Tamoxifen induced VEGFR2 knockout. VEGFR2 was inducibly knocked out in endothelial cells by crossing *vegfr2^{fl/fl}* with *Tie2Cre^{ERT2}* mice and treating with tamoxifen (*VEGFR2^{ECKO}*). [A] In lung tissue, VEGFR2 protein was detected in tamoxifen treated *vegfr2^{fl/fl}* mice lacking Cre (CTL), but not in *VEGFR2^{ECKO}* mice [B] Immunoblot densitometry demonstrated reduced VEGFR2 protein in the *VEGFR2^{ECKO}* mice compared with controls (* $p < 0.05$, =6 per group). [C-D] VE-cadherin expression was also reduced in lung tissue of the *VEGFR2^{ECKO}* mice (* $p < 0.05$, n=6 per group). [E] There was a reduction in VEGFR2 in the spinal cord of the *VEGFR2^{ECKO}* mice when compared to CTL mice (* $p < 0.05$, n=4 per group, this was accompanied by reductions in endothelial markers [F] VE-cadherin and [G] CD31 (** $p < 0.01$, n=4 per group. [H] Western blot of the lumbar spinal cord from *VEGFR2^{ECKO}* mice demonstrating [I] a reduction in VE-Cadherin expression when compared to CTL mice.




Figure 8 - Inducible endothelial cell *vegfr2* KO caused microvasculature loss in the dorsal horn of the lumbar region of the spinal and hyperalgesia. Representative images from the microvessels in [A lower power; B high power] *vegfr2^{fl/fl} Tie2Cre^{ERT2}* positive mice + vehicle (Vehicle) and [D-F] *VEGFR2^{ECKO}*. *VEGFR2^{ECKO}* mice had a significant reduction in microvasculature [C] volume and [D] diameter compared with controls (CTL and vehicle). (** $p < 0.000$, * $p < 0.01$, comparison between Vehicle and CTL vessel diameter $p = 0.5391$). [E] *VEGFR2^{ECKO}* mice had an increased number of smaller microvessels versus other experimental groups. [F] *VEGFR2^{ECKO}* mice showed a reduced withdrawal latency to heat when compared to control mice (** $p < 0.001$ comparison made at day 11 between CTL ($p = 0.003$), Vehicle ($P = 0.001$) and CTL +Vehicle ($P = 0.0008$) against *VEGFR2^{ECKO}* mice, $n = 10/11$ per group).




Figure 9 – Tamoxifen did not induce a loss in haematopoietic cells in the mouse spleen [A-B] Cells isolated from mouse spleens were analysed using flow cytometry to identify [C] F4/80 and [D] CD11b cell populations. In VEGFR2^{ecKO} + tamoxifen mice there were no changes in cell number in [E] F4/80 ($p=0.630$), [F] CD11b ($p>0.99$) and [G] F4/80/CD11b ($p>0.99$) cell populations or [H&I] median fluorescent intensity (F4/80= $p=0.11$; CD11b= 0.11) when compared with tamoxifen treated Tie2CRE mice ($n=4$ /group).

