

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16

**A life history perspective on athletes with low energy availability**

*Meghan K Shirley<sup>1</sup>; Daniel P. Longman<sup>2</sup>; Kirsty J. Elliott-Sale<sup>3</sup>; Anthony C. Hackney<sup>4</sup>; Craig Sale<sup>3</sup>; Eimear Dolan<sup>5</sup>*

- 1: Division of GI, Hepatology and Nutrition, The Children’s Hospital of Philadelphia, Philadelphia, PA, United States.
- 2: School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, United Kingdom.
- 3: Musculoskeletal Physiology Research Group, Sport Health and Performance Enhancement Research Centre, School of Science and Technology, Nottingham Trent University, Nottingham, United Kingdom.
- 4: Department of Exercise & Sport Science, Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC, United States.
- 5: Applied Physiology & Nutrition Research Group; School of Physical Education and Sport; Rheumatology Division; Faculdade de Medicina FMUSP, Universidade de Sao Paulo, Sao Paulo, SP, Brazil.

**Corresponding author:** Dr. Eimear Dolan ([eimeardol@gmail.com](mailto:eimeardol@gmail.com))

Meghan K. Shirley	ORCID ID: 0000-0001-8800-0707
Daniel P. Longman	ORCID ID: 0000-003-3025-7053
Kirsty J. Elliott-Sale	ORCID ID: 0000-0003-1122-5099
Anthony C. Hackney	ORCID ID: 0000-0002-6607-1472
Craig Sale	ORCID ID: 0000-0002-5816-4169
Eimear Dolan	ORCID ID: 0000-0002-1018-7601

17 **Abstract**

18 The energy costs of athletic training can be substantial, and deficits arising from costs unmet by adequate energy  
19 intake, leading to a state of low energy availability, may adversely impact athlete health and performance. Life  
20 history theory is a branch of evolutionary theory which recognizes that the way the body uses energy - and  
21 responds to low energy availability - is an evolved trait. Energy is a finite resource that must be distributed  
22 throughout the body to simultaneously fuel all biological processes. When energy availability is low, insufficient  
23 energy may be available to equally support all processes. As energy used for one function cannot be used for  
24 others, energetic “trade-offs” will arise. Biological processes offering the greatest immediate survival value will  
25 be protected, even if this results in energy being diverted away from others, potentially leading to their  
26 downregulation. Athletes with low energy availability provide a useful model for anthropologists investigating  
27 the biological trade-offs that occur when energy is scarce, while the broader conceptual framework provided by  
28 life history theory may be useful to sport and exercise researchers who investigate the influence of low energy  
29 availability on athlete health and performance. The goals of this review are: 1) to describe the core tenets of life  
30 history theory; 2) consider trade-offs that might occur in athletes with low energy availability in the context of  
31 four broad biological areas: reproduction, somatic maintenance, growth and immunity; and 3) use this  
32 evolutionary perspective to consider potential directions for future research.

33

34 **Key Points:**

- 35       ➤ Life history theory is a branch of evolutionary theory that describes how finite energy resources are  
36       distributed among competing biological processes. “Trade-offs” are predicted to arise because energy  
37       used to support one process cannot be used for others. Energy resources are allocated to tissues and  
38       functions in a hierarchical manner, with priority afforded to those that offer the greatest immediate  
39       survival value.
- 40       ➤ Because energetic trade-offs among biological processes are predicted to be heightened under conditions  
41       of scarcity, a life history lens is particularly relevant for athletes who may have low energy availability  
42       due to high exercise training energy expenditure, inadequate energy intake (resulting from conscious or  
43       inadvertent restriction), or a combination of both.
- 44       ➤ Energetic trade-offs are context specific and may be impacted by a range of factors, including sex, life-  
45       stage, training and health status, dietary composition and the length and severity of exposure to low  
46       energy availability.

47

48 **Table 1: Operational Definition of Terms** (*italicized in main text at first mention*)

Energy availability (EA)	Dietary energy intake minus energy expended in training, or the amount of energy available for biological processes after the demands of exercise training have been met [1,2]. Herein we refer to energy availability as “adequate” when sufficient to sustain usual functioning of all body systems, and “low” when energy availability cannot adequately sustain all systems.
Energy balance (EB)	Dietary energy intake minus total energy expenditure. An individual may be in energy balance while simultaneously experiencing low energy availability, as a downregulation of physiological systems can reduce total energy expenditure to match lower energy intakes [1,2].
Inadequate energy intake	A dietary energy intake that is insufficient to meet the physiological requirements of the body. Inadequate intakes may be conscious ( <i>e.g.</i> , intentional restricted intake to achieve a desired body composition), or inadvertent ( <i>e.g.</i> , unintended undereating or alimentary limitation, whereby individuals cannot consume sufficient quantities of food to meet energetic demands of training).
Evolutionary anthropology/biology	Evolutionary anthropology refers to the study of human evolution and variation, including in relation to human physiology and behaviour [3]. Evolutionary biology investigates the biological basis of evolution across species.
Life history theory	A branch of evolutionary theory describing the competitive allocation of limited energy resources among physiological functions [4,5].
Evolutionary fitness	The ability to successfully capture and use available resources to reproduce and pass genetic material to the next generation.
Phenotypic plasticity	The capacity to alter aspects of morphology, physiology, and behaviour in response to varying environmental circumstances.
Energy “trade-offs”	A reduction in energy allocation to one function as a result of energy being allocated to another function (considering that energy resources are finite).
Total daily energy expenditure (TDEE)	The total amount of energy expended in a day, which for adults will comprise basal metabolism, voluntary and involuntary activity, and the thermic effect of food.

49  
50  
51  
52  
53

54 **1. Introduction**

55 Meeting training demands with adequate energy intake can be challenging for many athletes, and evidence  
56 suggests that *low energy availability* (LEA; Table 1) is a common occurrence, with potentially negative  
57 consequences for health and performance [6–9]. These consequences have been conceptualized primarily in three  
58 models - the Female Athlete Triad [10], Relative Energy Deficiency in Sport (RED-S) [11], and the Exercise-  
59 Hypogonadal Male Condition [9] models. Despite ongoing investigation and important advances in knowledge,  
60 gaps in understanding of the physiology and pathophysiology of athletes with LEA remain, and these must be  
61 filled to progress screening, treatment, and return-to-play recommendations.

62 *Evolutionary biologists* recognize that the biological response to LEA is an evolved trait, and have long examined  
63 how finite energy is distributed among competing physiological processes in humans and other species within a  
64 theoretical framework called *life history theory* [4,5,12–15]. Classically used to understand species-level  
65 differences in the pattern and timing of important biological events (*e.g.*, maturation, reproduction, and death),  
66 life history theory's conceptual framework has increased understanding of which physiological functions are  
67 prioritized or sacrificed in humans under varying environmental circumstances, including energetic stressors such  
68 as LEA. This has broad implications for health in human populations and may hold specific relevance for athletes,  
69 many of whom struggle to achieve adequate energy availability due to high training energy expenditure and/or  
70 *inadequate energy intake* (which may occur via conscious restriction, reaching alimentary limits [16], or due to  
71 inadvertent under-eating [8,17]). Among athletes, those with particularly strenuous training schedules may incur  
72 extreme energetic costs that can increase their susceptibility to LEA. Many evolutionary anthropologists use  
73 exercise, sport and physical activity to investigate and inform theories of human evolution and adaptation [18–  
74 22]. Similarly, sport and exercise scientists may find the overarching conceptual framework of life history theory  
75 useful to consider the biological basis and variability of the human body's response to LEA. The aim of this  
76 review, therefore, is 1) to describe the core tenets of life history theory; 2) consider trade-offs that might occur in  
77 athletes with LEA in the context of four broad biological areas that are commonly considered within the life  
78 history literature, *i.e.*, reproduction, somatic maintenance, growth and immunity; and 3) use this evolutionary  
79 perspective to consider potential directions for future research.

80


81 **2. Brief Background to Evolutionary Life History Theory:**

82 The notion of 'competition' features prominently in Charles Darwin's theory of evolution by natural selection  
83 [23]. Darwin and his contemporary, Alfred Russel Wallace, described competition amongst individuals for the  
84 resources needed to successfully pass genetic material to subsequent generations - this constitutes '*fitness*' in  
85 evolutionary terms [24]. Life history theory, a branch of evolutionary theory, also predicts competition at an intra-  
86 individual level. This theory considers energy to be a finite resource that is competitively allocated between four  
87 principal areas: reproduction, growth, somatic maintenance, and immune defence [25–31]. Energy can also be  
88 used for physical activity, or stored in adipose tissue, or as liver or muscle glycogen or triglycerides. A key tenet  
89 of life history theory is that energy used for one function cannot be used for others, *i.e.*, energy resources are  
90 "*traded off*" among competing functions. Notably, competition among functions will be heightened during  
91 conditions of scarcity [30,32], as may be the case for athletes with LEA.

92 Like all aspects of human biology, the physiological mechanisms responsible for energy allocation among  
93 competing functions were shaped by humans' evolutionary journey. The evolutionary history of our species  
94 involved repeated cycles of colonization and dispersal, characterized by periods of food insecurity and other  
95 ecological challenges [33–35]. In the language of evolutionary biology, such challenges were 'selective pressures'  
96 that shaped the development of adaptive mechanisms to successfully survive periods of biological stress, including  
97 that caused by LEA. One such adaptive mechanism allows for an internal reallocation of energy during times of  
98 scarcity toward physiological processes providing the greatest immediate survival value. This capacity to  
99 dynamically redistribute energy resources derives from *phenotypic plasticity* [34,36,37], whereby morphological,  
100 physiological, and behavioural traits vary in response to changing environmental circumstances and stressors  
101 without alterations to the genetic code. Resultant trade-offs can occur (or be reversed if circumstances change) at  
102 several levels, and on a range of timescales. This could be considered at a behavioural level, where changes can  
103 take place immediately. It could also be considered at a cellular level with, for example, glycogen depletion  
104 resulting in a reduced capacity for performance over minutes or hours. At the organ level, deficits in, for example,  
105 bone mass or microarchitecture may accrue over months or years. Underlying all of these scenarios is the  
106 imperative to allocate available energy in the manner most likely to improve survival and reproductive success.

107 A useful analogy commonly employed in the life history literature considers the parallel between energy  
108 availability and financial budgets [18,38]. When an individual earns (or consumes) more than they spend (or  
109 expend), all expenses are paid and excess income is deposited in a savings account (*e.g.*, fat tissue). In situations  
110 where expenditure exceeds income, however (*e.g.*, if wages are cut, or expenses increase), funds must be allocated  
111 to the most immediately essential expenses, even if this comes at the cost of something else. Purchasing food, for  
112 example, may be traded off against purchasing clothing. Importantly, such trade-offs are context specific. If an  
113 individual already has adequate clothing, it makes little sense to use limited resources to buy more, particularly if  
114 this means going without food. In a cold climate with a well-stocked cupboard, however, the decision may be  
115 reversed. Alternatively, small savings can be made by marginally reducing expenditure in multiple areas, while  
116 short term deficits can be temporarily covered by withdrawing funds from one's savings account (although this  
117 strategy is not sustainable in the long-term). In a comparable way, athletes whose training load or dietary practices  
118 leave them with insufficient energy to cover the costs of all biological processes must allocate energy to certain  
119 processes at the expense of others (see Figure 1). We discuss this scenario in the following sections in the context  
120 of 4 broad categories commonly considered within the life history literature: reproduction, somatic maintenance,  
121 growth and immune defence [25–31].

122



123

124 **Figure 1.** Schematic depicting how low energy availability - whether arising via increased training expenditure,  
 125 inadequate intake, or both - requires the body to competitively allocate available energy among various biological  
 126 processes. Some processes will maintain their energy supply at the expense of others.

127

### 128 3. Reproduction

129 Life history theory posits that biological trade-offs at all stages of the life cycle ultimately serve to maximize  
 130 reproductive success, a term often used synonymously with evolutionary fitness [39–41]. Producing offspring that  
 131 go on to successfully reproduce themselves is the currency of evolutionary fitness. Energy allocation around  
 132 reproduction largely depends upon the relative pace of a species' life history, *i.e.*, its full lifespan, from conception  
 133 to death. Species with "faster" life histories generally prioritize current reproduction over somatic maintenance or  
 134 future reproduction. An example is the fruit fly, which may respond to environmental threats by increasing mating  
 135 effort to pass on its genetic material before death [42]. In contrast, larger, longer-lived mammals, who have a  
 136 "slower" life history pace typically implement a very different strategy in the face of environmental challenges  
 137 like LEA or pathogen exposure. Namely, environmental stressors that signal a short-lived fruit fly to reproduce  
 138 now may signal a larger, longer-lived mammal (*e.g.*, humans) to downregulate reproduction and wait for more  
 139 favourable conditions.

140 Reproductive downregulation is a well-established consequence of LEA in male and female athletes [9,43]. For  
 141 example, downregulation of the hypothalamic-pituitary-ovarian axis in exercising women can lead to cessation of  
 142 menses, or functional hypothalamic amenorrhea [10,44,45]. This represents a diversion of available energy away  
 143 from reproduction towards other functions and is often discussed in pathological terms. It is true that menstrual  
 144 disruption can serve to warn that energy demands of exercise training are not being met, and that prolonged  
 145 exposure to LEA may have clinical consequences (*e.g.*, for cardiovascular or bone health [46,47]). However, an  
 146 evolutionary perspective would not consider ovarian suppression to be a pathological condition *per se*. Instead,

147 interruption of the menstrual cycle can be viewed as a “*perfectly normal adaptive response to the prevailing*  
148 *environmental conditions when animals are challenged energetically*” [48]. Energy diverted away from current  
149 reproduction can be made available to other processes, while simultaneously conserving energy for future  
150 reproduction when energetic conditions improve [39,49]. This strategy makes sense in light of the high energetic  
151 costs of human female reproduction, which, in addition to pregnancy and lactation [50,51], also include caring for  
152 offspring [39,49,52]. Indeed, undertaking such an energetically expensive process in a state of LEA could  
153 jeopardize both mother and offspring health, and this appears to be a risk that humans have evolved to avoid.  
154 Functional hypothalamic amenorrhea in athletes appears to be a reversible phenomenon, with evidence suggesting  
155 affected athletes resume cycling once adequate energy is available [53–55]. A similar situation has been observed  
156 in a non-athletic context in response to seasonal fluctuations in workload-associated energy expenditure and food  
157 availability [49,56–58]. Little is currently known, however, about the longer-term reproductive consequences of  
158 transient periods of reproductive suppression in female athletes, nor how different lengths and extremes of LEA  
159 may impact subsequent reproductive capacity. Theoretically, the resumption of menses after a period of LEA-  
160 related amenorrhea could be hypothesized to impact subsequent fecundity positively, negatively, or not at all, but  
161 it appears this question is currently unaddressed in the literature.

162 Although much of the sport and exercise science literature focuses on the influence of LEA on the reproductive  
163 status of female athletes, it is important to highlight that men are also affected [9,43,59,60]. In addition to gamete  
164 production (which has minimal energetic costs), male reproduction also incurs costs related to child-care,  
165 attracting a mate and fending off male competitors, and reproductive ecologists have pointed to the modulation  
166 of physiological correlates of male mating effort (*e.g.*, testosterone, muscle mass) during times of energetic stress  
167 [39]. Evidence indicates that LEA can lead to downregulation of the hypothalamic-pituitary-testicular axis in  
168 exercising men [9,61]. Sexual dimorphism does, however, seem to exist in relation to trade-offs made under  
169 conditions of LEA [39,43,48,62,63]. This is unsurprising, given that the cost of reproduction is substantially  
170 greater for women than men [39,49,50], meaning that women in a state of LEA have a stronger biological  
171 imperative to weigh the benefits of reproductive investment against its energetic cost.

172 Perspectives differ on how indicators of energy status may predict or influence reproductive status. Reproductive  
173 ecologists postulate that *total daily energy expenditure* (sometimes referred to as “flux”) and reproductive success  
174 have a curvilinear relationship, with both very low and very high energy turnover negatively impacting  
175 reproductive success. For example, research conducted among rural Polish and Nepali women reported  
176 reproductive downregulation during more work-intensive seasons (*e.g.*, the monsoon season), despite comparable  
177 food availability [56,58]. These observations may potentially be explained by the “constrained downregulation”  
178 hypothesis, which posits that high total daily energy expenditure may render a woman less able to downregulate  
179 her metabolism, thus constraining her ability to meet the energetic demands of reproduction and lactation. Ovarian  
180 function may, therefore, be suppressed until energetic conditions change [49,58]. This is consistent with the  
181 broader constrained energy model proposed by Herman Pontzer and colleagues, which holds that human energy  
182 expenditure is constrained within physiological limits [64,65]. This model predicts that an excess of energy spent  
183 in one area (*e.g.*, training related energy expenditure) may result in a compensatory downregulation of another  
184 (*e.g.*, reproduction, or somatic maintenance) even when energy intake is adequate. Both the constrained  
185 downregulation hypothesis and the constrained energy model build upon life history theory, but interestingly, they



186 do not align well with the sport and exercise science literature. In the latter, high energy expenditure is only  
187 considered to become problematic when not met by adequate energy intake [45,66]. From this perspective, energy  
188 availability – rather than energy intake or expenditure *per se* – is considered the key regulator of reproductive  
189 function [45]. Ongoing investigation of the influence of LEA on athletic reproductive health, along with the factors  
190 that may influence this, is undoubtedly warranted.

191

#### 192 **4. Somatic Maintenance**

193 Somatic maintenance refers to “*preserving the body in good condition through diverse homeostatic processes,*  
194 *and thus promotes longevity and future opportunities for reproduction*” [29]. In conditions of LEA, certain  
195 essential maintenance processes cannot be traded off, lest survival be threatened. Other processes, however, may  
196 be temporarily downregulated without (or with fewer) negative consequences, and thus represent areas from  
197 which energy can be diverted when needed elsewhere [67]. More severe and/or prolonged periods of energy  
198 deficiency will decrease the flexibility of these energetic trade-offs, with potentially greater detriments to the  
199 body. This is particularly relevant for athletes, as undertaking, recovering from, and adapting to physical training  
200 are costly processes that may potentially result in higher somatic maintenance costs and different trade-offs than  
201 those experienced by the general population. In the following sections we consider from a life history perspective  
202 how LEA might impact the somatic maintenance of large body tissues and physical performance.

203

##### 204 *a. Bone*

205 A substantial body of research exists regarding the bone response to LEA, and this has been reviewed in detail  
206 elsewhere [68–70]. Briefly, cross-sectional and observational studies of athletes indicate that prolonged exposure  
207 to LEA manifests in lower bone mass, compromised bone micro-architecture, perturbed bone metabolism and  
208 lower estimated bone strength [71–74]. Perturbations in bone biomarkers have also been reported in individuals  
209 experimentally exposed to acute periods of LEA (*i.e.*, 3 – 5 days). This initially appears to manifest as a reduction  
210 in markers of bone formation [63,75], thus aligning with the prediction that energy will be diverted away from  
211 non-essential but energetically expensive metabolic processes during periods of LEA. Importantly, indicators of  
212 LEA are also associated with increased stress fracture risk [76–78], which has important practical and clinical  
213 implications for athlete health and performance.

214 When considered in the context of life history theory, diverting limited energy away from bone during periods of  
215 LEA could be considered a prudent strategy to preserve essential functions elsewhere, given that bone metabolism  
216 may be considered a temporarily “expendable” process [67]. Indeed, the “disposable soma” theory – an  
217 explanation of why we age that is closely related to life history theory - predicts that selection favours energetic  
218 investment in reproduction at the *expense* of longer-term bodily maintenance [79], which is likely to include the  
219 maintenance of bone tissue [80]. Bone mineral composition and micro-architecture are relatively stable in the  
220 short-term and measurable changes can take months or even years to develop, with bone-related disorders like  
221 osteoporosis primarily impacting older adults. Consequences of diverting energy away from bone metabolism

222 during periods of LEA may therefore not be felt until after the reproductive lifespan of the individual, and this  
223 could explain its apparent susceptibility to being traded off when energy resources are limited.

224 A recent review suggested that LEA-associated perturbations to bone metabolism could be offset by high-impact  
225 exercise [69]. The authors of this review posited that the benefits of high-impact exercise on bone may occur  
226 *despite* the negative influence of LEA, thus representing a potential countermeasure. A non-mutually exclusive,  
227 life history-oriented explanation is that the increased stress of high-impact loading on the skeleton may increase  
228 the necessity of skeletal preservation; potentially shifting bone higher up the hierarchy of preservation, thus  
229 rendering the tissue less likely to have resources diverted from it to other areas. Although there is a theoretical  
230 justification for a protective effect of high-impact exercise on bone during times of LEA, controlled trials are  
231 required to test the efficacy of this approach. Indeed, recent observational evidence indicates that the incidence of  
232 bone stress injury is higher in female distance runners with menstrual disturbances (assumed to be due to LEA)  
233 but that plyometric training did not influence this [81]. It is also pertinent to consider if mitigation strategies  
234 intended to protect against certain trade-offs might potentially lead to downstream trade-offs with unforeseen  
235 consequences.

236

237 *b. Muscle and Fat Mass:*

238 Muscle is the primary tissue driving human movement. It is therefore a critical determinant of athletic  
239 performance, in addition to human health [82]. Given this, one might expect muscle to maintain its energy supply  
240 during conditions of LEA. Muscle is, however, a metabolically expensive tissue accounting for approximately  
241 20% of resting energy expenditure in adult humans [83], and it incurs an even higher metabolic cost during  
242 physical activity. Substantial energy savings could, therefore, be made by reducing muscle mass when resources  
243 are scarce [9,84,85]. In addition, muscle provides an important reservoir of endogenous amino acids that can be  
244 liberated during periods of scarcity and used elsewhere, for example, to support enzyme and immune factor  
245 synthesis and maintenance of plasma protein concentrations [86]. This diversity in essential physiological  
246 functions renders muscle tissue responsive to a wide range of stimuli. The response of muscle to LEA may depend  
247 on which of its functions (*e.g.*, mechanical, regulatory or storage) proves to be most immediately essential. This  
248 in turn may depend on a host of factors, including sex, life-stage, health and training status, body composition  
249 phenotype, the length and severity of LEA, and the macro and micronutrient composition of available energy, as  
250 discussed elsewhere [87]. For example, biological sex is thought to influence the muscle response to periods of  
251 LEA, with men potentially more susceptible than women to muscle-loss during times of scarcity [88–90]. The  
252 proportion of muscle mass loss increases exponentially with decreasing body fat content [91]. Women have, on  
253 average, a larger “energy buffer” in the form of greater body fat stores than men [88,89,92]. This likely explains,  
254 at least in part, the sexually dimorphic response of muscle and fat to LEA. To the best of our knowledge, however,  
255 the relative loss of muscle and fat has not been directly tested in male or female athletes with LEA, some of whom  
256 may already be at, or close to, their lower biological adiposity limits. It is feasible that the high training status and  
257 baseline body composition of athletes may influence muscle-related trade-offs under conditions of LEA, but this  
258 requires further investigation.

259 Experimental evidence suggests that the muscle response to LEA is susceptible to manipulation by factors such  
260 as training or protein availability. Muscle mass is ultimately maintained, lost, or increased based on a dynamic  
261 balance between muscle protein synthesis and breakdown. In common with data on bone biomarkers, controlled  
262 laboratory studies indicate that short-term exposure to LEA typically manifests in a reduced rate of muscle protein  
263 synthesis [93–96]. This suggests that a portion of the energy typically allocated to building or maintaining muscle  
264 may be traded-off when resources are scarce. It appears that this can be prevented, or at least attenuated, by  
265 anabolic stimuli such as resistance exercise or amino acid availability, including the muscle protein synthesis-  
266 “triggering” amino acid leucine [93–95,97]. It has also been reported that intensive resistance training in  
267 conjunction with a high protein intake can stimulate muscle mass gains with simultaneous fat loss, despite  
268 individuals being experimentally exposed to an energy deficit of approximately 40% below estimated  
269 requirements [98]. This suggests that in certain situations the negative consequences of certain biological trade-  
270 offs may be susceptible to mitigation, given the right strategy. At the same time, as noted above regarding bone,  
271 it is important to consider whether such mitigation strategies could potentially lead to downstream trade-offs with  
272 unforeseen consequences, while the potential for the energy cost of resistance training to exacerbate the extent of  
273 LEA must also be considered.

274

275 *c. Physical performance:*

276 From an evolutionary perspective, physical activity during periods of energetic stress would have been essential  
277 to our ancestors’ survival by allowing them to hunt and gather the food required to restore adequate energy  
278 availability. Locomotion and physical activity have, however, been described as potentially reducible processes  
279 when energy availability is low [67], and substantial energy savings could be made by reducing non-essential  
280 activity-related expenditure. Convincing theoretical arguments could be made for either a positive or negative  
281 influence of LEA on exercise performance, with the actual response likely to depend upon factors including the  
282 type of activity, the training and nutritional status of the individual and the extent and duration of deficit.  
283 Intuitively, it seems likely that LEA may negatively impact exercise performance, given the importance of fuelling  
284 (both in terms of energy and individual nutrients) to optimise performance. On the other hand, and under certain  
285 circumstances, it seems plausible that performance could be maintained, or even improved, in the face of LEA.  
286 For example, a review of the influence of eating disorders on sporting performance suggested that shorter-term  
287 energy restrictions may induce transitory performance improvements [99]. This could be due to increased stress  
288 hormones like cortisol, which may have short-term stimulatory effects [100], or to increased power-mass ratios.  
289 A transitory increase in performance capacity in response to LEA could make sense from a survival perspective  
290 if it facilitates the catching or finding of needed calories. It has also been suggested that transient exposure to LEA  
291 within a periodized training and nutritional program may be desirable to drive certain phenotypic adaptations,  
292 such as body composition changes or increased efficiency of specific bioenergetic pathways [101].

293 Available intervention data directly investigating how LEA may impact exercise performance are conflicting,  
294 with both positive, neutral and negative effects shown. For example, Oliver et al. [102] reported that 48 hours of  
295 energy restriction reduced 30-minute treadmill time-trial performance [102], whereas Kimojo et al. [103] reported  
296 no influence of 3 days of LEA on time to exhaustion while running at 90% of  $VO_2$  max, despite reduced muscle

glycogen availability. In contrast, 4 weeks of energy restriction with protected protein intake led to improved countermovement jump height and 20-m sprint time in a group of male track and field jumpers and sprinters. At the same time, rapid, or gradual, weight-cutting (2.4 hours versus 3 weeks) did not impact 30m sprint or 1-minute Wingate performance or increased loaded vertical jump height [104] in a group of wrestlers and judo athletes. The type of exercise test may have been influential in these outcomes, and, plausibly, more explosive power/speed-type activities may be less susceptible to the influence of LEA than more prolonged endurance-type activities, although this hypothesis requires empirical testing. In contrast to this hypothesis, some evidence indicates that LEA may lead to an increase in exercise energy efficiency in aerobic activities [105,106]. For example, an observational study by Tornberg et al. [106] reported increased energy efficiency in a group of amenorrheic female runners, compared to their eumenorrheic counterparts, as evidenced by a lower caloric cost of pedalling on a cycle ergometer at 50 and 100 Watts.

The duration of exposure to LEA is likely to be an important determinant of its influence on exercise performance, particularly if it leads to consequences such as the suppression of reproductive hormones, muscle loss, or increased stress fracture risk. For example, observational studies have indicated that ovarian suppression as determined by states of oligo or amenorrhea may be detrimental to performance outcomes including knee muscle strength/endurance [106], 400m swim velocity [107] and training volume [108]. Athletes who increased their energy intake in response to increased training intensity had a superior performance on a running test than their counterparts who did not increase their energy intake to match the extra energy expended in training [109]. Collectively, these findings suggest that chronic LEA should be avoided to protect performance, although it is important to highlight that these are observational findings and that causality between LEA and exercise performance is as yet unconfirmed. Substantial further investigation is required to further elucidate how varying lengths and severity of LEA may impact exercise performance using a range of test protocols.

319

## 320 **5. Growth and Immune Defence**

The life history literature classically considers both growth and immune defence as important areas between which finite energy must be distributed [25–31]. Relatively little evidence is available, however, regarding the influence of LEA on growth in young athletes, or on how LEA impacts the ability to mount an immune defence in athletes of all ages. For this reason, we offer a short, combined discussion of these parameters in the hope of stimulating ideas for future research. With respect to growth, this may be threatened in young, still-growing athletes if energy availability is insufficient to meet the simultaneous demands of training, in addition to other essential maintenance functions, and growth. Research performed in rats and rabbits indicated negative impacts of nutritional deprivation on bone epiphyseal plates [110], with implications for longitudinal bone growth. Potential adverse effects of LEA on growth may be particularly relevant in sports that simultaneously emphasize early specialization and leanness and/or lightness (*e.g.*, dancers, gymnastics, and horse-racing jockeys), given that athletes may be required to maintain low body mass at pivotal developmental stages. Some evidence indicates that growth in young female gymnasts is impacted by intensive training, with catch-up growth potentially occurring when training intensity is reduced [111] (although differing perspectives on this topic do exist [112]). It seems likely that intensive training

334 will adversely affect the growth of young athletes only if the energetic demands of training are not met, thus  
335 leading to LEA, however this arguably requires further investigation according to age, sex and sport.

336 With respect to the immune system, energetic investment in maintenance and function is critical for survival,  
337 reproduction, and evolutionary fitness [26–28], but it is also energetically costly, “*particularly for those at the*  
338 *margins of nutritional adequacy*” [28]. This was recently demonstrated by Urlacher and colleagues, who found  
339 that among young Shuar forager-horticulturalists in Amazonian Ecuador, heightened immune activity alongside  
340 increased physical activity (compared to children from industrialized populations [113]) resulted in constraints on  
341 growth [114]. Interestingly, children with larger body fat levels were less susceptible to the growth-inhibiting  
342 effects of heightened immune activity, which was attributed to the greater energy reserve that higher body fat  
343 levels convey. Investigating the direct influence of LEA on immune function in athletes is challenging, given that  
344 high levels of training and competition may also influence the immune response, rendering it difficult to separate  
345 the effects of training and competition from those attributable to energy availability *per se* [65,115–117]. Because  
346 adequate energy from key fuel sources (*i.e.*, carbohydrates, amino acids and fatty acids) underlies the immune  
347 system’s ability to resist infection by various pathogens [116], it is intuitive that LEA might adversely affect  
348 immune response. Consistent with this, several studies have reported increased illness [118–120] and/or  
349 alterations to immune biomarkers [118,121] in athletes who were believed to have LEA at the time of data  
350 collection, in comparison to their energy-replete counterparts. Importantly, these observations are based on cross-  
351 sectional investigations, implying a more chronic adaptation to LEA (although of course causality cannot be  
352 confirmed from an observational study). In contrast, a recent study investigating potential trade-offs in response  
353 to the acute energetic stress of a 160km ultramarathon reported evidence for increased investment in innate  
354 immune function (indicated by increased bacterial killing ability and haemolytic complement activity) over  
355 reproductive effort (proxied by measures of testosterone and arousal) [122]. In common with many studies in this  
356 area, however, it is difficult to differentiate between LEA, versus the stress of the exercise event itself (or some  
357 other factor) as the causative agent. It is also interesting to consider that other populations with extreme LEA, *e.g.*,  
358 individuals with anorexia, are considered to be relatively robust to infection [123,124], implying that certain  
359 aspects of immunity (*e.g.*, innate versus acquired) may be more robust to trade-offs. Substantial investigation is  
360 required in this area to better understand whether and to what extent different aspects of immune function are  
361 traded-off when available energy is low.

362

## 363 **6. Perspectives for Future Research**

364 We believe that considering LEA in athletes from a life history perspective has substantial potential to inform  
365 research on this topic, both by providing an alternative viewpoint on existing data, while also allowing for  
366 discussion and hypothesis-generation to guide future research efforts. A core tenet of life history theory is that  
367 differential energy allocation will preserve – or sacrifice – different functions in different contexts, meaning that  
368 physiological processes will not be equally or simultaneously affected. A more holistic, whole-organism  
369 perspective that considers how LEA may impact specific processes *and* mediate trade-offs among other essential  
370 functions is preferable to a reductionist approach that considers functions in isolation. Additionally, a life history  
371 perspective emphasizes that LEA may differentially influence outcomes of interest (*e.g.*, bone metabolism or

372 immune function) based upon a myriad of factors, such as the life-stage of the individual, their broader nutritional  
373 status (*e.g.*, macro and micronutrient adequacy) and the type and intensity of their training schedule. Although  
374 undoubtedly challenging to adequately control and standardize for all relevant variables, researchers must remain  
375 cognizant of potential mediators and confounders and consider these in study design and interpretation.

376 Currently, limited information is available regarding the extent or duration of LEA that will provoke deleterious  
377 outcomes under different circumstances and this is an important avenue for ongoing research. Additionally, it is  
378 important to consider that LEA is extremely challenging to accurately measure [125], while varying perspectives  
379 exist related to factors such as the timeframe within which potential deficits should be considered. For example,  
380 some suggest that 3- or 7-day rolling averages are more useful, while others indicate that greater within-day  
381 imbalances may have increased consequences over similar intakes consumed more frequently throughout the day  
382 [126]. Validation of differing approaches to outcome assessment and consideration of how varying characteristics  
383 of LEA may impact health and related outcomes is an important direction for future research. This could include,  
384 for example, investigations of larger deficits across shorter time-periods versus smaller deficits across longer time  
385 periods, as well as imbalances arising primarily from higher training related expenditures versus more restricted  
386 energy intakes. It is also important to consider that a life history perspective considers the transient suppression  
387 of non-immediately essential physiological processes to be a normal, and necessary, adaptive response to  
388 conditions of energetic stress. As such, conclusions about the influence of LEA on health or performance related  
389 outcomes should only be made based on outcomes that directly relate to health or performance. Unsubstantiated  
390 assumptions that all trade-offs are negative should be avoided, as should over-extrapolation of mechanistic  
391 findings. Indeed, there may be situations where transient exposure to LEA within a periodized nutritional cycle  
392 may be desirable to drive certain phenotypic outcomes [101], while more moderate energy restriction throughout  
393 the life-course is hypothesized to increase longevity and health-span [127]. The topic of LEA in athletes is both  
394 complex and nuanced, and embracing these characteristics of the research topic may be key to the development  
395 of appropriate guidelines to support at-risk athletes.

396

## 397 **7. Conclusions and Practical Applications**

398 In summary, life history theory holds that finite energy must be competitively distributed among competing  
399 processes, and that this competition will be heightened when available energy is low. This theory has many  
400 practical applications for athletes and their support-teams. For example, life history theory considers trade-offs  
401 provoked by LEA to be context-specific, with their direction and magnitude impacted by factors including  
402 age/life-stage, sex, dietary composition, and health and training status. As such, each athlete may respond  
403 differently to a given situation, and their individual characteristics and requirements should be considered. A “one  
404 size fits all” approach is unlikely to suit the majority, and so individualization of strategies is recommended.  
405 Although there is little doubt that prolonged exposure to very LEA (*e.g.*, across months or years) may result in  
406 adverse effects on health, life history theory would not consider energetic trade-offs and their effects to always be  
407 pathological *per se*. As such, athletes and their support staff should not be overly concerned by brief periods of  
408 LEA (*e.g.*, as may occur during days or weeks of intensified training or competition) in otherwise healthy athletes,  
409 but rather should pay attention to the warning signs the body provides (*e.g.*, menstrual cycle changes) and adapt

410 training and nutritional strategies as required. Consideration of these and other factors discussed in this review  
411 may provide insight into how, and why, the body evolved to respond to periods of LEA, and ultimately serve to  
412 better inform screening, identification and treatment strategies for at-risk athletes.

413

#### 414 **Acknowledgements**

415 The authors thank Rafaela Silvério Pinto for her assistance with the development of Fig. 1.

416

#### 417 **Declarations**

418

#### 419 **Funding**

420 Eimear Dolan is financially supported by the Fundação de Ampara a Pesquisa do Estado do São Paulo  
421 (FAPESP: 2019/05616-6 and 2019/26899-6). No other sources of funding were used to assist in the preparation  
422 of this article.

423

#### 424 **Conflicts of Interest**

425 Meghan Shirley, Daniel Longman, Kirsty Elliott-Sale, Anthony Hackney, Craig Sale and Eimear Dolan declare  
426 that they have no conflicts of interest relevant to the content of this review.

427

#### 428 **Author Contributions**

429 MKS and ED conceived the original idea and developed the initial draft of this manuscript, with ongoing critical  
430 input from DPL, ACH, KES and CS. All authors read and approved the final version.

431

432

433 **REFERENCES:**

- 434 1. Loucks AB, Kiens B, Wright HH. Energy availability in athletes. *J Sports Sci.* 2011;29:37–41.
- 435 2. Loucks A. Exercise training in the normal female: Effects of low energy availability on reproductive function.  
436 In: Hackney A, Constantini N, editors. *Endocrinol Phys Act Sport.* 3rd ed. Springer Nature; 2020. p. 171–91.
- 437 3. Gibson M, Lawson D. Applying evolutionary anthropology. *Evol Anthropol.* 2015;24:3–14.
- 438 4. Stearns SC. Trade-offs in life-history evolution. *Funct Ecol.* 1989;3:259.
- 439 5. Gadgil M, Bossert WH, The S, Naturalist A, Feb NJ, Gadgil M, et al. Life historical consequences of natural  
440 selection. *Am Nat.* 1970;104:1–24.
- 441 6. Otis C, Drinkwater B, Johnson M, Loucks A, Wilmore J. American College of Sports Medicine Position  
442 Stand. The Female Athlete Triad. *Med Sci Sport Exerc.* 1997;29:1–9.
- 443 7. Mountjoy M, Sundgot-Borgen J, Burke L, Carter S, Constantini N, Lebrun C, et al. The IOC consensus  
444 statement: Beyond the Female Athlete Triad-Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med.*  
445 2014;48:491–7.
- 446 8. Areta J, Taylor H, Koehler K. Low energy availability: history, definition and evidence of its endocrine,  
447 metabolic and physiological effects in prospective studies in females and males. *Eur J Appl Physiol.*  
448 2021;121:1–21.
- 449 9. Hackney A. Hypogonadism in exercising males: Dysfunction or adaptive-regulatory adjustment? *Front*  
450 *Endocrinol (Lausanne).* 2020;11:11.
- 451 10. Nattiv A, Loucks AB, Manore MM, Sanborn CF, Sundgot-Borgen J, Warren MP. The female athlete triad.  
452 *Med Sci Sports Exerc.* 2007;39:1867–82.
- 453 11. Mountjoy M, Burke L, Ackerman KE, Blauwet C, Lebrun C, Melin A, et al. International Olympic  
454 Committee (IOC) consensus statement on relative energy deficiency in sport (RED-S): 2018 Update. *Int J Sport*  
455 *Nutr Exerc Metab.* 2018;28:316–31.
- 456 12. Reznick D, Nunney L, Tessier A. Big houses, big cars, superfleas and the costs of reproduction. *Trends Ecol*  
457 *Evol.* 2000;15:421–5.
- 458 13. Zera A, Harshman L. The physiology of life history trade- offs in animals. *Ecology.* 2001;32:95–126.
- 459 14. Bronson F. Mammalian reproduction: an ecological perspective. *Biol Reprod.* 1985;32:1–26.
- 460 15. Cody M. A General Theory of Clutch Size. *Evolution (N Y).* 1996;20:174–84.
- 461 16. Thurber C, Dugas L, Ocobock C, Carlson B, Speakman J, Pontzer H. Extreme events reveal an alimentary  
462 limit on sustained maximal human energy expenditure. *Sci Adv.* 2019;5:eaaw0341.
- 463 17. Gibbs JC, Williams NI, De Souza MJ. Prevalence of individual and combined components of the female  
464 athlete triad. *Med Sci Sports Exerc.* 2013;45:985–96.



- 465 18. Lieberman D. *Exercised: The science of physical activity, rest and health*. Penguin; 2021.
- 466 19. Lieberman DE, Bramble DM. Endurance running and the evolution of Homo. *Nature*. 2004;432:345–52.
- 467 20. Steudel-Numbers KL, Weaver TD, Wall-Scheffler CM. The evolution of human running: Effects of changes  
468 in lower-limb length on locomotor economy. *J Hum Evol*. 2007;53:191–6.
- 469 21. Pontzer H, Raichlen DA, Sockol MD. The metabolic cost of walking in humans, chimpanzees, and early  
470 hominins. *J Hum Evol*. Elsevier Ltd; 2009;56:43–54.
- 471 22. Longman DP, Wells JCK, Stock JT. Human athletic paleobiology; using sport as a model to investigate  
472 human evolutionary adaptation. *Am J Phys Anthropol*. 2020;1–18.
- 473 23. Darwin C. *On the origin of species by means of natural selection, or preservation of favoured races in the  
474 struggle for life*. London: John Murray; 1859.
- 475 24. Darwin C, Wallace A. On the tendency of species to form varieties; and on the perpetuation of varieties and  
476 species by natural means of selection. *J Proc Linn Soc London*. 1858;3:45–62.
- 477 25. Hill K. Life history theory and evolutionary anthropology. *Evol Anthropol Issues, News, Rev*. 1993;2:78–  
478 88.
- 479 26. Sheldon BC, Verhulst S. Ecological immunology: Costly parasite defences and trade-offs in evolutionary  
480 ecology. *Trends Ecol Evol*. 1996;11:317–21.
- 481 27. Lochmiller RL, Deerenberg C. Trade-offs in evolutionary immunology: Just what is the cost of immunity?  
482 *Oikos*. 2000;88:87–98.
- 483 28. McDade TW. Life history theory and the immune system: Steps toward a human ecological immunology.  
484 *Am J Phys Anthropol*. 2003;122:100–25.
- 485 29. Wells JCK. Evolutionary public health: How interventions may benefit from insights generated by life  
486 history theory. In: Schulkin J, Power M, editors. *Integr Evol Biol into Med Educ - Matern child Healthc  
487 students*, Clin Sci. Oxford: Oxford Scholarship Online; 2020.
- 488 30. Stearns S. *The evolution of life histories*. Oxford: Oxford University Press; 1992.
- 489 31. Wells JCK, Nesse RM, Sear R, Johnstone RA, Stearns SC. Evolutionary public health: introducing the  
490 concept. *Lancet*. Elsevier Ltd; 2017;390:500–9.
- 491 32. Bronson F. *Mammalian reproductive biology*. Chicago: University of Chicago Press; 1991.
- 492 33. Wells J, Stock J. The biology of the colonizing ape. *Am J Phys Anthropol*. 2007;45:191–222.
- 493 34. Flatt T, Heyland A. Mechanisms of life history evolution: The genetics and physiology of life history traits  
494 and trade-offs. Flatt T, Heyland A, editors. New York: Oxford University Press; 2011.
- 495 35. Ellison P. Endocrinology, energetics, and human life history: A synthetic model. *Horm Behav*. 2017;91:97–  
496 106.

- 497 36. West-Eberhard M. Phenotypic plasticity and the origins of diversity. *Annu Rev Ecol Syst.* 1989;20:249–78.
- 498 37. Pigliucci M, Murren C, Schlichting C. Phenotypic plasticity and evolution by genetic assimilation. *J Exp*  
499 *Biol.* 2006;209:2362–7.
- 500 38. Wells J. *The metabolic ghetto: An evolutionary perspective on nutrition, power relations, and chronic*  
501 *disease.* Cambridge: Cambridge University Press; 2016.
- 502 39. Ellison PT. Energetics and reproductive effort. *Am J Hum Biol.* 2003;15:342–51.
- 503 40. Lahdenpera M, Lummaa V, Helle S, Tremblay M, Russell A. Fitness benefits of prolonged post-  
504 reproductive lifespan in women. *Nature.* 2004;428:178–81.
- 505 41. Kaplan H, Hill K, Lancaster J, Hurtado A. A theory of human life history evolution: Diet, intelligence, and  
506 longevity. *Evol Anthropol.* 2000;156–85.
- 507 42. Polak M, Starmer WT. Parasite-induced risk of mortality elevates reproductive effort in male *Drosophila*.  
508 *Proc R Soc B Biol Sci.* 1998;265:2197–201.
- 509 43. De Souza M, Koltun K, Williams N. The role of energy availability in reproductive function in the female  
510 athlete triad and extension of its effects to men: An initial working model of a similar syndrome in male  
511 athletes. *Sport Med.* 2019;49:125–37.
- 512 44. Koltun K, De Souza M, Scheid J, Williams N. Energy availability is associated with luteinizing hormone  
513 pulse frequency and induction of luteal phase defects. *J Clin Endocrinol Metab.* 2020;105:185–93.
- 514 45. Loucks A. Energy availability, not body fatness, regulates reproductive function in women. *Exerc Sport Sci*  
515 *Rev.* 2003;31:144–8.
- 516 46. Tudor-Locke C, McColl R. Factors related to variation in premenopausal bone mineral status: a health  
517 promotion approach. *Osteoporos Int.* 2000;11:1–24.
- 518 47. O'Donnell E, Goodman J, Harvey P. Cardiovascular consequences of ovarian disruption: a focus on  
519 functional hypothalamic amenorrhea in physically active women. *J Clin Endocrinol Metab.* 2011;96:3638–48.
- 520 48. Wade G, Schneider J, Li H. Control of fertility by metabolic cues. *Am J Physiol.* 1996;1:1–19.
- 521 49. Jasienska G. *The fragile wisdom: An evolutionary view on women's biology and health.* Cambridge,  
522 Massachusetts: Harvard University Press; 2013.
- 523 50. Butte N, King J. Energy requirements during pregnancy and lactation. *Public Health Nutr.* 2005;8:1010–27.
- 524 51. Most J, Dervis S, Haman F, Adamo K, Redman L. Energy intake requirements in pregnancy. *Nutrients.*  
525 2019;11:1812.
- 526 52. Jasienska G. Costs of reproduction and ageing in the human female. *Philos Trans R Soc London Ser B Biol*  
527 *Sci.* 2020;375:21090615.
- 528 53. Areta J. Case study: Resumption of eumenorrhea in parallel with high training load after 4 years of

529 menstrual dysfunction: A 5-year follow-up of an elite female cyclist. *Int J Sport Nutr Exerc Metab.* 2020;1–6.

530 54. Cialdella-Kam L, Guebels C, Maddalozzo G, Manore M. Dietary intervention restored menses in female  
531 athletes with exercise-associated menstrual dysfunction with limited impact on bone and muscle health.  
532 *Nutrients.* 2014;6:3018–39.

533 55. Williams N, Mallinson R, De Souza M. Rationale and study design of an intervention of increased energy  
534 intake in women with exercise-associated menstrual disturbances to improve menstrual function and bone  
535 health: The REFUEL study. *Contemp Clin Trials Commun.* 2019;14:100325.

536 56. Panter-brick C, Lotstein DS, Ellison PT. Seasonality of reproductive function and weight loss in rural nepali  
537 women. *Hum Reprod.* 1993;8:684–90.

538 57. Jasienska G, Bribiescas R, Furberg A, Helle S, Nunez de la Mora A. Human reproduction and health: an  
539 evolutionary perspective. *Lancet.* 2017;390:510–20.

540 58. Jasienska G, Ellison P. Energetic factors and seasonal changes in ovarian function in women from rural  
541 Poland. *Am J Hum Biol.* 2004;16:563–80.

542 59. Tenforde A, Barrack M, Nattiv A, Fredericson M. Parallels with the female athlete triad in male athletes.  
543 *Sport Med.* 2016;46:171–82.

544 60. McGuire A, Warrington G, Doyle L. Low energy availability in male athletes: A systematic review of  
545 incidence, associations and effects. *Transl Sport Med.* 2020;3.

546 61. Lane A, Hackney A. Reproductive dysfunction from the stress of exercise training is not gender specific:  
547 The “exercise-hypogonadal male condition.” *J Endocrinol Diabetes.* 2014;1:4.

548 62. Koehler K, Hoerner N, Gibbs J, Zinner C, Braun H, De Souza M, et al. Low energy availability in exercising  
549 men is associated with reduced leptin and insulin but not with changes in other metabolic hormones. *J Sports*  
550 *Sci.* 2016;34:1–9.

551 63. Papageorgiou M, Elliott-Sale KJ, Parsons A, Tang JCY, Greeves JP, Fraser WD, et al. Effects of reduced  
552 energy availability on bone metabolism in women and men. *Bone.* 2017;105:191–9.

553 64. Pontzer H. Constrained total energy expenditure and the evolutionary biology of energy balance. *Exerc*  
554 *Sport Sci Rev.* 2015;43:110–6.

555 65. Pontzer H. Energy constraint as a novel mechanism linking exercise and health. *Physiol.* 2018;33:384–93.

556 66. Sundgot-Borgen J, Sundgot-Borgen C, Myklebust G, Solvberg N, Torstveit M. Elite athletes get pregnant,  
557 have healthy babies and return to sport early postpartum. *BMJ Open Sport Exerc Med.* 2019;21:e000652.

558 67. Wade G, Jones J. Neuroendocrinology of nutritional infertility. *Am J Physiol Regul Integr Comp Physiol.*  
559 2004;287:R1277-1296.

560 68. Papageorgiou M, Dolan E, Elliott KJ, Craig S. Reduced energy availability: Implications for bone health in  
561 physically active populations. *Eur J Nutr.* 2018;57:847–59.

- 562 69. Hutson M, O'Donnell E, Brooke-Wavell K, Sale C, Blagrove R. Effects of low energy availability on bone  
563 health in endurance athletes and high-impact exercise as a potential countermeasure: A narrative review. *Sport*  
564 *Med.* 2020;51:391–403.
- 565 70. Dolan E, Varley I, Ackerman K, Pereira R, Elliott-Sale K, Sale C. The bone metabolic response to exercise  
566 and nutrition. *Exerc Sport Sci Rev.* 2020;48:49–58.
- 567 71. Ackerman K, Nazem T, Chapko D, Russell M, Mendes N, Taylor A, et al. Bone microarchitecture is  
568 impaired in adolescent amenorrheic athletes compared with eumenorrheic athletes and nonathletic controls. *J*  
569 *Clin Endocrinol Metab.* 2011;96:3123–33.
- 570 72. Zanker CL, Swaine IL. Relation between bone turnover, oestradiol, and energy balance in women distance  
571 runners. *Br J Sports Med.* 1998;32:167–71.
- 572 73. Dolan E, McGoldrick A, Davenport C, Kelleher G, Byrne B, Tormey W, et al. An altered hormonal profile  
573 and elevated rate of bone loss are associated with low bone mass in professional horse-racing jockeys. *J Bone*  
574 *Miner Metab.* 2012;30:534–42.
- 575 74. De Souza MJ, West SL, Jamal SA, Hawker GA, Gundberg CM, Williams NI. The presence of both an  
576 energy deficiency and estrogen deficiency exacerbate alterations of bone metabolism in exercising women.  
577 *Bone.* 2008;43:140–8.
- 578 75. Ihle R, Loucks AB. Dose-response relationships between energy availability and bone turnover in young  
579 exercising women. *J Bone Miner Res.* 2004;19:1231–40.
- 580 76. Heikura I, Uusitalo A, Stellingwerff T, Bergland D, Mero A, Burke L. Low energy availability is difficult to  
581 assess but outcomes have large impact on bone injury rates in elite distance athletes. *Int J Sport Nutr Exerc*  
582 *Metab.* 2018;28:403–11.
- 583 77. Tenforde A, DeLuca S, Wu A, Ackerman K, Lewis M, Rauh M, et al. Prevalence and factors associated with  
584 bone stress injury in middle school runners. *PM&R journal Inj Funct Rehabil.* 2021;12673.
- 585 78. Barrack M, Gibbs J, De Souza M, Williams N, Nichols J, Rauh M, et al. Higher incidence of bone stress  
586 injuries with increasing female athlete triad-related risk factors: a prospective multisite study of exercising girls  
587 and women. *Am J Sports Med.* 2014;42:949–58.
- 588 79. Kirkwood T, Rose M. Evolution of senescence: late survival sacrificed for reproduction. *Philos Trans R Soc*  
589 *London Ser B Biol Sci.* 1991;332:15–24.
- 590 80. Madimenos F. An evolutionary and life-history perspective on osteoporosis. *Annu Rev Anthropol.*  
591 2015;44:189–206.
- 592 81. Hutson M, O'Donnell E, Petherick E, Brooke-Wavell K, Blagrove R. Incidence of bone stress injury is  
593 greater in competitive female distance runners with menstrual disturbances independent of participation in  
594 plyometric training. *J Sports Sci.* 2021;39:2558–66.
- 595 82. Wolfe R. The underappreciated role of muscle in health and disease. *Am J Clin Nutr.* 2006;84:475–82.

596 83. Elia M. Organ and tissue contribution to metabolic rate. In: Kinney J, Tucker H, editors. Energy Metab  
597 Tissue Determ Cell Corollaries. New York: Raven Press; 1992. p. 61–80.

598 84. Cunningham J. A reanalysis of the factors influencing basal metabolic rate in normal adults. *Am J Clin Nutr.*  
599 1980;33:2372–4.

600 85. Muchlinks M, Hemingway H, Pastor J, Omstead K, Burrows A. How the brain may have shaped muscle  
601 anatomy and physiology: A preliminary study. *Anat Rec.* 2018;301:528–37.

602 86. Carbone JW, McClung JP, Pasiakos SM. Recent advances in the characterization of skeletal muscle and  
603 whole-body protein responses to dietary protein and exercise during negative energy balance. *Adv Nutr.* Oxford  
604 University Press; 2019;10:70–9.

605 87. Oliveira-Junior G, Pinto R, Shirley M, Longman D, Koehler K, Saunders B, et al. The skeletal muscle  
606 response to energy deficiency: A life history perspective. *Adapt Hum Behav Physiol.* 2021;in press.

607 88. Henry C. The biology of human starvation: Some new insights. *Nutr Bull.* 2008;26:205–11.

608 89. Widdowson E. The response of the sexes to nutritional stress. *Proc Nutr Soc.* 1976;35:175–80.

609 90. PrayGod G, Range N, Faurholt-Jepsen D, Jeremiah K, Faurholt-Jepsen M, Aabye M, et al. Predictors of  
610 body composition changes during tuberculosis treatment in Mwanza, Tanzania. *Eur J Nutr.* 2015;69:1125–32.

611 91. Forbes G. Body fat content influences the body composition response to nutrition and exercise. *Ann N Y*  
612 *Acad Sci.* 2000;904:359–65.

613 92. Ocobock C. Body fat attenuates muscle mass catabolism among physically active humans in temperate and  
614 cold high altitude environments. *Am J Hum Biol.* 2017;29:23013.

615

616 93. Areta JL, Burke LM, Camera DM, West DWD, Crawshay S, Moore DR, et al. Reduced resting skeletal  
617 muscle protein synthesis is rescued by resistance exercise and protein ingestion following short-term energy  
618 deficit. *Am J Physiol - Endocrinol Metab.* 2014;306:989–97.

619 94. Hector AJ, McGlory C, Damas F, Mazara N, Baker SK, Phillips SM. Pronounced energy restriction with  
620 elevated protein intake results in no change in proteolysis and reductions in skeletal muscle protein synthesis  
621 that are mitigated by resistance exercise. *FASEB J.* 2018;32:265–75.

622 95. Murphy C, Churchward-Venne T, Mitchell C, Kolar N, Kassis A, Karagounis L, et al. Hypoenergetic  
623 diet-induced reductions in myofibrillar protein synthesis are restored with resistance training and balanced daily  
624 protein ingestion in older men. *Am J Physiol Endocrinol Metab.* 2015;308:734–43.

625 96. Pasiakos S, Vislocky L, Carbone J, Altieri N, Konopelski K, Freake H, et al. Acute energy deprivation  
626 affects skeletal muscle protein synthesis and associated intracellular signaling proteins in physically active  
627 adults. *J Nutr.* 2010;140:745–51.

628 97. Wilkinson D, Hossain T, Hill D, Phillips B, Crossland H, Williams J, et al. Effects of leucine and its  
629 metabolite  $\beta$ -hydroxy- $\beta$ -methylbutyrate on human skeletal muscle protein metabolism. *J Physiol.*

630 2013;591:2911–23.

631 98. Longland T, Oikawa S, Mitchell C, Devries M, Phillips S. Higher compared with lower dietary protein  
632 during an energy deficit combined with intense exercise promotes greater lean mass gain and fat mass loss: a  
633 randomized trial. *Am J Clin Nutr.* 2016;103:738–46.

634 99. Ghoch M, Soave F, Calugi S, Dalle Grave R. Eating disorders, physical fitness and sport performance: A  
635 systematic review. *Nutrients.* 2013;5:140–60.

636 100. Hackney A, Walz E. Hormonal adaptation and the stress of exercise training: the role of glucocorticoids.  
637 *Trends Sport Sci.* 2013;4:165–71.

638 101. Stellingwerff T, Morton J, Burke L. A framework for periodized nutrition for athletes. *Int J Sport Nutr*  
639 *Exerc Metab.* 2019;29:141–51.

640 102. Oliver S, Laing S, Wilson S, Bilzon J, Walsh N. Endurance running performance after 48 hours of  
641 restricted fluid and/or energy intake. *Med Sci Sport Exerc.* 2007;39:316–22.

642 103. Kojima C, Ashibashi A, Tanabe Y, Iwayama K, Kamei A, Takahashi H, et al. Muscle glycogen content  
643 during endurance training under low energy availability. *Med Sci Sport Exerc.* 2020;52:187–95.

644 104. Fogelholm G, Koskinen R, Laakso J, Rankinen T, Ruokonen I. Gradual and rapid weight loss: Effects on  
645 nutrition and performance in male athletes. *Med Sci Sport Exerc.* 1993;25:371–7.

646 105. Pons V, Riera J, Capo X, Martorell M, Sureda A, Tur J, et al. Calorie restriction regime enhances physical  
647 performance of trained athlete. *J Int Soc Sports Nutr.* 2018;15:12.

648 106. Tornberg A, Melin A, Manderson Koivula F, Johansson A, Skouby S, Faber J, et al. Reduced  
649 neuromuscular performance in amenorrheic elite endurance athletes. *Med Sci Sport Exerc.* 2017;49:2478–85.

650 107. Vanheest J, Rodgers C, Majoney C, De Souza M. Ovarian suppression impairs sport performance in junior  
651 elite female swimmers. *Med Sci Sport Exerc.* 2014;46:156–66.

652 108. Ihalainen J, Kettunen O, McGawley K, Solli G, Hackney A, Mero A, et al. Body composition, energy  
653 availability, training, and menstrual status in female runners. *Int J Sports Physiol Perform.* 2021;16:1043–8.

654 109. Schaal K, VanLoan M, Hausswirth C, Casazza G. Decreased energy availability during training overload is  
655 associated with non-functional overreaching and suppressed ovarian function in female runners. *Appl Physiol*  
656 *Nutr Metab.* 2021;46:1179–88.

657 110. da Silva K, Silva C, Costa R, De Moraes S. How does protein malnutrition or food deprivation interfere  
658 with the growth of the epiphyseal plate in animals? *Int J Morphol.* 2013;31:584–9.

659 111. Caine D, Lewis R, O'Connor P, Howe W, Bass S. Does gymnastics training inhibit growth of females. *Clin*  
660 *J Sport Med.* 2001;11:260–70.

661 112. Baxter-Jones A, Maffuli N, Mirwald R. Does elite competition inhibit growth and delay maturation in  
662 some gymnasts? Probably not. *Pediatr Exerc Sci.* 2003;15:373–82.

663 113. Urlacher S, Snodgrass J, Dugas L, Sugiyama L, Liebert M, Joyce C, et al. Constraint and trade-offs  
664 regulate energy expenditure during childhood. *Sci Adv.* 2019;5:eaax1065.

665 114. Urlacher S, Ellison P, Sugiyama L, Pontzer H, Eick G, Liebert M, et al. Tradeoffs between immune  
666 function and childhood growth among Amazonian forager-horticulturalists. *Proc Natl Acad Sci U S A.*  
667 2018;115:E3914–24.

668 115. Campbell JP, Turner JE. Debunking the myth of exercise-induced immune suppression: Redefining the  
669 impact of exercise on immunological health across the lifespan. *Front Immunol.* 2018;9.

670 116. Walsh NP. Nutrition and athlete immune health: New perspectives on an old paradigm. *Sport Med.*  
671 Springer International Publishing; 2019;49:153–68.

672 117. Simpson R, Campbell J, Gleeson M, Kruger L, Nieman D, Pyne D, et al. Can exercise affect immune  
673 function to increase susceptibility to infection? *Exerc Immunol Rev.* 2020;26:8–22.

674 118. Shimizu K, Suzuki N, Nakamura M, Aizawa K, Imai T, Suzuke S, et al. Mucosal immune function  
675 comparison between amenorrheic and eumenorrheic distance runners. *J Strength Cond Res.* 2012;26:1402–6.

676 119. Drew MK, Vlahovich N, Hughes D, Appaneal R, Peterson K, Burke L, et al. A multifactorial evaluation of  
677 illness risk factors in athletes preparing for the Summer Olympic Games. *J Sci Med Sport. Sports Medicine*  
678 *Australia;* 2017;20:745–50.

679 120. Hagmar M, Hirschberg AL, Berglund L, Berglund B. Special attention to the weight-control strategies  
680 employed by olympic athletes striving for leanness is required. *Clin J Sport Med.* 2008;18:5–9.

681 121. Sarin H V., Gudelj I, Honkanen J, Ihalainen JK, Vuorela A, Lee JH, et al. Molecular pathways mediating  
682 immunosuppression in response to prolonged intensive physical training, low-energy availability, and intensive  
683 weight loss. *Front Immunol.* 2019;10.

684 122. Longman DP, Prall SP, Shattuck EC, Stephen ID, Stock JT, Wells JCK, et al. Short-term resource  
685 allocation during extensive athletic competition. *Am J Hum Biol.* 2018;30:1–11.

686 123. Slotwinska S, Slowwinski R. Immune disorders in anorexia. *Cent Eur J Immunol.* 2017;42:294–300.

687 124. Marcos A. Eating disorder: A situation of malnutrition with peculiar changes in the immune system. *Eur J*  
688 *Clin Nutr.* 2000;54:61–4.

689 125. Burke L, Lundy B, Fahrenholtz I, Melin A. Pitfalls of conducting and interpreting estimates of energy  
690 availability in free-living athletes. *Int J Sport Nutr Exerc Metab.* 2018;28:350–63.

691 126. Fahrenholtz I, Sjodin A, Benardot D, Tornberg A, Skouby S, Faber J, et al. Within-day energy deficiency  
692 and reproductive function in female endurance athletes. *Scand J Med Sci Sport.* 18AD;28:1139–46.

693 127. Speakman J. Why does caloric restriction increase life and healthspan? The “clean cupboards” hypothesis.  
694 *Natl Sci Rev.* 2020;7:1153–6.

695