

1 **The influence of boot design on exercise associated surface temperature of** 2 **tendons in horses**

3 *L. Hopegood¹, L. Sander², A. D. Ellis³*

4 ¹ *Nottingham Trent University, School of Animal, Rural and Environmental Sciences,*
5 *Brackenhurst, Southwell, Nottinghamshire, NG25 0QF, UK,*

6 ² *Hofgut Beutig 1, 04749, Ostrau, Germany.*

7 ³ *UNEQUI, Research, Education, Innovation; Southwell, NG25 0DS, UK.*

8 *Corresponding author: lyn.hopegood@ntu.ac.uk*
9

10

11 **Abstract**

12 Sport horses frequently injure tendons of the lower limb. Tendon boots are commonly applied
13 for structural support and trauma prevention during competitions. However these boots may
14 increase heat stress in the area. Two separate studies were carried out with the aim to
15 improve understanding of the effect of boots on heat around the tendon area. Study 1
16 measured heat emitted from two types of boots (traditional and perforated, cross over design)
17 covering the superficial digital flexor tendon (SDFT) in 4 horses during a set ridden and
18 lunged exercise test. Study 2, a Field test, measured the effect of boot style (traditional,
19 perforated and open fronted) on skin surface temperature in 131 horses, after completing a
20 cross country event test (either a BE 100 three day event or a CCI* - two day short format
21 event). The Raytek Raynger ST20 (infrared thermometer) was used to measure temperatures
22 during both studies. The MobIR[®]M4 Thermal Imager was also used in Study 1 to compare
23 measurement methods. A significant correlation was found between both measurement types
24 ($p < 0.001$; $R^2 = 0.94$). Boots designed with perforations demonstrated greater heat emissions
25 than traditional (non-perforated) boots (+ 3.5° C, $p < 0.01$). In Study 2 mean tendon surface
26 temperature for perforated type boots (28.0°C) was significantly lower than for traditional
27 boots (32.3°C) and for open fronted tendon boots (31.1°C) ($P < 0.001$). As this was an applied

28 field study, additional environmental factors, such as speed and fitness level of horses, may
29 have influenced results.

30

31 *Take home message*

32 Although exact mechanisms leading to these findings and the link between heat and tendon
33 injury need to be researched further, it is advisable to design boots to minimise tendon
34 exposure to high temperatures, which may contribute to tendon injury.

35 Keywords: heat, equine, protective equipment, thermal imaging

36

37 **Introduction**

38 Injuries of the superficial digital flexor tendon are one of the most common causes of
39 lameness in the Thoroughbred race horse (Patterson-Kane and Firth 2009) and in athletic
40 sports horses (Murray *et al.* 2006). In National Hunt racing (over jumps) 89% of all ligament
41 and tendon injuries are to the SDFT (Ely *et al.* 2009). Singer *et al.* (2008) found 24% of all
42 injuries (0.45% of starts recorded at events) during the cross country phase in eventing are
43 related to tendons and ligaments. In addition 43 % of injuries sustained during training for
44 eventing are tendon or ligament injuries and 36% of these involved the SDFT (Singer *et al.*
45 2008). The human Achilles tendon (AT) is considered functionally equivalent to the equine
46 SDFT and in humans the AT has a pivotal role in saving energy during high-speed
47 locomotion, by reducing muscular work, which leads to increased heat in the area (Malvankar
48 and Khan, 2011).

49

50 Wilson and Goodship (1994) measured heat produced inside tendons *in vivo* and found
51 temperatures of 45°C in the SDFT and concluded that this heat could be a major contributor
52 to degenerative changes in tendons of equine and human athletes. At temperatures of 45°C -
53 48°C over a period of 10 minutes a rapid decline of tendon fibroblast activity takes place,

54 resulting in cell death (Birch *et al.* 1997; Burrows *et al.* 2008). Yamasaki *et al.* (2001) also
55 showed *in vitro* that when tendons were exposed to 45°C for 10 minutes only 27% of
56 tenocytes survived and showed *in vivo* that Temperatures of 45°C were reached after a short
57 gallop on the track. Although tenocytes have a higher heat resistance than other cells these
58 temperatures at shorter time periods are likely to influence tendon matrix quality leading to
59 some damage of tenocytes (Smith 2004; Patterson-Kane and Firth 2009). The extrapolation of
60 the *in vitro and in vivo* data on core temperatures to *in vivo* tendon injuries should be made
61 with caution, as this has not yet been directly linked.

62
63 In order to prevent mechanical injury to the lower limb from over reach, hitting jumps or
64 during a fall, various types of boots are worn routinely during jumping competitions (Murphy,
65 2008). The encasement of the lower limb by the boot may increase heat stress to soft tissue
66 but to date *in vivo* research in this area is scarce. The design and structure of horse boots
67 varies according to their purpose. Traditional boots, such as brushing boots enclosing the
68 limb distal to the carpus and proximal to the metacarpophalangeal joint used to be made of
69 leather with leather straps. Open-fronted tendon boots were generally worn by show jumpers,
70 providing extra padding around the tendons but being open dorsally to ‘remind horses to pick
71 up their feet’ over jumps (Murphy, 2008). Modern boots are often made of a mix of more
72 pliable and softer cushioning materials which are cheaper, easier to clean and maintain.
73 Polyvinyl Chloride (PVC) is a water-proof rigid thermoplastic polymer with added
74 plasticizers for flexibility. It has a thermal conductivity of 0.14-0.17 W/m-K (the lower the
75 number the more insulation it provides). Polycarbonate, another thermoplastic polymer, with
76 a thermal conductivity of 0.19-0.22 W/m-K is used because of its light weight, impact
77 strength (Izod 600-850 J/m) and temperature resistance (Rouabah *et al.* 2007). Most modern
78 horse boots combine a softer inner layer such as neoprene (polychloroprene, thermal
79 conductivity of 0.054 W/m-K - high insulation) and a synthetic rubber which also provides

80 insulation (e.g used in wetsuits), and aids prevention of rubbing (Bardy *et al.* 2006).
81 Thermoplastic elastomers (generally a mixture of plastic and rubber) are used for their
82 softness and durability (Holden *et al.* 2000) imparting high elastic properties and these have a
83 thermal conductivity of 0.209-0.251 W/m-K. Combining these materials in a boot will
84 prevent endogenous heat dispersal. Heat dissipation from the surface of the limb without a
85 boot occurs through evaporation and radiation. With a boot further conduction is required
86 through the boot material followed by radiation from the boot surface.

87
88 In recent years a focus on prevention of injury and a deeper understanding of tendon injuries
89 has led to the development of boots with perforation holes (often called ‘air-cooled’ boots),
90 which are marketed as ‘allowing better dissipation of heat through air circulation to the limb’.
91 However, no research, to the knowledge of these authors, has been published which tests this
92 theory.

93

94 *Aims*

95 The overall aim of the studies presented here was to measure the effect of fully closed versus
96 perforated boots on SDFT skin surface temperature in exercising horses.

97

98 The aim of Study 1 was to measure heat emitted from either traditional closed boots or novel
99 air-perforated boots after two controlled exercise tests in 4 horses.

100

101 The aim of Study 2 was to measure the effect of boot style used for 131 horses on tendon skin
102 surface temperature following a cross country Field test.

103

104 **Materials and Methods**

105

106 *Experimental Design*

107 The study passed the procedures of the Ethical Review Committee at Nottingham Trent
108 University. Study 1 employed a controlled cross over design with 4 horses. The Treatment
109 consisted of traditional designed tendon boots or a perforated (air-flow) boot. Study 2 was a
110 field study and measured SDFT skin surface temperature of 131 horses immediately after
111 completing a high intensity exercise (Cross Country Event).

112

113 *Thermometer*

114 The Raytek Raynger ST20 Laser Thermometer (Berlin, Germany) (non-contact infrared
115 thermometer with a temperature range of -32°C to 535°C) was used to measure temperatures.
116 The Raytek Raynger ST20 is usable in all weather conditions and was used in both studies for
117 speed and ease of measurement. Temperatures were taken approximately 5 cm from the limb
118 according to manufacturers' guidelines (Raytek Raynger).

119

120 *Study 1*

121 Four sound German Warmbloods (550 ± 50 kg Bodyweight, 6-13 years old) were selected.
122 The study consisted of two parts, a lunging test and a ridden exercise test which occurred over
123 four days with two different phases (cross-over design). The Eskadron (Werther, Germany)
124 more 'traditional' cross country boot and the New Equine Wear 'perforated' boots
125 (Chippenham, UK) were tested on all four limbs. Both boots were chosen as they were made
126 similarly with a PVC upper surface and Neoprene cushioning material underneath. Each test
127 was performed twice and horses wore one set of boots (e.g. traditional) on left limbs while
128 wearing the other set (e.g. perforated) on the opposite leg in a further cross-over design.
129 Therefore, each leg was used as a separate unit with the opposite leg as control treatment so
130 that measures were taken simultaneously reducing any environmental effects over time.
131 Measurements were always performed in the same order to eliminate timing results (i.e.

132 temperature was taken from left fore and hind legs first, then from right fore and hind
133 resulting in a further cross-over between boot types). The set exercise tests (same rider or
134 handler) were carried out in an arena with a sand surface and included 10 minutes warm up in
135 walk followed by 5 minutes of trot in each direction, and a canter for a) Ridden test: 2.5
136 minutes on each rein and b) Lunging test: 1 minute on each rein.

137

138 After the exercise, temperatures were taken in three areas to assess heat emissions with the
139 boot still in place at the lateral aspect of the boot (area of SDFT): just 1 cm below the top
140 edge of the boot (top), the mid-point (middle – midway between the carpus and the
141 metacarpal-phalangeal joint) and 1 cm above the bottom edge of the boot (bottom) on all four
142 legs with both methods of temperature measurement.

143

144

145 *Study 2*

146 The field test was carried out at the International Horse Trials at Aldon. Horses from two
147 classes were used: A British Eventing (BE100) – three day event: length: 3000 m;
148 height/No.of jumps: 1.5 m/25, n=61; and a Fédération Equestrienne Internationale (FEI)
149 Concours Internationale Combined – One Star (CIC)* - two day short format event – length:
150 3000 m; height/No.of jumps: 1.3 m/24, n=69). Horses in the three day BE 100 event also
151 have completed the field and track phases (incl. 4000 m Endurance) on the same day prior to
152 the cross country event. Twenty-one types of boots could be distinguished according to style
153 and manufacturer. Results for boots were pooled into three groups according to style: a
154 traditional boot design (closed all around the leg = traditional, n=93), boots with a design
155 using holes, perforations or mesh design to allow for air to cool the leg (perforated, n=24),
156 open fronted tendon boots (tendon boots, n=12) and no boots (n=2).

157

158 Temperature of the left front leg only was taken, mid-way between the carpus and the
159 metacarpo-phalangeal joint on the SDFT, immediately after horses had crossed the finish line
160 after the boot had been removed (one leg only was used purely because of speed and the
161 horses needing to be cooled down). A few horse owners insisted on removal of boots
162 immediately after the finish line of the Field test and therefore temperature had to be
163 measured in all horses after removal of the boot.

164

165

166 *Statistical Analysis*


167 Statistical analyses was carried out using IBM SPSS (v17.00). The significance level was set
168 as $P < 0.05$. Data were analysed for normal distribution (Kolmogorov-Smirnov) and Analysis
169 of Variance (ANOVA) was applied, testing for differences between boots and for effects and
170 interactions between left and right, fore and hind, exercise and phases (Study 1). For study 2
171 ANOVA was used to test for effect of boots and cross country class. Unless stated otherwise,
172 means are reported with standard errors.

173

174 **Results**

175 *Study 1*

176 There were no differences between left and right legs, fore and hind legs or according to
177 phases and no interaction was identified. Therefore independent leg data were pooled and
178 when all temperature areas (top, middle and bottom) were considered together there was no
179 significant difference in heat emissions between the two boots but there was a significant
180 difference between areas irrelevant of boots ($P < 0.001$; $n=8$; Figure 1).



181

182 Figure 1. Distribution of mean heat emissions of outer surface of boots for horses undergoing
 183 two exercise tests depending on measurement area (ab – significantly different $P=0.000$,
 184 $F=18$; ANOVA, $n=8$)

185

186 There was a significant difference in heat emissions from the middle area of boots between
 187 exercises and between boots ($P<0.01$) (Table 1).

188

189 Table 1. Mean heat emissions from the outer surface in the middle of the boot according to
 190 boots and exercise for 4 horses

191


	Treatment	Temperature (°C)	s.d.	P-value ¹	n
Traditional	Lunging	11.63	19	$P=0.064$, $F=5.1$	4
	Ridden	8.54	19		
Perforated	Lunging	15.46	2.0	$P=0.074$, $F=4.7$	4
	Ridden	12.00	2.9		

Boots	Traditional	10.08	2.4	P=0.005, F=11	8 ²
	Perforated	13.73	3.0		
Exercise	Lunging	13.54	2.7	P=0.009, F=9.6	8 ³
	Ridden	10.27	3.0		

192 ¹Univariate Anova ²x 2 Exercises ³x 2 Boots
193

194 Study 2

195 There was no significant difference in limb temperatures between the two eventing classes
196 although the perforated boots showed a much narrower distribution of measures for the CIC*
197 Event (Figure 2).



198

199 Figure 2. Distribution of temperatures under the tendon boots at the finish line of two Cross
200 Country Events (BE 100 – 3 day event with roads and tracks, n=61; CCI* = 2 day show
201 jumping and cross country short format event, n=69)

202

203 There was a significantly lower temperature under perforated boots compared to open-fronted
204 tendon boots ($p < 0.001$) and compared to traditional boots ($p < 0.001$, Table 2). The difference

205 between open-fronted and traditional boots was slight but significant ($p<0.05$) and much
 206 lower temperatures were recorded for the two horses, who did not wear boots.

207

208 Table 2. Mean, minimum and maximum temperatures of skin surface according to type of
 209 boot for 129 horses upon completion of the cross country phase of a one day event

210

Boot Type	Mean ¹	s.e.	Min	Max
Traditional	32.33 ^a	±0.17	29.3	36.5
Perforated	28.66 ^b	±0.32	25.6	32.6
Open-fronted	31.10 ^c	±0.47	28.9	33.4
No Boot ²	21.70		21.2	22.2

211 ¹Anova, Bonferroni: Superscripts ab, bc $P<0.001$; ac $P<0.05$

212 ² n=2, not included in statistical analysis

213

214

215 Discussion

216 The aims of this study were to measure the effect of boot style on heat emissions from the
 217 surface of boots following light exercise and on skin surface temperature under the boots
 218 following heavy exercise. The RayTek Laser Infra-red thermometer was ideal for temperature
 219 measurements in the competition environment, as it could be used quickly and easily. The
 220 slightly unconventional application of different boots on opposing legs was used in Study 1 to
 221 help eliminate environmental influences.

222

223 As the thermal image showed a large change in temperatures between the centre of the boots
 224 and the top and bottom, it was decided to take mean measurements from these three points for
 225 both measurement devices. In Study 1 there were significantly lower heat emissions from the
 226 middle of both boots ($p<0.01$) compared to the top and bottom showing a stronger insulation
 227 effect of boots in this area which may be due to a greater insulating effect or it could be due to
 228 less heat in the leg underneath those areas relative to the more proximal and distal areas.
 229 Although this area of the of the SDFT has been found at higher risk of developing lesions,

230 Birch *et al.* (2002) did not find a change in strength or composition of tendons in this area.
231 These authors actually pointed towards the possibility of hypoxia and/or hyperthermia in this
232 area causing site-specific tendon lesions in the SDFT (Birch *et al.*, 2002). Further research on
233 the SDFT in various areas with and without a boot needs to be carried out to establish a
234 possible link.

235

236 The boots chosen in Study 1 were made of the same material and thickness, with the main
237 difference being the perforations for the air cooled boot. Results for the perforated boots may
238 indicate that greater heat emission and thus heat dissipation took place by a mean of $+3.5^{\circ}\text{C}$
239 (± 1.1 s.e.) from traditional boots. This interpretation is further supported by a reduced
240 temperature ($-3.7^{\circ}\text{C} \pm 0.13$ s.e.) measured in the same metacarpal area on removal of the
241 perforated boots compared to the traditional boots in Study 2. These two studies have to be
242 treated separately due to differences in exercise level and design as well as the different range
243 of boots used by competitors in Study 2. In favour of some comparison is the high number of
244 participants of the field study and that the results seem to show some convergence. It is
245 unfortunate that, due to circumstances beyond our control, it was not possible to take similar
246 measurements in both studies.

247

248 The difference in the temperature lost from the middle of the boots during lunging was higher
249 than when horses were ridden ($+3.1^{\circ}\text{C} \pm 1.3$ s.e.), although exercise duration was 3 minutes
250 shorter when horses were lunged. The balanced design of Study 1, with repetition in each
251 horse, as well as no phase effects point towards this being a true effect. Differences may be
252 due to changes in biomechanical use of the limb due to working on a tighter circle or due to
253 speed and collection via rider/handler influence.

254

255 To date there is a lack of data available in measuring differences in leg temperatures between
256 horses according to their heart rates and speed and other factors in competitions. Study 2 can
257 be summed up as a pilot field study to record what types of boots are used and to measure *in*
258 *vivo* temperatures after 130 horses completed a cross country event. The range of
259 temperatures measured was between 29-37°C (traditional boots) and 26-33°C (perforated
260 boots). Differences within and between groups could be due to boot design, speed and fitness
261 level. To eliminate or measure these factors would be desirable in future research. The
262 insulation effect of boots overall was shown by the limb temperatures of the two horses
263 competing without boots (21-23°C) which were considerably lower than those wearing any
264 type of boots (mean 30.7°C). In an ideal situation a higher sample population with no boots
265 could confirm these results. The open-fronted tendon boots did not result in a relevant
266 reduction in temperature (-1.2°C) compared to traditional fully enclosed boots.

267
268 Although long term training and mechanical stress during exercise over time may play a
269 primary role in injury (Ely *et al.* 2009), additional heat stress may contribute to final aetiology
270 (Wilson and Goodship, 1994; Birch *et al.*, 2002). Birch *et al.* (1997) reported that a tendon
271 temperature of 39°C already leads to a 4% decrease of tendon fibroblast viability. Wilson and
272 Goodship (1994) reported that the skin temperature was around 5.4°C cooler than the tendon
273 core temperature in their *in vivo* study at an ambient temperature of 2°C, when horses
274 galloped for 2 minutes on a treadmill. The maximum skin surface temperature in Study 2 was
275 36.5°C for traditional boots, however, the ambient temperature during the horse trial was
276 much warmer at 9°C (Met Office; Weather station, Yeovilton). In addition the thermometer
277 was only commercially calibrated and had not been re-calibrated against absolute temperature
278 measurements prior to commencement of the trial. This limits direct temperature comparison
279 between studies. However, following these results, the possible link between heat stress and
280 boot design warrants further exploration.

281
282 Future research could focus on measuring temperatures during the actual exercise with a
283 motion thermal imaging camera or electrode surface temperature time-lapse recording
284 equipment while looking at speed and heart rate of horses, together with final performance.
285 This would allow for evaluation of temperature development from onset of exercise and could
286 also look at the effect of cooling down periods and the influence of water fences on the limb
287 temperature. Furthermore, temperatures developed under the boots or bandages of national
288 hunt and racehorses should be evaluated. In addition effect of boot material on heat insulation
289 needs to be investigated further.

290

291 **Conclusion**

292 During both studies differences in heat retention according to the type of boot were recorded.
293 This points towards a possible cooling effect of air perforated boots but further more detailed
294 research is required to test other influencing factors and to test the protective properties of
295 these boots. Although exact mechanisms leading to these findings and the link between heat
296 and tendon injury needs to be proven *in vivo*, based on current knowledge, it is advisable to
297 design boots to minimise tendon exposure to high temperatures.

298

299

300 **Acknowledgements**

301 Thanks go to the whole team at Aldon Horse Trials for accommodating the study. Further
302 thanks go to all the riders that participated.

303

304 **Conflict of Interest Statement**

305 None of the authors of this paper has a financial relationship with any organisation that could
306 inappropriately influence or bias the content of this paper.

307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356

References

- Bardy, E., Mollendorf, J. and Pendergast, D. 2006. A comparison of the thermal resistance of a foam neoprene wetsuit to a wetsuit fabricated from aerogel-syntactic foam hybrid insulation Journal of Physics D: Applied Physics 39: 18, 4068-4076 (9).
- Birch, H.L., Wilson, A.M. and Goodship, A.E. 1997. The effect of exercise induced hyperthermia on tendon cell survival. The Journal of Experimental Biology. 200. 1703-1708.
- Birch H.L., Smith T.J., Poulton C., Peiffer D., Goodship A.E. 2002. Do regional variations in flexor tendons predispose to site-specific injuries? Equine Veterinary Journal. Supplement 34, 288-292.
- Burrows, S., Patterson-Kane, J.C., Fleck, R.A., Becker, D.L. 2008. Alterations in gap junction communication in tenocyte monolayers following an episode of hyperthermia. Transactions of the Orthopaedic Research Society. 33: 0323.
- Dowling, B.A., Dart, A.J., Hodgson, D.R., Rose, R.J. and Walsh, W.R. 2002. Recombinant equine growth hormone does not affect the *in vitro* biomechanical properties of equine superficial digital flexor tendons. Veterinary Surgery. 31, 325-330.
- Eddy, A.L., van Hoogmoed, L.M. and Snyder, J.R., 2001. The role of thermography in the management of equine lameness. The Veterinary Journal. 162. 172-181.
- Ely, E.R., Avella, C.S., Price, J.S., Smith, R.K.W., Wood, J.L.N. and Verheyen, K.L.P. 2009. Descriptive epidemiology of fracture, tendon and suspensory ligament injuries in National Hunt racehorses in training. Equine Veterinary Journal. 41:4, 372-378.
- Holden, G., Kricheldorf, H. R. and Quirk, R. P. 2000. Thermoplastic Elastomers, Hanser 3rd edition.
- Liduin, S., Meershoek, L.S. and Lanovaz, J.L. 2001. Sensitivity analysis and application to trotting of a non-invasive method to calculate flexor tendon forces in the equine forelimb. American Journal of Veterinary Research. 62. 1594-1598.
- Malvankar, S. and Khan, W.S. 2011. Evolution of the Achilles tendon: the athlete's Achilles heel? Foot (Edinburgh). 21. 193-197.
- Murphy, J. 2008 Boots on Horses: Limb Protection or Hyperflexion Training Aids in the Showjumping Horse. Journal of Applied Animal Welfare Science. 11:3, 223-227.
- Murray, R. C., Dyson, S. J., Tranquille, C. and Adams, V. 2006. Association of type of sport and performance level with anatomical site of orthopaedic injury diagnosis, Equine Veterinary Journal, 38, 411-416.
- Patterson-Kane, J.C. and Firth, E.C. 2009. The pathobiology of exercise-induced superficial digital flexor tendon injury in Thoroughbred racehorses, The Veterinary Journal. 181:2, 79-89.

- 357 Purohit, R.C. and McCoy, M.D. 1980. Thermography in the diagnosis of inflammatory
358 processes in the horse. *American Journal of Veterinary Research*. 41, 1167-1174.
359
- 360 Rouabah F., Fois M., Ibos L., Boudenne A., Dadache D., Haddaoui N. and Ausset P. 2007.
361 Mechanical and thermal properties of polycarbonate. II. Influence of titanium dioxide content
362 and quenching on pigmented polycarbonate, *Journal of Applied Polymer Science*. 106, 2710 –
363 2717.
364
- 365 Singer, E.R., Barnes, J., Saxby, F. and Murray, J.K. 2008. Injuries in the event horse: training
366 versus competition. *Veterinary Journal*. 175, 76-81.
367
- 368 Smith R.K.W. 2004. Equine tendon adaptation to training: which type of exercise does what?
369 Conference of Equine Sports Medicine and Science, CESMAS 2004, Oslo.
370
- 371 Verna, M., Turner, T.A. and Anderson, K.L. 2005. Scintigraphic, radiographic, and
372 thermographic appearance of the metacarpal and metatarsal regions of adult healthy horses
373 treated with nonfocused extracorporeal shock wave therapy – a pilot study. *Veterinary*
374 *Therapeutics*. 6:3, 268-276.
375
- 376 Wilson, A.M. and Goodship, A.E. 1994. Exercise-induced hyperthermia as a possible
377 mechanism for tendon degeneration, *Journal of Biomechanics*, 27, 899-905.
378
- 379 Yamasaki, H., Goto, M., Yoshihara, T., Sekigushi, M., Konno, K., Momoi, Y. and Iwasaki, T.
380 2001. Exercise- induced Superficial Digital Flexor Tendon Hyperthermia and the effect of
381 cooling sheets on thoroughbreds, *Journal of Equine Science*. 12. 85-91.
382
- 383 Yang, W. and Yang, P.P.T. 1992 Literature survey on biomedical applications of
384 thermography. *Bio-Medical Materials and Engineering*. 2. 7-18.