

Psychological morbidity and return to work after injury: multicentre cohort study

Kendrick D,¹ Dhiman P,² Kellezi B,³ Coupland C,¹ Whitehead J,¹ Beckett K,⁴ Christie N,⁵ Sleney J,⁶ Barnes J,⁷ Joseph S,⁸ Morriss R.⁹

¹ School of Medicine, Division of Primary Care, Tower Building, University Park, Nottingham, NG7 2RD.

² School of Medicine, Research Design Service East Midlands (RDS EM), Queen's Medical Centre, Nottingham, NG7 2UH

³ Department of Psychology, Nottingham Trent University, NG1 4BU, UK

⁴ University of the West of England, Oakfield House
Bristol, BS8 2BN, UK

⁵ Centre for Transport Studies, University College London, WC1E 6BT, UK

⁶ Department of Sociology, University of Surrey, GU2 7XH, UK

⁷ Loughborough Design School, Loughborough University, LE11 3TU, UK

⁸ School of Education, University of Nottingham, NG8 1BB, UK

⁹ Division of Psychiatry and Applied Psychology, University of Nottingham, NG7 2TU, UK

Correspondence to: Professor Denise Kendrick

(ORCID ID orcid.org/0000-0003-3603-6542)

E mail: denise.kendrick@nottingham.ac.uk

Tel: 0115 8466914

Word count: 2482

Abstract word count: 247

Key words: injuries, work, mental health, cohort study

Abstract

Background: The benefits of work for physical, psychological and financial wellbeing are well documented. Return to work (RTW) after unintentional injury is often delayed, and psychological morbidity may contribute to this delay. The impact of psychological morbidity on RTW after a wide range of unintentional injuries in the UK has not been adequately quantified.

Aims: To quantify the role of psychological factors including anxiety, depression and post-traumatic distress on RTW following unintentional injuries.

Design and Setting: Longitudinal multi-centre prospective study in Nottingham, Bristol, Leicester and Guildford, UK

Method: Participants (n=273) were 16-69 year olds admitted to hospital following unintentional injury and, in paid employment prior to injury. They were surveyed at baseline, 1, 2, 4 and 12 months following injury on demographic and injury characteristics, psychological morbidity and RTW status. Associations between demographic, injury and psychological factors and RTW status were quantified using random effects logistic regression.

Results: The odds of RTW reduced as depression scores one month post-injury increased (OR 0.87, 95%CI 0.79, 0.95) and as length of hospital stay increased (OR 0.91, 95%CI 0.86, 0.96). Those experiencing threatening life events following injury (OR 0.27, 95%CI 0.10, 0.72) and with higher scores on the crisis social support scale (OR 0.93, 95%CI 0.88, 0.99) had a lower odds of RTW. Multiple imputation analysis found similar results except crisis social support did not remain significant.

Conclusion: Primary care professionals can identify patients at risk of delayed RTW who may benefit from management of psychological morbidity and support to RTW.

How this fits in

Injuries are common in working age-adults; resulting in a delayed return to work (RTW) for many.

The benefits of work for physical, psychological and financial wellbeing are well documented.

Depression occurring early in the recovery period, threatening life events after the injury and a longer stay in hospital significantly reduce the odds of RTW in the first year after injury.

Primary care professionals can identify patients at risk of delayed RTW, detect and manage psychological morbidity and provide support to RTW.

Introduction

Being in work benefits physical and mental health,[1] while being out of work can have financial, physical and psychological consequences.[2] The importance of identifying modifiable factors which, if addressed, may help individuals to remain in work is highlighted by The Black Report, "Working for a healthier tomorrow".[2] In England in 2014-15, 319,000 adults aged 16-69 years were admitted to hospital with traumatic injuries or poisoning (external cause codes V01-X59).[3] The annual NHS costs of care in the first 12 months after such injuries has recently been estimated at £1.53 billion.[4] Hospital treated injuries result in substantial health-related work absence; with 17% of emergency department attenders and 43% of hospital admissions not RTW 4 months post-injury.[5] Injuries also account for 10% of sick notes in the UK,[2] and 14% of benefit claimants.[6]

Depression,[7-11] anxiety[12] and post-traumatic stress disorder (PTSD)[13] are common after traumatic injury. For example, a review focussing on road traffic injuries[14] estimated prevalence at 1 year ranged from 21-67% for depression, 4-87% for anxiety and 0-100% for PTSD. These conditions impact negatively on ability to RTW.[8, 13, 15-20] This is illustrated by one study,[20] which found fewer patients with depression (52%) or PTSD (47%) were working 6 months after injury than those without depression (73%) or PTSD (78%).

As depression, anxiety and PTSD are detectable and treatable, it is important to quantify their impact on RTW amongst injured working-age adults in the UK. This paper reports findings on psychological morbidity and RTW from the Impact of Injuries Study, to inform identification and management of these diagnoses post-injury in primary care, and other services such as occupational health.

Methods

Study methods are fully described in the published protocol.[21]

Study design

Multi-centre longitudinal cohort study in four NHS acute hospital trusts with emergency departments in Nottingham, Bristol, Leicester and Guildford, UK. These sites were chosen due to prior experience of recruiting to similar studies.

Participants

Participants were recruited from June 2010-June 2012 within three weeks of hospital admission for unintentional injury. Those aged 16-70 years with a fixed address (to enable follow-up) were eligible. Those with loss of consciousness, amnesia or a Glasgow coma scale of <15 at presentation were excluded due to difficulty distinguishing between head injury sequelae and psychological morbidity.[22] Participants were recruited face-to-face, by post and by phone. Quota sampling by injury type was used from June 2010-May 2011, but subsequently due to slow recruitment, all eligible patients could participate. Only participants in paid employment at the time of injury were included in this paper's analyses.

Data collection

Participants completed self-administered questionnaires at recruitment (baseline) and at 1, 2, 4 and 12 months post-injury. Baseline questionnaires assessed socio-demographic characteristics (age, marital status, ethnicity, number of cars in household, living alone, employment status, area-level deprivation (Index of Multiple Deprivation (IMD) 2010);[23] injury details, long term health conditions, anxiety and depression [24] alcohol problems,[25] substance use,[26] and social functioning.[27] Injury severity was assessed from medical records using the Abbreviated Injury Scale (AIS)[28] grouped into: minor (AIS=1), moderate (AIS=2) and serious to maximum (AIS=3-6) and based on the most severe injury for participants with multiple injuries. Follow-up questionnaires included questions on time off work since injury, self-reported recovery,[29] post-traumatic distress,[30] threatening life events related to the

injury,[31] social support,[32] positive and negative changes in outlook[33] and legal proceedings or compensation claims due to injury. More information about each scale can be found in table 1. Researchers administered structured clinical interviews (SCID)[34] measuring psychological morbidity for all participants at baseline.

[INSERT TABLE 1]

Outcomes

The primary outcome was whether a participant reported RTW. This was a binary outcome at each time point (1, 2, 4 and 12 months post-injury). RTW was defined as being in full or part-time paid employment, working at the specific time point and not prevented from working because of their injury since the previous follow-up time point.

The sample for this analysis comprised participants in paid employment at baseline, who returned the 1 month questionnaire, and at least one subsequent follow-up questionnaire (at 2, 4 or 12 months).

Statistical Analysis

Baseline characteristics were described and compared between the sample described above (responders) and those employed at baseline who returned the 1 month questionnaire but did not return any subsequent follow-up questionnaires (non-responders). Categorical data were compared using chi-square tests, and continuous data using t-tests or Mann-Whitney U tests dependent on distributions.

Proportions of participants who had RTW were calculated and a RTW trajectory was developed.[35] Non-responders and those with missing RTW data were categorised as 'unknown'.

Univariate and multivariable odds ratios with 95% confidence intervals were estimated for RTW using random effects logistic regression to account for repeated measures of RTW at 2, 4 and 12 months. Linearity of continuous predictors was assessed, and non-

linear predictors were categorised into quintiles. Models were built using predictors described in box 1.

[INSERT BOX 1]

Block 2 psychological predictors at 1 month were added in order of statistical significance on univariate analysis, to Block 1 and retained if the likelihood ratio test (LRT) p-value was <0.05 . Correlations between predictors in Blocks 3 and 4 with Block 2 psychological predictors were assessed. Those with correlation coefficients ≥ 0.5 or ≤ -0.5 were excluded from the analysis. Block 3 predictors were added and retained in the model if the LRT $p < 0.05$ or if their removal resulted in a $>10\%$ change in the 1 month psychological predictor odds ratio. Block 4 predictors were added and retained as for Block 3. Interactions between psychological predictors at 1 month and age, sex, and follow-up time were assessed based on LRT $p < 0.01$. Collinearity was assessed by the covariance correlation matrix and variance inflation factors. Model assumptions were checked using deviance residuals. Multiple imputation with chained equations was used to impute missing data for all participants employed at baseline. The imputation model included all predictors in the univariate analysis and the outcome (RTW at 2, 4, and 12 months). Ten datasets were created and combined using Rubin's rules.[36, 37]

Results

Three fifths (393, 59%) of the total 668 study participants were employed at the time of injury. Of these 299 (76%) returned the 1 month follow-up questionnaire; and 273 (91%) returned at least one subsequent follow-up questionnaire and so formed the sample for these analyses. Figure 1 shows the flow of participants in the study.

[INSERT FIGURE 1]

Table 2 shows baseline characteristics of the study sample: 52% were men, ages ranged from 16-69 years and 53% were aged 45 to 64; 66% had an injury of moderate severity; 43% reported single injuries; 62% had an injury of the lower limb; injuries most commonly occurred at work (30%), and were most frequently caused by falls (58%).

[INSERT TABLE 2]

Few (13%) participants returning follow-up questionnaires had RTW at 1 month, 23% had RTW at 2 months, 52% at 4 months, and two thirds at 12 months (67%). Only 6% of participants had fully RTW at all time points; 4% initially RTW, but had not RTW at a later time point; over 50% had a delayed RTW; 8% had not RTW at any time point and 29% had insufficient information to categorise RTW over the full 12 month period.

Univariate and multivariable associations with RTW are shown in tables 3 and 4 respectively. In the final model, a one unit increase in the HADS depression score reduced the odds of RTW by 13% (OR 0.87, 95% CI: 0.79, 0.95), a one unit increase in the number of nights in hospital reduced the odds of RTW by 9% (OR 0.91, 95% CI: 0.86, 0.96), the odds of RTW reduced by 7% per unit increase in crisis support (OR 0.93, 95% CI: 0.88, 0.99); and by 73% for those experiencing threatening life events since the injury (OR 0.27, 95% CI: 0.10, 0.72). Social functioning and negative changes in outlook were excluded due to high correlations with the HADS-D at 1 month. No significant interactions between depression at 1 month and age, sex, and time were found. Variance inflation factors ranged from 1.03 to 3.08.

[INSERT TABLES 3 and 4]

Non-responders were significantly younger ($p < 0.001$), significantly more likely to be male ($p < 0.001$), single ($p < 0.001$), live in disadvantaged areas ($p < 0.001$), and have

scores indicating greater problems with alcohol ($P=0.001$) and drug use ($p=0.01$). Results from the multiple imputation analysis (table 3) were similar to the complete case analysis. Associations between depression at 1 month (OR 0.91, 95%CI 0.85, 0.99), nights in hospital (OR 0.92, 95%CI 0.88, 0.97) and threatening life events (OR 0.42, 95%CI 0.19, 0.92) remained significant. Crisis support no longer remained significantly associated with RTW (OR 0.97, 95%CI 0.92, 1.03).

Discussion

Main findings

One third of participants had not returned to work at 12 months post-injury. A higher depression score at one month post-injury, longer hospital stay, subsequent threatening life events and higher crisis support were associated with significantly lower odds of RTW after injury. Other socio-demographic and injury characteristics were not independently associated with RTW status. Findings for depression, length of stay and threatening life events remained significant in multiple imputation analyses.

Strengths and limitations

The strengths of our study are that it is the first prospective multicentre UK study to quantify the impact of early psychological morbidity on RTW in adults aged 16-69 years admitted to hospital following a wide range of injuries. Our study addressed some limitations of previous studies by including a general injury population with injuries of varying levels of severity, measuring a series of psychological predictors of RTW, and adjusting for many potential confounders. Despite responders differing from non-responders on a range of characteristics, multiple imputation analyses showed most of our findings were robust to missing data. Follow-up rates were higher than, or comparable to studies using similar recruitment methods.[38-42]

The limitations of our study include potential selection bias as thirty percent of eligible patients approached took part and participants and non-participants may have differed

in likelihood of RTW. We were unable to explore the impact of injuries on changes in jobs and in hours of work; and further work is required to address this. The number of participants with some types of injuries was small, limiting analyses to broad injury groupings (upper limb, lower limb, upper and lower limbs, and other injuries; chosen because limb injuries are a major cause of work disability[43-45] and consistent with a systematic review of prognostic factors associated with RTW post-injury).[46] Similarly, small numbers required broad grouping of injury mechanisms. Our study sites came from the midlands and southern England. Occupations vary across the country and “blue-collar work” has previously been associated with lower rates of RTW post injury;[46] hence some care should be taken in generalising our findings to the north of the country. Black and minority ethnic groups were under-represented, potentially limiting generalisability of our findings. Work injuries are likely to have poorer work-related outcomes than those occurring elsewhere.[42, 47, 48] This may be partly explained by depression and PTSD which may be more common after occupational injuries.[8, 49-51] Small numbers of work-related injuries in our study precluded analysis of occupational injury outcomes and their predictors.

The finding of a lower odds of RTW with increasing crisis support was unexpected. The scale we used measured support provided in a crisis, not longer-term social support. Higher levels of crisis support may reflect greater emotional distress or physical impairment, which could both reduce the odds of RTW. Also the short-term nature of crisis support may not provide the buffering effect on depression often seen with longer term social support.

Comparison with other studies

Non-UK studies show post-injury depression and PTSD are associated with delayed RTW[19, 52, 53] but differences in occupations, benefits and compensation systems limit comparability with the UK. A 2010 systematic review of RTW prognostic factors after acute orthopaedic trauma[46] included only 2 small UK studies and we have not found

more recent UK studies. The first cohort study from 2002,[42] recruited 154 injured male hospital admissions and found greater PTSD symptoms were associated with a reduced odds of RTW 18 months post-injury (regression coefficient IES-R avoidance subscale 0.47, 95%CI not reported, $p < 0.001$). The second study from 1992, a review of records of 101 road traffic injury patients receiving compensation[54] found undefined psychological problems (regression coefficient 3.24, standard error (SE) 1.54) and older age (regression coefficient -0.77, SE 0.39) were associated with a reduced likelihood of RTW. We found no association between PTSD symptoms and RTW, once depression was in the multivariable model. This may be explained by depression and PTSD often co-existing.[55] Most (81%) of our participants with moderate or severe PTSD met borderline or case criteria for depression and/or anxiety. In addition, injuries due to assaults[42] and road traffic injuries[54] were more frequent in these studies than in ours, and PTSD may be more common after such injuries. Differences in study populations may explain variation in findings for gender[42] and age[54] and our larger sample size may explain our significant finding for length of hospital stay.

Implications for practice and research

Patients consult frequently in primary care after injury[56] hence a range of primary care professionals, including GPs, are well placed to identify psychological morbidity post injury. Most injured patients have not RTW one month after injury and would be eligible for the Fit For Work service.[57] This provides occupational health assessments and develops a RTW Plan with patients and our findings suggest assessments should include identifying and responding to psychological morbidity. We found a small number of key factors (depression, longer inpatient stays, threatening life events subsequent to injury and greater crisis support) predicted longer work absence. GPs, occupational health services and the Fit For Work service can use these factors to identify those who may benefit from additional help to RTW.

Some GPs may regard depressive symptoms almost as “normal” after an injury and are reluctant to “medicalise unhappiness”. [58] Watchful waiting may be appropriate for short-lived symptoms, but our study clearly shows the negative impact of depressive symptoms lasting one month or more post-injury. The National Institute for Health and Care Excellence (NICE) guideline on recognising and managing depression in adults with chronic physical health problems highlights the high risk of depression, where there is functional impairment. [59] Traumatic injuries requiring hospital admission frequently result in functional impairments for many months; [41, 60] hence these NICE guidelines should be applied. However traumatic injuries present additional challenges for GPs in managing depression, such as impaired mobility limiting access to group-based peer support, talking therapies or to undertaking physical activity, whilst analgesics and adjuvant pain medications may interact with antidepressants.

Further work exploring GPs’ perceptions of psychological problems post-injury, the extent and ways in which they identify, manage and coordinate care for these patients and barriers and facilitators to doing so would be useful. Longitudinal studies assessing the impact of psychological problems on RTW after acute traumatic occupational injuries would also be useful, because this group may be at particular risk of psychological problems.

Funding:

This study was funded by the National Institute for Health Research (NIHR) Collaboration for Leadership in Applied Health Research and Care (CLAHRC) Nottinghamshire Derbyshire and Lincolnshire. RM is currently funded by NIHR CLAHRC East Midlands. The views expressed are those of the authors and not necessarily of the National Health Service, NIHR or the Department of Health.

Ethical approval: for the study was provided by Nottingham Research Ethics Committee 1 (number: 09/H0407/29).

Competing interests: None

Box 1: Predictors and potential predictors of RTW included in the multivariable model building

Block 1 (a-priori predictors): age, sex, study centre, and time.

Block 2 (psychological predictors at 1 month): HADS depression (HADS-D), HADS anxiety (HADS-A), IES avoidance (IES-A), IES intrusion (IES-I), AUDIT and DAST.

Block 3 (potential predictors at baseline): number of psychological morbidities, HADS-D, HADS-A, AUDIT, DAST, long standing illness, work status, ethnicity, deprivation, marital status, length of hospital stay, injury characteristics (severity, number, body part, mechanism and location).

Block 4 (potential predictors at 1 month): social functioning, social support, changes in outlook (positive and negative), threatening life events since injury, pain visual analogue scale, compensation, and litigation.

Table 1: Description and reliability scores for scales used

Variables	Scale used	Description of scale	Cronbach's alpha
Anxiety and Depression	Hospital Anxiety and Depression Scale (HADS)	Measure consists of 7 items measuring depression and 7 items measuring anxiety. Scores range from 0 to 21 for each subscale. Higher scores indicate higher severity of anxiety and depression.	.80 (anxiety) .70 (depression) .83 (overall)
Alcohol problems	Alcohol Use Disorder Identification Test (AUDIT)	Measure consists of 10 items measuring alcohol problems. Scores range from 0 to 40. Higher scores indicate higher levels of excessive or harmful drinking.	.79
Substance use	Drug Abuse Screening Test (DAST)	Measure consists of 10 items measuring drug use. Scores range from 0-10. Higher scores indicate higher levels of abuse of substances other than alcohol.	.67
Social functioning	Social Functioning Questionnaire (SFQ)	Measure consists of 8 items measuring social functioning. Score ranging from 0 to 24. Higher scores indicate greater social dysfunction	.72
Post-traumatic distress	Impact of Events Scale (IES)	Measure consists of 7 items measuring intrusion and 8 measuring avoidance symptoms. Scores range from 0 to 75 overall. Higher scores indicate higher severity of post-traumatic symptoms.	.89 (intrusion) .84 (avoidance) .92 (overall)
Social support	Crisis Support Scale (CSS)	Measure consists of 6 items measuring social support. Scores range from 6-42. Higher scores indicate higher social support.	.76
Positive and negative changes in outlook	Change in Outlook Questionnaire, (CiOQ)	Measure consists of 5 items measuring positive changes and 5 items measuring negative changes. Scores range from 5 to 30 for each subscale. Higher scores indicate respective greater positive and negative changes.	.78 (negative) .87 (positive)

Table 2. Baseline characteristics of study participants.

Characteristics measured at baseline	Participants employed at baseline, returned 1 month questionnaire and at least 1 follow-up questionnaire (n=273)
Centre	
Nottingham	99 (36.3)
Loughborough	74 (27.1)
Bristol	71 (26.0)
Surrey	29 (10.6)
Age	
16-24	29 (10.6)
25-44	92 (33.7)
45-64	145 (53.1)
65-69	7 (2.6)
Sex	
Female	132 (48.4)
Male	141 (51.7)
Number of psychiatric diagnoses in past (obtained from the SCID)	
0	237 (86.8)
1	27 (9.9)
≥2	9 (3.3)
HADS-D	[1]
Median (IQR)	0 (0, 2)
HADS-A	[1]
Median (IQR)	2 (0, 4)
AUDIT scale	[5]
Median (IQR)	4 (2, 6)
DAST scale	[2]
Median (IQR)	0 (0, 0)
Social functioning scale	
Median (IQR)	1 (0, 3)
Pain visual analogue scale	[1]
Median (IQR)	0 (0, 2)
Long standing illness	[1]
No	230 (84.3)
Yes	42 (15.4)
Ethnic group	
White	265 (97.1)
Black or ethnic minority	8 (2.9)
Deprivation (IMD)	[3]
Median (IQR)	12 (7, 20)
Marital status	
Single	74 (27.1)
Married/partnership	164 (60.1)
Divorced/widowed	35 (12.8)

Nights in hospital	[11]
Median (IQR)	5.5 (3, 8)
Injury severity*	[1]
Minor	15 (5.5)
Moderate	206 (75.5)
Serious or worse	51 (18.7)
Number of injuries	
1	117 (42.9)
2	91 (33.3)
≥3	65 (23.8)
Body part injured	
Other	28 (10.3)
Upper limb	49 (18.0)
Lower limb	170 (62.3)
Upper and lower limbs	26 (9.5)
Injury mechanism	
Other	28 (10.3)
Falls	159 (58.2)
Traffic	63 (23.1)
Struck	23 (8.4)
Place of injury	[1]
Other	42 (15.4)
Home	39 (14.3)
Work	83 (30.4)
Road	29 (10.6)
Countryside	38 (13.9)
Sports facilities	41 (15)

[] missing values. SCID= structured clinical interview; HADS-D=HADS depression; HADS-A= HADS anxiety; AUDIT=alcohol use disorder identification test; DAST=drug abuse screening test; IMD = Index of Multiple Deprivation. *Injury severity measured using the Abbreviated Injury Scale (AIS); minor = AIS=1, Moderate = AIS=2, Serious or worse = AIS> =3.

Table 3. Unadjusted odds ratios for potential factors associated with RTW (with 95% confidence intervals).

Variables	Unadjusted OR (95% CI)
A-priori confounders	
Centre	
Nottingham	1.00
Loughborough	0.82 (0.51, 1.31)
Bristol	0.92 (0.57, 1.48)
Surrey	1.96 (1.00, 3.84)
Age	
16-24	1.00
25-44	1.29 (0.67, 2.51)
45-64	0.95 (0.51, 1.76)
65-69	0.65 (0.19, 2.28)
Sex	
Female	1.00
Male	1.04 (0.71, 1.51)
Follow-up Time	
2 months	1.00
4 months	11.72 (6.06, 22.70)
12 months	77.39 (30.91, 193.77)
Psychological predictors at 1 month post-injury	
HADS-D	0.92 (0.88, 0.96)
HADS-A	0.92 (0.88, 0.97)
IES-A	0.96 (0.93, 0.98)
IES-I	0.96 (0.94, 0.99)
AUDIT scale	
1 (0)	1.00
2 (1-2)	1.08 (0.64, 1.83)
3 (2.2-3)	1.73 (0.95, 3.16)
4 (3.3-6)	2.12 (1.19, 3.76)
5 (7-25)	1.16 (0.65, 2.05)
DAST scale	1.08 (0.69, 1.67)
Psychological, social-demographic, and injury characteristics at baseline	

Number of psychiatric diagnoses in past (SCID)	
0	1.00
1	0.88 (0.46, 1.68)
≥2	0.27 (0.09, 0.80)
HADS-D	0.94 (0.85, 1.04)
HADS-A	0.99 (0.93, 1.05)
AUDIT scale	1.00 (0.96, 1.04)
DAST scale	0.92 (0.61, 1.37)
Long standing illness	
No	1.00
Yes	0.76 (0.46, 1.27)
Ethnic group	
White	1.00
Black or minority ethnic group	0.43 (0.12, 1.48)
Deprivation (IMD)	0.98 (0.97, 1.00)
Marital status	
Single	1.00
Married/partnership	1.43 (0.92, 2.22)
Divorced/widowed	1.17 (0.62, 2.20)
Nights in hospital	0.93 (0.90, 0.97)
Injury severity*	
Minor	1.00
Moderate	0.73 (0.32, 1.63)
Serious or worse	0.42 (0.17, 1.02)
Number of injuries	
1	1.00
2	1.12 (0.73, 1.72)
≥3	0.60 (0.38, 0.97)
Body part injured	
Other	1.00
Upper limb	0.95 (0.46, 1.96)
Lower limb	0.55 (0.29, 1.04)
Upper and lower limbs	0.45 (0.19, 1.03)
Injury mechanism	
Other	1.00
Falls	1.00 (0.54, 1.84)
Traffic	0.75 (0.38, 1.49)

Struck	1.10 (0.47, 2.57)
Place of injury	
Other	1.00
Home	0.63 (0.32, 1.22)
Work	0.70 (0.40, 1.24)
Road	0.90 (0.44, 1.85)
Countryside	1.28 (0.64, 2.55)
Sports facilities	0.61 (0.31, 1.17)
Other predictors at 1 month post-injury	
Social functioning scale (Quintiles)	
1 (0-4.6)	1.00
2 (5-6.9)	0.42 (0.24, 0.73)
3 (7-8)	0.31 (0.18, 0.54)
4 (9-10)	0.29 (0.16, 0.52)
5 (10.3-18.3)	0.23 (0.13, 0.42)
CCS scale	1.00 (0.97, 1.04)
CiOQ-P scale	0.96 (0.94, 0.98)
CiOQ-N scale	0.97 (0.93, 1.01)
Threatening life events since injury	
No	1.00
Yes	0.38 (0.22, 0.66)
Pain visual analogue scale	0.98 (0.98, 0.99)
Seeking compensation	
No	1.00
Yes	0.59 (0.38, 0.90)
Involved in litigation	
No	1.00
Yes	0.57 (0.35, 0.94)

SCID= structured clinical interview; HADS-D=HADS depression; HADS-A=HADS anxiety; IES-A=IES avoidance; IES-I=IES intrusion; AUDIT=alcohol use disorder identification test; DAST=drug abuse screening test; IMD = Index of Multiple Deprivation; CSS=Crisis Support Scale; CiOQ-P=Change in outlook questionnaire (positive); CiOQ-N=Change in outlook questionnaire (negative). *Injury severity measured using the Abbreviated Injury Scale (AIS); minor = AIS=1, Moderate = AIS=2, Serious or worse = AIS>3

Table 4. Adjusted odds ratios for psychological predictors at 1 month post-injury associated with RTW in complete case and multiple imputation analysis.

Characteristics	Final model (complete case analysis)	Final model (multiple imputation analysis)
	Odds Ratio (95% CI)	Odds Ratio (95% CI)
A-priori confounders		
Centre		
Nottingham	1.00	1.00
Loughborough	0.86 (0.40, 1.85)	0.90 (0.48, 1.71)
Bristol	0.68 (0.31, 1.48)	0.95 (0.52, 1.73)
Surrey	2.62 (0.92, 7.50)	1.97 (0.81, 4.80)
Age		
16-24	1.00	1.00
25-44	1.13 (0.39, 3.29)	0.95 (0.44, 2.08)
45-64	0.44 (0.16, 1.24)	0.60 (0.27, 1.31)
65-69	0.27 (0.04, 1.99)	0.31 (0.06, 1.68)
Sex		
Female	1.00	1.00
Male	0.73 (0.39, 1.34)	0.79 (0.45, 1.38)
Time		
2 months	1.00	1.00
4 months	10.80 (5.62, 20.76)	7.16 (3.61, 14.19)
12 months	72.17 (29.02, 179.45)	31.78 (13.87, 72.81)
Psychological predictors at 1 month post-injury		
HADS-D	0.87 (0.79, 0.95)	0.91 (0.85, 0.99)
Psychological, socio-demographic, and injury characteristics at baseline		
Nights in hospital	0.91 (0.86, 0.96)	0.92 (0.88, 0.97)
Other predictors at 1 month post-injury		
CSS scale	0.93 (0.88, 0.99)	0.97 (0.92, 1.03)
Threatening life events since injury		
No	1.00	1.00
Yes	0.27 (0.10, 0.72)	0.42 (0.19, 0.92)

HADS-D=HADS depression; CSS=Crisis Social Support.

References

1. Waddell G, Burton AK. Is work good for your health and well-being? - an evidence review London: The Stationery Office; 2006.
2. Black C. 'Working for a healthier tomorrow' - Dame Carol Black's review of the health of Britain's working age population. London: The Stationery Office, 2008.
3. NHS Digital. Hospital Episode Statistics, Admitted Patient Care - England, 2014-15: External causes. Available from <http://content.digital.nhs.uk/searchcatalogue?productid=19420&q=title%3a%22Hospital+Episode+Statistics%2c+Admitted+patient+care+-+England%22&sort=Relevance&size=10&page=1#top> [Accessed 3/1/17].
4. Kellezi B, Baines DL, Coupland C, Beckett K, Barnes J, Slaney J, et al. The impact of injuries on health service resource use and costs in primary and secondary care in the English NHS. *Journal of Public Health*. 2015.
5. Kendrick D, Vinogradova Y, Coupland C, Christie N, Lyons RA, Towner EL. Getting back to work after injury: the UK Burden of Injury multicentre longitudinal study. *BMC public health*. 2012;12:584.
6. Black C, Frost D. Health at work – an independent review of sickness absence. London: Department for Work and Pensions, 2011.
7. Bultmann U, Franche R-L, Hogg-Johnson S, Cote P, Lee H, Severin C, et al. Health status, work limitations, and return-to-work trajectories in injured workers with musculoskeletal disorders. *Quality of Life Research*. 2007;16(7):1167-78. PubMed PMID: 17616838. Pubmed Central PMCID: PMC2039824.
8. Carnide N, Franche R-L, Hogg-Johnson S, Côté P, Breslin FC, Severin CN, et al. Course of Depressive Symptoms Following a Workplace Injury: A 12-Month Follow-Up Update. *J Occup Rehabil*. 2016;26(2):204-15.
9. Lin K-H, Shiao JS-C, Guo N-W, Liao S-C, Kuo C-Y, Hu P-Y, et al. Long-Term Psychological Outcome of Workers After Occupational Injury: Prevalence and Risk Factors. *Journal of Occupational Rehabilitation*. 2014;24(1):1-10.
10. Vles WJ, Steyerberg EW, Essink-Bot M-L, van Beeck EF, Meeuwis JD, Leenen LPH. Prevalence and determinants of disabilities and return to work after major trauma. *Journal of Trauma-Injury Infection & Critical Care*. 2005;58(1):126-35. PubMed PMID: 15674163.
11. Wall CL, Ogloff JR, Morrissey SA. Psychological consequences of work injury: Personality, trauma and psychological distress symptoms of noninjured workers and injured workers returning to, or remaining at work. *The International Journal of Disability Management Research*. 2007;2(2):37-46. PubMed PMID: Peer Reviewed Journal: 2008-18561-002.
12. Noble J, Gomez M, Fish JS. Quality of life and return to work following electrical burns. *Burns*. 2006;32(2):159-64. PubMed PMID: 16448771.
13. Zatzick D, Jurkovich G, Rivara F, Wang J, Fan M, Joesch J, et al. A national US study of posttraumatic stress disorder, depression, and work and functional outcomes after hospitalization for traumatic injury. *Ann Surg*. 2008;248.
14. Blaszczyński A, Gordon K, Silove D, Sloane D, Hillman K, Panasetis P. Psychiatric morbidity following motor vehicle accidents: A review of methodological issues. *Comprehensive Psychiatry*. 1998;39(3):111-21.
15. Ash P, Goldstein SI. Predictors of Returning to Work. *Journal of the American Academy of Psychiatry and the Law Online*. 1995;23(2):205-10.
16. Franche RL, Carnide N, Hogg-Johnson S, Cote P, Breslin FC, Bultmann U, et al. Course, diagnosis, and treatment of depressive symptomatology in workers following a workplace injury: a prospective cohort study. *Can J Psychiatry*. 2009;54(8):534-46.
17. Matthews LR. Work potential of road accident survivors with post-traumatic stress disorder. *Behav Res Ther*. 2005;43(4):475-83.

18. Opsteegh L, Reinders-Messelink HA, Schollier D, Groothoff JW, Postema K, Dijkstra PU, et al. Determinants of return to work in patients with hand disorders and hand injuries. *J Occup Rehabil.* 2009;19(3):245-55.
19. Toien K, Skogstad L, Ekeberg O, Myhren H, Schou Bredal I. Prevalence and predictors of return to work in hospitalised trauma patients during the first year after discharge: a prospective cohort study. *Injury.* 2012;43(9):1606-13. PubMed PMID: 21489524.
20. Ching AC, Wessler L, Blizzard SR, Zusman NL, Yoo JU. The Relation between Depression, PTSD and the Likelihood of Returning to Work After Spine Trauma. *The Spine Journal.*14(11):S6.
21. Kendrick D, O'Brien C, Christie N, Coupland C, Quinn C, Avis M, et al. The impact of injuries study. Multicentre study assessing physical, psychological, social and occupational functioning post injury-a protocol. *BMC public health.* 2011;11(1):963.
22. Jacobson RR. The post-concussional syndrome: physiogenesis, psychogenesis and malingering. An integrative model. *J Psychosom Res.* 1995;39(6):675-93.
23. Department for Communities and Local Government. English indices of deprivation 2010. Available from <https://www.gov.uk/government/statistics/english-indices-of-deprivation-2010>, [Accessed 18/11/16].
24. Bjelland I, Dahl AA, Haug TT, Neckelmann D. The validity of the Hospital Anxiety and Depression Scale: An updated literature review. *Journal of Psychosomatic Research.* 2002;52(2):69-77.
25. Saunders JB, Aasland OG, Babor TF, De La Fuente JR, Grant M. Development of the Alcohol Use Disorders Identification Test (AUDIT): WHO Collaborative Project on Early Detection of Persons with Harmful Alcohol Consumption-II. *Addiction.* 1993;88:791-804.
26. Maisto SA, Carey MP, Carey KB, Gordon CM, Gleason JR. Use of the AUDIT and the DAST-10 to identify alcohol and drug use disorders among adults with a severe and persistent mental illness. *Psychol Assess.* 2000;12(2):186-92. PubMed PMID: 10887764.
27. Tyrer P, Nur U, Crawford M, Karlsen S, MacLean C, Rao B, et al. The Social Functioning Questionnaire: a rapid and robust measure of perceived functioning. *International Journal of Social Psychiatry.* 2005;51(3):265-75.
28. Association of the Advancement of Automotive Medicine. Abbreviated Injury Scale (AIS) 2005 – Update 2008 Manuals. Barrington: AAAM; 2008.
29. Harms L. After the accident: Survivors' perceptions of recovery following road trauma. *Australian Social Work.* 2004;57(2):161-74.
30. Horowitz M, Wilner N, Alvarez W. Impact of Event Scale: a measure of subjective stress. *Psychosom Med.* 1979;41(3):209-18.
31. Brugha T, Bebbington P, Tennant C, Hurry J. The List of Threatening Experiences: a subset of 12 life event categories with considerable long-term contextual threat. *Psychol Med.* 1985;15:189-94.
32. Joseph S, Andrews B, Williams R, Yule W. Crisis support and psychiatric symptomatology in adult survivors of the Jupiter cruise ship disaster. *British Journal of Clinical Psychology.* 1992;31(1):63-73.
33. Joseph S, Linley PA, Andrews L, Harris G, Howle B, Woodward C, et al. Assessing positive and negative changes in the aftermath of adversity: psychometric evaluation of the changes in outlook questionnaire. *Psychological assessment.* 2005;17(1):70.
34. First MB, Spitzer RL, Gibbons M, Williams JBW. Structurd Clinical Interview for DSM-IV Axis I Disorders, research version, Non-Patient Edition APA. 1997.
35. Harcombe H, Langley J, Davie G, Derrett S. Functional status following injury: What recovery pathways do people follow? *Injury.* 2015;46(7):1275-80.
36. Rubin DB. Multiple Imputation for Nonresponse in Surveys. New York: John Wiley & Sons, 2004.
37. Royston JP. Multiple imputation of missing values. *Stata Journal.* 2004;4:227-41.

38. Polinder S, van Beeck EF, Essink-Bot ML, Toet H, Looman CWN, Mulder S, et al. Functional Outcome at 2.5, 5, 9, and 24 Months After Injury in the Netherlands. *Journal of Trauma: Injury, Infection and Critical Care* 2007;62(1):133-41.
39. Wilson R, Hansen P, Langley J, Derrett S. A comparison of injured patient and general population valuations of EQ-5D health states for New Zealand. *Health Qual Life Outcomes*. 2014;12(1):1-7.
40. Kendrick D, Vinogradova Y, Coupland C, Mulvaney CA, Christie N, Lyons RA, et al. Recovery from injury: the UK Burden of Injury Multicentre Longitudinal Study. *Inj Prev*. 2013;19:370-81.
41. Meerding WJ, Looman CWN, Essink-Bot M-L, Toet H, Mulder S, van Beeck EF. Distribution and Determinants of Health and Work Status in a Comprehensive Population of Injury Patients. *Journal of Trauma and Acute Care Surgery*. 2004;56(1):150-61.
42. Mason S, Wardrope J, Turpin G, Rowlands A. Outcomes after Injury: A Comparison of Workplace and Nonworkplace Injury. *J Trauma*. 2002;53(1):98-103.
43. MacKenzie EJ, Bosse MJ, Kellam JF, Pollak AN, Webb LX, Swiontkowski MF, et al. Early predictors of long-term work disability after major limb trauma. *J Trauma*. 2006;61(3):688-94.
44. Pichora D, Grant H. Upper extremity injured workers stratified by current work status: an examination of health characteristics, work limitations and work instability. *The international journal of occupational and environmental medicine*. 2010;1(3):124-31.
45. Read KM, Kufera JA, Dischinger PC, Kerns TJ, Ho SM, Burgess AR, et al. Life-Altering Outcomes after Lower Extremity Injury Sustained in Motor Vehicle Crashes. *Journal of Trauma-Injury Infection & Critical Care*. 2004;57(4):815-23.
46. Clay FJ, Newstead SV, McClure RJ. A systematic review of early prognostic factors for return to work following acute orthopaedic trauma. *Injury*. 2010;41(8):787-803.
47. Schneider JC, Bassi S, Ryan CM. Employment Outcomes After Burn Injury: A Comparison of Those Burned at Work and Those Burned Outside of Work. *J Burn Care Res*. 2011;32(2):294-301. PubMed PMID: 01253092-201103000-00015.
48. Lilley R, Davie G, Langley J, Ameratunga S, Derrett S. Do outcomes differ between work and non-work-related injury in a universal injury compensation system? Findings from the New Zealand Prospective Outcomes of Injury Study. *BMC public health*. 2013;13:995-. PubMed PMID: PMC4015720.
49. Kim J. Depression as a psychosocial consequence of occupational injury in the US working population: findings from the medical expenditure panel survey. *BMC public health*. 2013;13(1):1-10.
50. Ghisi M, Novara C, Buodo G, Kimble MO, Scozzari S, Di Natale A, et al. Psychological Distress and Post-Traumatic Symptoms Following Occupational Accidents. *Behavioral Sciences*. 2013;3(4):587-600. PubMed PMID: PMC4217600.
51. Buodo G, Novara C, Ghisi M, Palomba D. Posttraumatic and Depressive Symptoms in Victims of Occupational Accidents. *Depression Research and Treatment*. 2012;2012:8.
52. Zatzick D, Jurkovich GJ, Rivara FP, Wang J, Fan M-Y, Joesch J, et al. A national US study of posttraumatic stress disorder, depression, and work and functional outcomes after hospitalization for traumatic injury. *Ann Surg*. 2008;248(3):429-37. PubMed PMID: 18791363.
53. Richmond TS, Amsterdam JD, Guo W, Ackerson T, Gracias V, Robinson KM, et al. The Effect of Post-Injury Depression on Return to Pre-injury Function: A Prospective Cohort Study. *Psychol Med*. 2009;39(10):1709-20. PubMed PMID: PMC2741535.
54. Cornes P. Return to work of road accident victims claiming compensation for personal injury. *Injury*. 1992;23(4):256-60. PubMed PMID: 1618568. Epub 1992/01/01.
55. O'Donnell ML, Creamer M, Pattison P. Posttraumatic Stress Disorder and Depression Following Trauma: Understanding Comorbidity. *American Journal of Psychiatry*. 2004;161(8):1390-6. PubMed PMID: 15285964.
56. Kendrick D, Vinogradova Y, Coupland C, Christie N, Lyons R, Towner E. Getting back to work after injury: the UK Burden of Injury multicentre longitudinal study. *BMC public health*. 2012;12(584).

57. Fit For Work: Guidance for GPs.
https://www.gov.uk/government/uploads/system/uploads/attachment_data/file/458954/gp-fit-for-work-gps.pdf. [Accessed 29/9/16].
58. Christie N, Beckett K, Earthy S, Kellezi B, Sleney J, Barnes J, et al. Seeking support after hospitalisation for injury: a nested qualitative study of the role of primary care. *Br J Gen Pract*. 2016;66(642):e24-e31.
59. National Institute for Health and Care Excellence. Depression in adults with a chronic physical health problem: recognition and management. NICE guidelines [CG91] Published date: October 2009.
60. Holbrook TL, Anderson JP, Sieber WJ, Browner D, DB. H. Outcome after major trauma: 12-month and 18-month follow-up results from the Trauma Recovery Project. *Journal of Trauma-Injury Infection & Critical Care* 1999;46(5):765-71; discussion 71-3.