



## Revisiting the Role of Impulsivity and Compulsivity in Problematic Sexual Behaviors

Beáta Bőthe, István Tóth-Király, Marc N. Potenza, Mark D. Griffiths, Gábor Orosz & Zsolt Demetrovics

To cite this article: Beáta Bőthe, István Tóth-Király, Marc N. Potenza, Mark D. Griffiths, Gábor Orosz & Zsolt Demetrovics (2018): Revisiting the Role of Impulsivity and Compulsivity in Problematic Sexual Behaviors, *The Journal of Sex Research*, DOI: [10.1080/00224499.2018.1480744](https://doi.org/10.1080/00224499.2018.1480744)

To link to this article: <https://doi.org/10.1080/00224499.2018.1480744>



Published with license by Taylor & Francis Group, LLC



Published online: 18 Jun 2018.



Submit your article to this journal



View related articles



View Crossmark data

## Revisiting the Role of Impulsivity and Compulsivity in Problematic Sexual Behaviors

Beáta Bőthe<sup>ID</sup>

Doctoral School of Psychology, ELTE Eötvös Loránd University and Institute of Psychology, ELTE Eötvös Loránd University

István Tóth-Király<sup>ID</sup>

Doctoral School of Psychology, ELTE Eötvös Loránd University and Institute of Psychology, ELTE Eötvös Loránd University and Institute of Psychology, ELTE Eötvös Loránd University

Marc N. Potenza<sup>ID</sup>

School of Medicine, Yale University and Connecticut Mental Health Center

Mark D. Griffiths<sup>ID</sup>

Psychology Department, International Gaming Research Unit, Nottingham Trent University

Gábor Orosz<sup>ID</sup>

Institute of Psychology, ELTE Eötvös Loránd University and Institute of Cognitive Neuroscience and Psychology, Hungarian Research Centre for Natural Sciences

Zsolt Demetrovics<sup>ID</sup>

Institute of Psychology, ELTE Eötvös Loránd University

*Impulsivity and compulsivity are transdiagnostic features associated with clinically relevant aspects of psychiatric disorders, including addictions. However, little research has investigated how impulsivity and compulsivity relate to hypersexuality and problematic pornography use. Thus, the aims of the present study were to investigate (a) self-reported impulsivity and compulsivity with respect to hypersexuality and problematic pornography use and (b) the similarities and possible differences between hypersexuality and problematic pornography use in these domains. Utilizing structural equation modeling (SEM) in a large community sample (N = 13,778 participants; female = 4,151, 30.1%), results indicated that impulsivity ( $\beta = .28$ ,  $\beta = .26$ ) and compulsivity ( $\beta = .23$ ,  $\beta = .14$ ) were weakly related to problematic pornography use among men and women, respectively. Impulsivity had a stronger relationship ( $\beta = .41$ ,  $\beta = .42$ ) with hypersexuality than did compulsivity ( $\beta = .21$ ,  $\beta = .16$ ) among men and women, respectively. Consequently, impulsivity and compulsivity may not contribute as substantially to problematic pornography use as some scholars have proposed. On the other hand, impulsivity might have a more prominent role in hypersexuality than in problematic pornography use. Future research should examine further social and situational factors associated with problematic pornography use.*

---

Correspondence should be addressed to Beáta Bőthe, Eötvös Loránd University, Institute of Psychology, Izabella utca 46, Budapest H-1064, Hungary. E-mail: [bothe.beata@ppk.elte.hu](mailto:bothe.beata@ppk.elte.hu)

Color versions of one or more of the figures in the article can be found online at [www.tandfonline.com/hjsr](http://www.tandfonline.com/hjsr).

This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial-NoDerivatives License (<http://creativecommons.org/licenses/by-nc-nd/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited, and is not altered, transformed, or built upon in any way.

Online pornography has become more anonymous, accessible, and affordable over time. One popular pornography Web site's statistics reported approximately 81 million visitors per day and approximately 28.5 billion visits in 2017 (Pornhub.com, 2018). In most cases, pornography viewing is not problematic. However, for some viewers, pornography use may become problematic (estimated at approximately 3.6% of pornography users; Bőthe et al., 2018) and negatively impact functioning by generating impairments in romantic relationships, meeting obligations, and/or achieving other goals (Kraus, Meshberg-Cohen,

Martino, Quinones, & Potenza, 2015; Twohig, Crosby, & Cox, 2009). According to recent case studies (Bostwick & Bucci, 2008; Kraus, Meshberg-Cohen et al., 2015), difficulties in controlling urges to engage in pornography use, a feature which may involve impulsive and compulsive tendencies, may represent a strong obstacle for people to overcome when attempting to reduce problematic pornography use. Problematic pornography use may represent a prominent manifestation of hypersexuality (also referred to as *sexual compulsivity*, *sexual addiction*, or *excessive sexual behavior* in the literature; Kafka, 2010; Karila et al., 2014; Wéry & Billieux, 2017) because in several studies more than 80% of people with hypersexuality have reported excessive/problematic pornography use (Kafka, 2010; Reid et al., 2012). An improved understanding of the similarities and possible differences between problematic pornography use and hypersexuality may help with respect to developing improved interventions. Because both impulsivity and compulsivity have been highly associated with non-substance-related addictive behaviors such as gambling (American Psychiatric Association, 2013; el-Guebaly, Mudry, Zohar, Tavares, & Potenza, 2012; Leeman & Potenza, 2012; World Health Organization, 2017), questions have arisen regarding the extent to which these features may be associated with problematic pornography use and hypersexuality. The aim of the present study was, for the first time, to simultaneously examine the relationships between the two transdiagnostic measures of self-reported impulsivity and compulsivity and specific forms of problematic sexual behaviors (i.e., problematic pornography consumption and hypersexuality).

### The Proposed Obsessive-Compulsive Spectrum Model as Related to Problematic Sexual Behaviors

Over two decades ago, an obsessive-compulsive spectrum model was proposed (Hollander, 1993; Hollander & Wong, 1995) with the conceptualization that different addictions could be juxtaposed on a continuum or spectrum. Disorders were proposed to lie along this spectrum with underestimation of harm being on the impulsive end and overestimation of harm being on the compulsive end (American Psychiatric Association, 2013; Hollander & Benzaquen, 1997). According to the metatheory of Hollander and Wong (1995), sexuality-related compulsions or addictions are closer to the impulsive end of the spectrum. Just over a decade later, Mick and Hollander (2006) proposed that problematic sexual behaviors had both impulsive and compulsive characteristics. However, these models were proposed largely in the absence of empirical data supporting this continuum of impulsivity and compulsivity lying along extreme ends of a continuous spectrum. When examining gambling and substance use disorders, both impulsive and compulsive features have been observed, and individuals with gambling disorders score high on measures of both impulsivity and compulsivity (Leeman & Potenza, 2012; Potenza, 2007). Consequently, questions exist regarding

empirically derived relationships between problematic sexual behaviors and impulsivity and compulsivity.

According to Lochner et al.'s (2005) study, hypersexual disorder may belong to the reward-deficiency cluster rather than impulsive or somatic clusters on the basis of complex clinical interviews with patients with obsessive-compulsive spectrum disorders. However, the *ICD-11* (beta version of the eleventh version of the *International Statistical Classification of Diseases and Related Health Problems*) Working Group on obsessive-compulsive and related disorders suggested that compulsive sexual behavior disorder (hypersexual disorder) should be included under the classification of impulse control disorders in the *ICD-11* (Grant et al., 2014; Kraus et al., 2018; Stein et al., 2016; World Health Organization, 2017) due to its conceptualization and symptomatology (e.g., repeated failure to resist the impulse to engage in sexual behavior despite its long-term negative consequences). However, such a classification has been questioned because compulsive sexual behavior has similar neurobiological features to substance use disorders, indicating that compulsive sexual behavior could be considered an addictive disorder (Potenza, Gola, Voon, Kor, & Kraus, 2017). Thus, there is currently no consensus whether sexuality-related disorders or problems (such as problematic pornography use or hypersexuality) relate to impulsive or compulsive features or whether they should be considered behavioral addictions (e.g., Griffiths, 2016; Kraus, Voon, & Potenza, 2016; Potenza et al., 2017), while noting that these possibilities are not mutually exclusive. Given that no prior studies have simultaneously examined impulsivity and compulsivity as related to hypersexuality and problematic pornography use, there is currently a knowledge gap in this area.

One study investigated compulsivity and impulsivity together as related to pornography use (Wetterneck, Burgess, Short, Smith, & Cervantes, 2012). However, in this study, sexual compulsivity, as opposed to general compulsivity, was assessed. According to that study's results, impulsivity-related features (risk taking and sensation seeking) were positively and weakly correlated with self-reported positive and negative effects of pornography use and the frequency of pornography use. However, after dividing the sample into problematic and nonproblematic users, there were no significant differences between the groups regarding their level of impulsivity. With respect to sexual compulsivity, the positive and negative effects of pornography use and the frequency of pornography use were positively and moderately associated with sexual compulsivity, and there was a significant difference between the problematic and nonproblematic user groups, because individuals in the problematic groups reported 1.5-fold higher levels of sexual compulsivity than the nonproblematic group. This study is the only one that has assessed both impulsivity and (sexual) compulsivity in one model, with few studies having separately examined impulsivity or

compulsivity as related to problematic sexual behaviors, such as hypersexuality and problematic pornography consumption, as discussed in the next section.

### **Impulsivity, Hypersexuality, and Pornography Consumption**

Impulsivity has been related to multiple behaviors relevant to psychiatric problems and disorders (alcohol drinking, Anestis, Selby, & Joiner, 2007; Fischer, Anderson, & Smith, 2004; Fischer & Smith, 2008; compulsive buying, Billieux, Rochat, Rebetez, & Van Der Linden, 2008; eating disorders, Claes, Vandereycken, & Vertommen, 2005; Fischer et al., 2004; Fischer & Smith, 2008) and specific problematic online behaviors or online addictions (such as Internet addiction, Burnay, Billieux, Blairy, & Larøi, 2015; problematic online gaming, Billieux et al., 2011; Zsila et al., 2017; Facebook overuse and problematic series watching, Orosz, Vallerand, Bóthe, Tóth-Király, & Paskuj, 2016). According to Whiteside and Lynam (2001), impulsivity is defined via four dimensions: *sensation seeking* (openness to experiences that might be dangerous and enjoyment of exciting activities), *negative urgency* (the tendency to engage in impulsive behaviors in order to diminish negative emotions and affects, despite the potentially harmful long-term consequences), *lack of perseverance* (difficulties with staying focused on tasks that might be boring and with finishing projects or tasks if distracting stimuli are present), and *lack of premeditation* (acting before thinking about the possible consequences). This original four-dimensional impulsivity model was later complemented with a fifth dimension, namely, *positive urgency* (Billieux et al., 2012; Lynam, Smith, Whiteside, & Cyders, 2006). Positive urgency refers to the tendency to act rashly when experiencing intensive positive emotions. Most research examining the relationships between pornography use and impulsivity or hypersexuality and impulsivity have either applied a unidimensional impulsivity concept or emphasized the role of sensation seeking.

Within the field of hypersexuality, previous research on heterosexual, bisexual, and homosexual men and women has identified a positive but weak association between self-reported impulsive tendencies and hypersexuality. This suggests that people with higher impulsivity are more likely to engage in hypersexual behaviors (Walton, Cantor, & Lykins, 2017). However, in a study examining a combined sample of hypersexual males and healthy community controls, a moderate positive association was found between impulsivity and the level of hypersexuality, a relationship persisting when anxiety, depression, vulnerability, and mindfulness were taken into consideration (Reid, Bramen, Anderson, & Cohen, 2014). In the case of highly sexually active gay and bisexual males, a similar positive, moderate association has been observed between self-reported impulsivity and levels of hypersexuality (Pachankis, Rendina, Ventuneac, Grov, & Parsons, 2014).

However, when hypersexual and nonhypersexual males were compared regarding their level of impulsivity, only a trend toward significance was found in relation to impulsivity (Mulhauser et al., 2014). In another study comparing

impulsivity between hypersexual gay men and nonhypersexual gay men (Miner et al., 2016), only one significant difference was observed. Hypersexual gay men showed elevated levels of nonplanning impulsivity compared to non-hypersexual gay men. There were no significant differences between the two groups in their levels of attentional and motor impulsivity. The aforementioned findings suggest that hypersexuality is related to generalized impulsivity and that hypersexual males are not a homogenous group regarding impulsivity levels (Miner et al., 2016; Mulhauser et al., 2014). However, the findings suggest that impulsivity importantly relates to hypersexuality (Pachankis et al., 2014; Reid et al., 2014; Walton et al., 2017).

Regarding pornography use, sensation seeking is arguably the most prevalently examined impulsivity-related characteristic studied to date. Sensation seeking has been found to be positively related to the frequency of pornography consumption (Beyens, Vandenbosch, & Eggemont, 2015; Peter & Valkenburg, 2010). For men, experience seeking has also been found to be positively related to online pornography use (Paul, 2009). According to Cooper, Delmonico, and Burg's (2000) research, sexually compulsive people and individuals with cybersex addictions score higher on sexual and nonsexual sensation-seeking scales than do non-sexually compulsive and moderately sexually compulsive people. In sum, people with higher levels of sensation seeking may use pornography more intensively as manifested by either an increased amount of time spent with online pornography or the development of problematic online pornography use. Regarding the four other proposed dimensions of impulsivity (negative urgency, positive urgency, lack of perseverance, and lack of premeditation), no previous research has ever examined associations between these variables and online pornography use.

With respect to general impulsivity, the frequency of pornography use has been found to be negatively related to impulsivity among men (i.e., losing one's temper or easily getting irritated), but this was not the case for women (Carroll et al., 2008). In a separate study, low self-control (including impulsivity) explained only a small amount of the variance concerning the frequency of visiting pornography Web sites and downloading pornographic material after controlling for gender and age (Buzzell, Foss, & Middleton, 2006). Other research has found that motivations for using pornography positively and moderately relate to impulsivity across all motivational dimensions investigated (Reid, Li, Gilliland, Stein, & Fong, 2011). Consequently, these data suggest weak but complex relationships between pornography use and impulsivity that do not appear entirely consistent across studies.

In sum, the empirical evidence shows that impulsivity is weakly or moderately related to several aspects of pornography use, such as frequency of pornography use or motivation for pornography viewing (e.g., Beyens et al., 2015; Carroll et al., 2008; Peter & Valkenburg, 2010; Reid et al., 2011). However, little research has focused on the relationship

between impulsivity and problematic pornography use. On the other hand, data suggest that impulsivity relates to hypersexuality, with other personality-related measures also showing relationships (Miner et al., 2016; Mulhauser et al., 2014; Pachankis et al., 2014; Reid et al., 2014; Walton et al., 2017).

### **Compulsivity, Hypersexuality, and Pornography Consumption**

Compulsivity is another personality-related characteristic that has been associated with psychiatric disorders and behaviors (e.g., substance use and gambling disorders, Leeman & Potenza, 2012; compulsive overeating, Davis & Carter, 2009; alcohol abuse and dependence, Modell, Glaser, Mountz, Schmaltz, & Cyr, 1992; bulimia nervosa, Engel et al., 2005). Compulsivity is characterized by the “performance of repetitive and functionally impairing overt or covert behavior without adaptive function, performed in a habitual or stereotyped fashion, either according to rigid rules or as a means of avoiding perceived negative consequences” (Fineberg et al., 2014, p. 70). Therefore, compulsivity may refer to the engagement in ritualistic, repetitive behaviors and actions to prevent or reduce distress or eliminate feared consequences of an individual’s behaviors. However, this feeling of alleviation may be temporary, leading to a vicious cycle where the individual regularly engages in ritualistic actions (Deacon & Abramowitz, 2005).

Few studies have examined associations between compulsivity and hypersexuality. Among males with nonparaphilic hypersexual disorder, the lifetime prevalence of obsessive-compulsive disorder—a psychiatric disorder characterized by compulsivity—ranges from 0% to 14% (Kafka, 2015). Obsessiveness—which may be associated with compulsive behavior (Minnesota Multiphasic Personality Inventory 2 (MMPI-2); Butcher, Dahlstrom, Graham, Tellegen, & Kaemmer, 1989)—in treatment-seeking men with hypersexuality has been found to be elevated relative to a comparison group, but the effect size of this difference was weak (Reid & Carpenter, 2009). When the association between the level of obsessive-compulsive behavior—assessed by a subscale of the Structured Clinical Interview for DSM-IV (SCID-II) (First, Gibbon, Spitzer, Williams, & Benjamin, 1997)—and the level of hypersexuality was examined among treatment-seeking males with hypersexual disorder, a trend toward a positive, weak association was found (Carpenter, Reid, Garos, & Najavits, 2013). On the basis of the aforementioned results, compulsivity appears to contribute in a relatively small manner to hypersexuality.

Sexual compulsivity (more so than general compulsivity) has been associated with pornography use. Among male students, pornography viewing has been found to be positively and moderately associated with sexual compulsivity, with sexual compulsivity mediating the positive association between pornography viewing and problematic behavioral outcomes (Twohig et al., 2009). In line with the effects of thought suppression observed in obsessive-compulsive disorder (e.g., Abramowitz, Tolin, & Street, 2001; Tolin,

Abramowitz, Przeworski, & Foa, 2002), these results suggest that the unwanted urges to use pornography may influence pornography use, leading to ego-dystonic viewing (i.e., pornography viewing in conflict with an individual’s personality and beliefs), which in turn could lead to negative outcomes (i.e., problematic viewing). A moderate positive association between sexual compulsivity and problematic pornography use was reported in a separate convenience sample of males and females (Grubbs, Exline, Pargament, Hook, & Carlisle, 2015). Other research has found that craving for pornography was also positively and moderately related to sexual compulsivity (Kraus & Rosenberg, 2014). These results are consistent with the notion that hypersexuality includes elements of sexual compulsivity (e.g., Kafka, 2010).

In one study, general compulsivity was examined in relation to problematic pornography use among men, showing positive but weak associations (Egan & Parmar, 2013). When investigated in a more complex model, the relationship between general compulsivity and problematic pornography use was mediated by sexual addiction and Internet addiction, as well as an addiction more generally (Egan & Parmar, 2013). Taken together, the associations between compulsivity and hypersexuality and compulsivity and problematic use appear relatively weak (Carpenter et al., 2013; Egan & Parmar, 2013).

### **Examining Impulsivity and Compulsivity With Respect to Hypersexuality and Problematic Pornography Consumption**

Building on prior work (Wetterneck et al., 2012), a subsequent step is the simultaneous examination of general compulsivity and impulsivity and how each of the constructs may relate to problematic pornography use and hypersexuality in the case of men and women. The aims of the present study were to examine impulsivity and compulsivity relative to hypersexuality and problematic pornography use to identify possible similarities and differences in relationships with hypersexuality and problematic pornography use in a large, nonclinical sample and using validated and well-established measures. It was hypothesized that impulsivity and compulsivity would each positively correlate with problematic pornography use and hypersexuality, and that these relationships would be relatively weak but stronger for hypersexuality.

### **Method**

#### **Participants and Procedure**

The present study was conducted in accordance with the approval of the institutional review board (IRB) of the related university and following the Declaration of Helsinki. Informed consent was obtained from all participants. Data collection was conducted in January 2017 via an online questionnaire that was advertised on one of the largest Hungarian news portals as

a research study examining sexual activities. Only individuals aged 18 years old or older were invited to participate in the present study. Participants received detailed information about the aims of the study (i.e., investigation of sexual habits and behaviors of people), and they were assured of anonymity and confidentiality. Subsequently, the participants read and provided informed consent. Completing the questionnaire took approximately 30 minutes.

Overall, 24,372 individuals agreed to participate. However, 7,282 participants quit before completing the scales used in these analyses. Four requirements were established for being included in the present analysis: (1) having watched pornography at least once in the past year, (2) completing the hypersexuality-related scale, (3) completing the compulsivity-related scale, and (4) completing the impulsivity-related scale. Out of 17,090 participants, 1,602 had not watched pornography at least once in the past year; 469 did not complete the hypersexuality-related scale; 899 did not complete the compulsivity-related scale, and 342 did not complete the impulsivity-related scale. Therefore, 13,778 participants met the aforementioned criteria (female = 4,151, 30.1%; did not indicate gender = 72, 0.5%) and were aged between 18 and 76 years ( $M_{age} = 33.52$ ,  $SD_{age} = 10.93$ ). Regarding residency, 7,505 (54.5%) lived in the capital city, 2,133 (15.5%) in county towns, 2,881 (20.9%) in towns, and 1,259 (9.1%) in villages. Regarding level of education, 350 (2.5%) had primary school degrees or less, 541 (3.9%) had vocational degrees, 4,383 (31.8%) had high school degrees, and 8,504 (61.7%) had higher education degrees (bachelor's, master's, or doctoral). Regarding relationship status, 3,198 were single (23.2%), 5,932 were in a relationship (43.1%), 556 were engaged (4.0%), 3,430 were married (24.9%), 384 were divorced (2.8%), 67 were widows/widowers (0.5%), and 211 indicated the "other" option (1.5%). A previously established question was asked to assess the sexual orientation of the participants (Træen, Nilsen, & Stigum, 2006). Based on responses to this question, 11,388 were heterosexual (82.7%), 1,401 were heterosexual with same-sex orientation to some extent (10.2%), 380 were bisexual (2.8%), 99 were homosexual with heterosexuality to some extent (0.7%), 384 were same-sex orientation (2.8%), 16 were asexual (0.1%), 73 were unsure about their sexual orientation (0.5%), and 37 indicated the "other" option (0.3%). Regarding past-year pornography use, participants watched online pornography weekly, and reported spending 26.4 minutes per session ( $SD = 20.5$ ).

## Measures

**UPPS-P Impulsive Behavior Scale (UPPS-P).** The Short UPPS-P Impulsive Behavior Scale (Zsila, Bőthe, Demetrovics, Billieux, & Orosz, 2017) was developed by Billieux et al. (2012) from the original 59-item UPPS-P (Lynam et al., 2006). The Short UPPS-P is a 20-item scale comprising five different impulsivity aspects with four items per dimension: negative urgency (e.g., "When I am upset I often act without thinking"), positive urgency (e.g., "When I am really excited, I tend not to think about the consequences of

my actions"), sensation seeking (e.g., "I sometimes like doing things that are a bit frightening"), lack of premeditation (e.g., "I usually think carefully before doing anything"), and lack of perseverance (e.g., "I generally like to see things through to the end"). All items were scored on a four-point Likert scale (from 1 = *I agree strongly* to 4 = *I disagree strongly*). The facets negative urgency, positive urgency, and sensation seeking include reversed items. Descriptive statistics and the internal consistencies of the scale are shown in Table 1.

## Structured Clinical Interview for DSM Disorders.

The SCID-II (First et al., 1997; Szádóczky, Unoka, & Rózsa, 2004) comprises 140 items covering 10 personality disorders included in *The Diagnostic and Statistical Manual of Mental Disorders*, Fourth Edition (DSM-IV), Axis II and two personality disorders listed in the appendix for diagnoses requiring further studies. In the present research, only the compulsivity subscale was used, which assesses compulsive behavior, utilizing nine true (1) or false (0) items (e.g., "Do you have trouble throwing things out because they might come in handy someday?"). Descriptive statistics and the internal consistency of the scale are shown in Table 1.

**Hypersexual Behavior Inventory (HBI).** The HBI (Bőthe, Bartók et al., 2018; Reid et al., 2011) comprises 19 items assessing the level of hypersexual behavior across three dimensions. The *control* factor (eight items; e.g., "I engage in sexual activities that I know I will later regret") assesses the lack of self-control in sexuality-related behaviors, such as an individual's failed attempts to change his or her sexual behavior. The *coping* factor (seven items; e.g., "Doing something sexual helps me cope with stress") refers to sexual behaviors as a response to emotional distress, such as frustration, sadness, or daily life worries. The *consequences* factor (four items; e.g., "My sexual thoughts and fantasies distract me from accomplishing important tasks") refers to perceived consequences of sexual urges, thoughts, and behaviors, such as sexual activities interfering with important tasks, studies, or work. This scale was translated on the basis of the protocol outlined by Beaton, Bombardier, Guillemin, and Ferraz (2000). All items are scored on a five-point Likert scale (from 1 = *Never* to 5 = *Very often*). Descriptive statistics and the internal consistency of the scale are shown in Table 1.

**Problematic Pornography Consumption Scale (PPCS).** The PPCS (Bőthe, Tóth-Király et al., 2018) is based on a proposed six-component addictions model (Griffiths, 2005). The PPCS is an 18-item scale that assesses problematic pornography consumption using six factors, with three items relating to each factor. *Tolerance* captures when increasing amounts of the activity are required to achieve the same mood-modifying effects (e.g., "I felt that I had to watch more and more porn for satisfaction"). *Salience* refers to the relevance of pornography in an individual's life (e.g., "I felt that porn is an important part of my life"). *Mood modification* is an either arousing or relaxing subjective experience that users report as a consequence of viewing pornography (e.g., "I

**Table 1.** Descriptive Statistics, Reliability Indices and Correlations Between the Aspects of Impulsivity, Compulsivity, Hypersexuality and Problematic Pornography Use

Scales	Skewness (SE)	Kurtosis (SE)	Range	<i>M</i> ( <i>SD</i> )	<i>a</i>	1	2	3	4	5	6	7	8
1. PPCS total	1.61 (0.02)	2.61 (0.04)	1–7	1.92 (1.00)	.94	—							
2. HBI total	1.24 (0.02)	1.85 (0.04)	1–5	1.76 (0.57)	.89	.57*	—						
3. UPPS-P total	0.06 (0.02)	-0.10 (0.04)	1–4	2.28 (0.41)	.85	.15*	.31*	—					
4. UPPS-P negative urgency	0.17 (0.02)	-0.57 (0.04)	1–4	2.38 (0.73)	.83	.13*	.24*	.76*	—				
5. UPPS-P positive urgency	-0.05 (0.02)	-0.25 (0.04)	1–4	2.61 (0.63)	.73	.13*	.29*	.80*	.64*	—			
6. UPPS-P sensation seeking	-0.10 (0.02)	-0.24 (0.04)	1–4	2.58 (0.63)	.77	.04*	.15*	.47*	.17*	.39*	—		
7. UPPS-P lack of premeditation	0.38 (0.02)	-0.03 (0.04)	1–4	1.90 (0.58)	.82	.04*	.14*	.67*	.36*	.35*	.04*	—	
8. UPPS-P lack of perseverance	0.41 (0.02)	-0.05 (0.04)	1–4	1.91 (0.59)	.83	.14*	.18*	.52*	.18*	.16*	-.09*	.49*	—
9. SCID-II compulsivity	-0.15 (0.02)	-0.41 (0.04)	0–1	0.54 (0.21)	.49 <sup>a</sup>	.13*	.14*	.02*	.17*	.11*	.04*	-.19*	-.11*

Note. PPCS = Problematic Pornography Consumption Scale; HBI = Hypersexual Behavior Inventory; UPPS-P = UPPS-P Impulsive Behavior Scale; SCID-II = Structured Clinical Interview for DSM Disorders; *a* = Cronbach's alpha; *M* = mean; *SD* = standard deviation; SE = standard error.

<sup>a</sup>Internal consistency was examined by assessment of Kuder-Richardson formula 20 (KR-20) for this dichotomous scale.


\**p* < .001.

released my tension by watching porn"). *Conflict* includes interpersonal conflicts between problematic users and their significant others, intrapsychic conflicts (e.g., knowing the activity is generating problems but finding difficulties consuming less or quitting), and occupational or educational concerns (e.g., "I felt porn caused problems in my sexual life"). *Relapse* is the tendency for returning to pornography quickly after abstinence or control (e.g., "I unsuccessfully tried to reduce the amount of porn I watch"). Last, *withdrawal* refers to the unpleasant feelings and emotional states that occur when the particular activity is decreased or ceased (e.g., "I became stressed when something prevented me from watching porn"). All items are scored on a 7-point Likert scale (from 1 = *Never* to 7 = *Very often*). Descriptive statistics and the internal consistency of the scale are shown in Table 1.

## Statistical Analyses

For the statistical analysis, SPSS 21 and Mplus 7.3 (Muthén & Muthén, 1998–2015) were used. Normality was assessed by the investigation of skewness and kurtosis. Reliability was assessed using Cronbach's alpha (Nunnally, 1978) in the case of continuous scales. For the one dichotomous scale used (i.e., compulsivity subscale of SCID-II), internal consistency was examined with Kuder-Richardson formula 20 (KR-20, Kuder & Richardson, 1937). Structural equation modeling (SEM) was used to explore the associations between impulsivity, compulsivity, hypersexuality, and problematic pornography use. Items were treated as categorical indicators, because they had significant floor effects (on the basis of kurtosis and skewness). Consequently, the mean- and variance-adjusted weighted least squares estimator (WLSMV) was applied (Finney & DiStefano, 2006). Commonly used goodness-of-fit indices (Brown, 2015; Kline, 2011) were observed (Bentler, 1990; Brown, 2015; Browne & Cudeck, 1993; Hu & Bentler, 1999; Schermelleh-Engel, Moosbrugger, & Müller, 2003; Tabachnick & Fidell, 2001) to assess the acceptability of the proposed model. The analyses examined the comparative fit index (CFI; ≥ .95 for good, ≥ .90 for acceptable), the Tucker-Lewis index (TLI; ≥ .95 for good, ≥ .90 for acceptable), and the root mean square error of approximation (RMSEA; ≤ .06 for good, ≤ .08 for acceptable) with a 90% confidence interval (CI).

In the case of the SCID-II compulsivity subscale and the HBI items, a parceling approach was conducted due to the fact that these latent variables were assessed using many items. Parcels are aggregated items that were used in the present model as assessed variables. This approach is acceptable in the case of theoretically unidimensional scales (e.g., Bandalo & Finney, 2001; Little, Cunningham, Shahar, & Widaman, 2002; Orosz et al., 2016), and it can minimize the issues related to non-normally distributed data (Bandalo, 2002; Matsunaga, 2008). In the case of the SCID-II compulsivity subscale, Rogers and Schmitt's (2004) exploratory factor analysis-based algorithm was applied in the parcel construction. For the HBI, a facet-representative approach was employed (Little, Rhemtulla, Gibson, & Schoemann, 2013) and each subscale (namely, coping, control, and



**Figure 1.** The impulsivity and compulsivity background of hypersexuality and problematic pornography use ( $N_{\text{total}} = 13,778$ ;  $N_{\text{males}} = 9,555$ ;  $N_{\text{females}} = 4,151$ ). All variables presented in ellipses are latent variables. For the sake of clarity, indicator variables related to them are not depicted in this figure. One-headed arrows represent standardized regression weights and two-headed arrows represent correlations. The first numbers on the arrows indicate the path coefficients of the total sample, the second numbers indicate the path coefficients of the male sample, and the third numbers indicate the path coefficients of the female sample. All pathways were significant at level  $p < .01$ .

consequences) was averaged. Consequently, three indicators were constructed.

## Results

Descriptive data, reliability indices, and correlations between the aspects of impulsivity, compulsivity, hypersexuality, and problematic pornography use are shown in Table 1. According to the correlations, there were only small differences between the correlations of problematic pornography use, hypersexuality, and the specific aspects of impulsivity. Therefore, for the sake of simplicity, the total score of impulsivity was used in further analyses.

By using SEM, the associations between impulsivity, compulsivity, hypersexuality, and problematic pornography use were investigated in the total sample and in the case of men and women in separate models as well. The models with standardized estimates are shown in Figure 1.

In the total sample model, the fit indices were acceptable ( $\text{CFI} = .941$ ,  $\text{TLI} = .937$ ,  $\text{RMSEA} = .055$  [90% CI = .054–.055]). Both impulsivity and compulsivity were related positively but weakly to problematic pornography use ( $\beta = .17$ ,  $p < .01$ , and  $\beta = .19$ ,  $p < .01$ , respectively). The proportion of explained variance of problematic pornography use was 6.6%. In the case of hypersexuality, compulsivity was also positively but weakly related to hypersexuality ( $\beta = .19$ ,  $p < .01$ ). However, impulsivity was positively but moderately related to hypersexuality ( $\beta = .37$ ,  $p < .01$ ). The proportion of explained variance of hypersexuality was 18.1%.

In the male sample model, the fit indices were acceptable ( $\text{CFI} = .929$ ,  $\text{TLI} = .924$ ,  $\text{RMSEA} = .059$  [90% CI = .058–.059]). Both impulsivity and compulsivity were related positively but weakly to problematic pornography use ( $\beta = .28$ ,  $p < .01$ , and  $\beta = .23$ ,  $p < .01$ , respectively). The proportion of explained variance of problematic pornography use was 13.2%. In the case of hypersexuality, compulsivity was also positively but weakly related to hypersexuality ( $\beta = .21$ ,  $p < .01$ ). However, impulsivity was positively but moderately related to hypersexuality ( $\beta = .41$ ,  $p < .01$ ). The proportion of explained variance of hypersexuality was 21.7%.

.23,  $p < .01$ , respectively). The proportion of explained variance of problematic pornography use was 13.2%. In the case of hypersexuality, compulsivity was also positively but weakly related to hypersexuality ( $\beta = .21$ ,  $p < .01$ ). However, impulsivity was positively but moderately related to hypersexuality ( $\beta = .41$ ,  $p < .01$ ). The proportion of explained variance of hypersexuality was 21.7%.

In the female sample model, the fit indices were acceptable ( $\text{CFI} = .914$ ,  $\text{TLI} = .908$ ,  $\text{RMSEA} = .055$  [90% CI = .054–.056]). Both impulsivity and compulsivity were related positively but weakly to problematic pornography use ( $\beta = .26$ ,  $p < .01$ , and  $\beta = .14$ ,  $p < .01$ , respectively). The proportion of explained variance of problematic pornography use was 9.1%. In the case of hypersexuality, compulsivity was also positively but weakly related to hypersexuality ( $\beta = .16$ ,  $p < .01$ ). However, impulsivity was positively but moderately related to hypersexuality ( $\beta = .42$ ,  $p < .01$ ). The proportion of explained variance of hypersexuality was 21.0%.

In sum, the associations between problematic pornography use and impulsivity and compulsivity, respectively, were weak, and the explained variance of problematic pornography use by impulsivity and compulsivity was relatively low (6.6% to 13.2%) in the total sample, as well as among men and women. In the case of hypersexuality, impulsivity had a stronger effect on hypersexual behavior than compulsivity, with hypersexuality having an explained variance of approximately 20% by impulsivity and compulsivity in the total sample, as well as among men and women.

## Discussion

There is a current debate regarding how best to consider problematic sexual behaviors (such as hypersexuality and problematic pornography use), with competing models

proposing classifications as impulse-control disorders, obsessive-compulsive spectrum disorders, or behavioral addictions (e.g., Griffiths, 2016; Kraus et al., 2016; Potenza et al., 2017). Relationships between transdiagnostic features of impulsivity and compulsivity and problematic sexual behaviors should inform such considerations, although both impulsivity and compulsivity have been implicated in addictions (Fineberg et al., 2014; Leeman & Potenza, 2012). The present study contributes to the ongoing debate by examining and identifying differences in the relationships between measures of self-reported impulsivity, compulsivity, hypersexuality, and problematic pornography consumption.

The findings of the present study showed that impulsivity was moderately and positively related to hypersexual behavior, while compulsivity was only weakly related, suggesting that impulsivity contributes more strongly to hypersexuality than compulsivity in both men and women. However, impulsivity and compulsivity related only weakly to problematic pornography use among both genders. From a statistical perspective, impulsivity and compulsivity both positively predicted problematic pornography use, but the effect sizes were small in both cases and the proportion of explained variance of problematic pornography use did not reach 15%, suggesting that more emphasis should be put on other factors (e.g., social and society related) in research and clinical interventions in the case of problematic pornography use. On the other hand, the finding that impulsivity related moderately to hypersexuality provides support both for the classification of compulsive sexual behavior disorder (as proposed for *ICD-11*; World Health Organization, 2017) as an impulse-control disorder or as a behavioral addiction. In considering the other disorders currently being proposed as impulse-control disorders (e.g., intermittent explosive disorder, pyromania, and kleptomania) and the central elements of compulsive sexual behavior disorder and proposed disorders due to addictive behaviors (e.g., gambling and gaming disorders), the classification of compulsive sexual behavior disorder in the latter category appears better supported.

The findings of the present study suggest that problematic pornography use may differ from hypersexuality more generally. As such, considering specific forms of excessive or problematic sexual behaviors will be important because different individuals with different temperamental features may be vulnerable to, and experience problems with, different types of sexual behavior.

### **The Role of the Impulsivity and Compulsivity in Hypersexuality and Problematic Pornography Consumption**

Impulsivity and compulsivity are among the most frequently examined personality-related factors in the case of problematic behaviors with addictive potential (e.g., Billieux et al., 2008; Davis & Carter, 2009; Deckman & DeWall, 2011; Engel et al., 2005; Leeman & Potenza, 2012; Mottram & Fleming, 2009). However, little research has examined the associations of impulsivity, compulsivity,

and problematic sexual behaviors (such as hypersexuality and problematic pornography use). This small body of work reports relatively small effect sizes and inconsistent results. However, no prior study to this one has ever simultaneously investigated the relationships of impulsivity and compulsivity with hypersexuality and problematic pornography use.

Regarding pornography use motivations (Reid et al., 2011), impulsivity was positively and moderately related to almost all motivational factors, whereas, in the case of frequency of pornography use, a less consistent pattern was observed, from positive associations to no association (e.g., Beyens et al., 2015; Carroll et al., 2008; Peter & Valkenburg, 2011). Only one study (i.e., Wetterneck et al., 2012) has investigated the associations between impulsivity, compulsivity, and problematic pornography use simultaneously. Similar to the results of the present study, positive but weak associations between the variables were observed, and after dividing the sample into problematic and nonproblematic users, no significant differences were found between the groups regarding levels of impulsivity. Therefore, impulsivity may not be as relevant to problematic pornography use as previously proposed (e.g., Hollander & Wong, 1995; Mick & Hollander, 2006).

In the case of hypersexuality, research has shown that impulsivity is weakly or moderately associated with hypersexual behaviors, fantasies, and urges (Pachankis et al., 2014; Reid et al., 2014; Walton et al., 2017). However, comparisons of hypersexual and nonhypersexual individuals have not shown consistent results (Miner et al., 2016; Mulhauser et al., 2014). The results of the present study corroborate the findings of Pachankis et al. (2014) and Reid et al. (2014) because the associations between impulsivity and hypersexuality are positive and moderate, suggesting that impulsivity may contribute importantly to the development and maintenance of hypersexuality.

Regarding compulsivity, associations between pornography use and sexual compulsivity have been more widely investigated than those between pornography use and general compulsivity. Not surprisingly, when sexual compulsivity was assessed in relation to pornography viewing (e.g., Grubbs, Exline et al., 2015; Twohig et al., 2009; Wetterneck et al., 2012), the association was moderate and positive. Several possible reasons for this relationship have been proposed. First, context-specific compulsivity may be expected to be more strongly related to problematic pornography use than context-free (i.e., general) compulsivity. Second, hypersexuality by definition may include sexual compulsivity (e.g., Kafka, 2010). However, when general compulsivity has been assessed as an antecedent of problematic pornography use, similar to the results of the present study, positive but weak associations were observed (Egan & Parmar, 2013). Previously, general compulsivity or obsessiveness was only weakly related or unrelated to hypersexuality (e.g., Carpenter et al., 2013; Reid & Carpenter, 2009). In the present study, similar relationships were observed because general compulsivity (from a statistical

perspective) significantly predicted hypersexuality, but the effect size was low.

In the present study, the five-facet model of impulsivity (Billieux et al., 2012; Lynam et al., 2006) was examined in relation to problematic pornography use and hypersexuality. The five facets—namely, negative urgency, positive urgency, lack of premeditation, lack of perseverance, and sensation seeking—were generally positively but weakly related to problematic pornography use and positively and moderately to hypersexuality, demonstrating a consistent relationship pattern between the aspects of impulsivity and problematic pornography use and hypersexuality. Therefore, the total score of impulsivity was used as a statistical predictor of problematic pornography use and hypersexuality. As expected, impulsivity was positively associated with problematic pornography use and hypersexuality. However, the extent of the association between impulsivity and problematic pornography use was rather small.

### Possible Explanations for the Weak Associations Between Impulsivity, Compulsivity, and Problematic Pornography Use

Several factors may explain why impulsivity and compulsivity only weakly statistically predicted the level of problematic pornography use while hypersexuality was moderately statistically predicted by impulsivity. It is possible that impulsivity and compulsivity do not have a strong direct impact on problematic pornography use but have stronger effects via mediating variables. In the case of impulsivity, Reid et al. (2011) found that impulsivity had positive moderate associations with all four motivations of pornography use. From their four motivational factors, emotional avoidance had the strongest relationship with impulsivity, with excitement seeking being the second strongest and sexual pleasure being the third strongest, whereas sexual curiosity had the weakest relationship with impulsivity. Based on these results, emotional avoidance motivation may represent a mediator between impulsivity and problematic pornography use, although direct investigation of this possibility is needed to confirm the hypothesis.

Moreover, the frequency of pornography use may also serve as a potential mediator between impulsivity and problematic pornography use. For men, impulsivity has been found to be positively related to the frequency of pornography viewing; for females, it was not related (Carroll et al., 2008). As men tend to have higher levels of impulsivity (e.g., Chapple & Johnson, 2007; Cross, Copping, & Campbell, 2011; Waldeck & Miller, 1997), it could be hypothesized that this elevated level of impulsivity may lead to an increased frequency of pornography use, which in turn may lead to problematic pornography use (e.g., Brand et al., 2011; Grubbs, Exline et al., 2015; Grubbs, Volk et al., 2015; Twohig et al., 2009). For women, impulsivity was not related to the frequency of pornography use (Carroll et al., 2008); therefore, it could be assumed that their impulsivity may not reflect in the frequency of pornography leading to problematic pornography use, but

problematic pornography use could develop via different pathways (e.g., Lewczuk, Szmyd, Skorko, & Gola, 2017). In Egan and Parmar's (2013) study, the association between compulsivity and problematic pornography use was mediated by sexual addiction, Internet addiction, and addiction more generally. Therefore, a similar mediational pattern could be hypothesized regarding the association between compulsivity and hypersexuality.

Likewise, self-efficacy may also mediate possible relationships between impulsivity, compulsivity, and problematic pornography use. In previous studies (e.g., Kraus, Rosenberg, Martino, Nich, & Potenza, 2017; Kraus, Rosenberg, & Tompsett, 2015), self-efficacy in reducing pornography use and self-efficacy in avoiding possibly tempting situations were identified as important factors in reducing problematic pornography use. Therefore, one might hypothesize that people with high levels of impulsivity or compulsivity may control their urges because of their high level of self-efficacy to avoid tempting situations, which in turn may result in lower levels of problematic pornography use.

Nevertheless, it is possible that levels of impulsivity and compulsivity in relationships with problematic sexual behaviors (such as problematic pornography use and hypersexuality) have been overestimated. According to a number of scholars (e.g., Conway, Kane, Ball, Poling, & Rounsville, 2003; Griffiths, 2017; Kerr, 1996; Szalavitz, 2016), no single personality trait or set of traits may lead to problematic behaviors or addictions. The three cornerstones of online pornography use (anonymity, affordability, and accessibility; Cooper, 1998) may create situations that facilitate the increased use of pornography, and these may also contribute to the development of problematic pornography use. The careful, experimental examination of these cornerstones may significantly contribute to the understanding of problematic pornography use. Furthermore, situation-related factors that may affect individuals in a given life stage, such as loneliness (e.g., Bozoglan, Demirer, & Sahin, 2013; Ceyhan & Ceyhan, 2008) or perceived stress (e.g., Grubbs, Volk, et al., 2015; Levin, Lillis, & Hayes, 2012; Paul & Shim, 2008; Reid et al., 2011), may also influence the level of addictive online behaviors such as problematic pornography use. Finally, it should also be noted that societal factors such as regulations and policies that influence the accessibility, affordability, and anonymity of pornography may in turn promote or hinder the emergence of those specific situations in which (problematic or nonproblematic) pornography use may have significant psychosocial impacts.

### Future Studies and Limitations

Further measures are needed in future studies that may directly assess respondents' behaviors while respecting individuals' privacy. The extent of problematic pornography consumption and the level of hypersexuality may be temporally stable or it may change over time. It is possible that an individual may temporarily use pornography more intensively

or in a more problematic manner, but this behavior may change. Therefore, longitudinal studies are needed to answer the question of stability. Future experimental studies with well-established designs are needed to determine a potentially causal role of individual differences and situational factors in the development and maintenance of hypersexuality and problematic pornography use, such as sex mind-set beliefs (Böthe, Tóth-Király, Demetrovics, & Orosz, 2017), reward deficiency syndrome (Comings & Blum, 2000; Lochner et al., 2005), perceived stress (Grubbs, Volk, Exline, & Pargament, 2015), or basic psychological needs (Tóth-Király, Morin, Böthe, Orosz, & Rigó, 2018). Finally, it should be kept in mind that the outcomes studied in the present study pertain only to specific problematic aspects of sexuality (i.e., problematic online pornography use and hypersexuality). Developing measures that can assess nonproblematic aspects of pornography use may be useful in further research. Stronger cooperation between pornography Web sites—which may provide behavioral data—and the scientific community may be beneficial in providing predictive validity of related measures. Future studies should focus on preventions and interventions that emphasize not only self-reported individual differences but also social and situational factors related to the development and maintenance of problematic sexual behaviors.

Some limitations of the present study should be noted. The use of self-report cross-sectional methods have possible biases that need be considered when interpreting the findings. Moreover, causality cannot be inferred from the present cross-sectional findings. The internal consistency of the compulsivity subscale of the SCID-II was not adequate; therefore, it is possible that the low level of internal consistency may have distorted the findings. In addition, self-reported compulsivity was assessed via SCID-II methods. Other assessments of compulsivity (e.g., via the Padua Inventory or other assessments; Andrews et al., 2011; Scherrer, Xian, Slutske, Eisen, & Potenza, 2015) may have yielded different results. Similar concerns exist regarding the UPPS-P and other self-report measures of impulsivity. In addition, because self-report measures differ with behavioral measures of constructs (e.g., Krishnan-Sarin et al., 2007), it is important for future studies to investigate both behavioral and self-report measures related to hypothesized predictors of the given behavior (e.g., using the cued go/no-go task [Fillmore, 2003] or the stop signal task [Logan, 1994] with a self-reported measure in the case of impulsivity [Ding et al., 2014]). It will also be important to concurrently assess the behavior itself (e.g., actual amount of pornography use employing tracking-data approaches in collaboration with pornography Web site operators as has been done in other fields such as gambling; Griffiths, 2014).

### Conclusion and Implications

In sum, impulsivity and compulsivity did not contribute as importantly and directly to problematic pornography use as

previously proposed in the literature, and impulsivity may have a more prominent role in hypersexuality. Furthermore, these results have several conceptual and research implications. First, several issues arise regarding the categorization of problematic pornography use. One issue is whether problematic pornography use may be considered a subcategory of hypersexuality if relationships with impulsivity and compulsivity are not as strong as previously hypothesized. A second issue—which may be related to the categorization of problematic pornography use under the umbrella of hypersexuality—is how problematic pornography use (and especially problematic online pornography use) may best be categorized (Griffiths, 2016; Kraus et al., 2016; Potenza et al., 2017).

From a research perspective, self-reported tendencies may have a stronger impact on problematic sexual behaviors via mediating variables such as motivations, frequency and time spent with the activity, frustration relating to psychological needs, beliefs about the malleability of the given activities, topic-relevant self-efficacy beliefs, and/or other factors. All of these possibilities warrant direct examination. Moreover, it is important to consider the complex etiologies of addictions. More specifically, it is likely that a complex set of personality factors, other individual difference factors, and social and situational factors lead to the development and maintenance of problematic sexual behaviors and that these may vary according to the type of problematic sexual behavior. Additional research is needed to understand factors related to specific problematic sexual behaviors and translate the factors into improved prevention, treatment, and policy initiatives.

### Funding

The research was supported by the Hungarian National Research, Development, and Innovation Office (PD116686, FK124225, K111938, KKP126835) and the Hungarian Academy of Sciences (Lendület Project LP2012-36). The second author (ITK) was supported by the ÚNKP-17-3 New National Excellence Program of the Ministry of Human Capacities. MNP's involvement was supported by the National Center for Responsible Gaming through a Center of Excellence grant. The funding agencies did not have input into the content of the manuscript, and the views described in the manuscript reflect those of the authors and not necessarily those of the funding agencies.

### Conflict of Interest

The authors declare no conflict of interest with respect to the content of this manuscript. Dr. Potenza has consulted for and advised Rivermend Health, Opiant/Lightlake Therapeutics, and Jazz Pharmaceuticals; received research support (to Yale) from the Mohegan Sun Casino and the National Center for Responsible Gaming; consulted for legal and gambling entities on issues related to impulse control and addictive behaviors.

The other authors report no financial relationships with commercial interests.

## ORCID

- Beáta Bóthe  <http://orcid.org/0000-0003-2718-4703>  
 István Tóth-Király  <http://orcid.org/0000-0003-2810-3661>  
 Marc N. Potenza  <http://orcid.org/0000-0002-6323-1354>  
 Mark D. Griffiths  <http://orcid.org/0000-0001-8880-6524>  
 Gábor Orosz  <http://orcid.org/0000-0001-5883-6861>  
 Zsolt Demetrovics  <http://orcid.org/0000-0001-5604-7551>

## References

- Abramowitz, J. S., Tolin, D. F., & Street, G. P. (2001). Paradoxical effects of thought suppression: A meta-analysis of controlled studies. *Clinical Psychology Review*, 21, 683–703. doi:10.1016/S0272-7358(00)00057-X
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Washington, DC: Author.
- Andrews, M. M., Meda, S. A., Thomas, A. D., Potenza, M. N., Krystal, J. H., Worhunsky, P., ... Pearlson, G. D. (2011). Individuals family history positive for alcoholism show functional magnetic resonance imaging differences in reward sensitivity that are related to impulsivity factors. *Biological Psychiatry*, 69, 675–683. doi:10.1016/j.biopsych.2010.09.049
- Anestis, M. D., Selby, E. A., & Joiner, T. E. (2007). The role of urgency in maladaptive behaviors. *Behaviour Research and Therapy*, 45, 3018–3029. doi:10.1016/j.brat.2007.08.012
- Bandalos, D. L. (2002). The effects of item parceling on goodness-of-fit and parameter estimate bias in structural equation modeling. *Structural Equation Modeling*, 9, 78–102. doi:10.1207/S15328007SEM0901\_5
- Bandalos, D. L., & Finney, S. J. (2001). Item parceling issues in structural equation modeling. In G. A. Marcoulides & R. E. Schumacker (Eds.), *New developments and techniques in structural equation modeling* (pp. 269–296). London, UK: Lawrence Erlbaum.
- Beaton, D. E., Bombardier, C., Guillemin, F., & Ferraz, M. B. (2000). Guidelines for the process of cross-cultural adaptation of self-report measures. *Spine*, 25, 3186–3191.
- Bentler, P. M. (1990). Comparative fit indexes in structural models. *Psychological Bulletin*, 107, 238–246. doi:10.1037/0033-2909.107.2.238
- Beyens, I., Vandenbosch, L., & Eggermont, S. (2015). Early adolescent boys' exposure to internet pornography relationships to pubertal timing, sensation seeking, and academic performance. *Journal of Early Adolescence*, 35, 1045–1068. doi:10.1177/0272431614548069
- Billieux, J., Chanal, J., Khazaal, Y., Rochat, L., Gay, P., Zullino, D., & Van Der Linden, M. (2011). Psychological predictors of problematic involvement in massively multiplayer online role-playing games: Illustration in a sample of male cybercafe players. *Psychopathology*, 44, 165–171. doi:10.1159/000322525
- Billieux, J., Rochat, L., Ceschi, G., Carré, A., Offerlin-Meyer, I., Defeldre, A. C., ... Van Der Linden, M. (2012). Validation of a short French version of the UPPS-P impulsive behavior scale. *Comprehensive Psychiatry*, 53, 609–615. doi:10.1016/j.comppsych.2011.09.001
- Billieux, J., Rochat, L., Rebetez, M. M. L., & Van Der Linden, M. (2008). Are all facets of impulsivity related to self-reported compulsive buying behavior? *Personality and Individual Differences*, 44, 1432–1442. doi:10.1016/j.paid.2007.12.011
- Bostwick, J. M., & Bucci, J. A. (2008). Internet sex addiction treated with naltrexone. *Mayo Clinic Proceedings*, 83, 226–230.
- Bóthe, B., Bartók, R., Tóth-Király, I., Reid, R. C., Griffiths, M. D., Demetrovics, Z., & Orosz, G. (2018). Hypersexuality, gender, and sexual orientation: A large-scale psychometric survey study. *Archives of Sexual Behavior*. doi:10.1007/s10508-018-1201-z
- Bóthe, B., Tóth-Király, I., Demetrovics, Z., & Orosz, G. (2017). The pervasive role of sex mindset: Beliefs about the malleability of sexual life is linked to higher levels of relationship satisfaction and sexual satisfaction and lower levels of problematic pornography use. *Personality and Individual Differences*, 117, 15–22. doi:10.1016/j.paid.2017.05.030
- Bóthe, B., Tóth-Király, I., Zsila, Á., Griffiths, M. D., Demetrovics, Z., & Orosz, G. (2018). The development of the problematic pornography consumption scale (PPCS). *Journal of Sex Research*, 55, 395–406. doi:10.1080/00224499.2017.1291798
- Bozoglan, B., Demirer, V., & Sahin, I. (2013). Loneliness, self-esteem, and life satisfaction as predictors of internet addiction: A cross-sectional study among Turkish university students. *Scandinavian Journal of Psychology*, 54(4), 313–319. doi:10.1111/sjop.12049
- Brand, M., Laier, C., Pawlikowski, M., Schächtle, U., Schöler, T., & Altstötter-Gleich, C. (2011). Watching pornographic pictures on the internet: Role of sexual arousal ratings and psychological–Psychiatric symptoms for using internet sex sites excessively. *Cyberpsychology, Behavior, and Social Networking*, 14, 371–377. doi:10.1089/cyber.2010.0222
- Brown, T. A. (2015). *Confirmatory factor analysis for applied research* (2nd ed.). New York, NY: Guilford Press.
- Browne, M. V., & Cudeck, R. (1993). Alternative ways of assessing model fit. In K. A. Bollen & J. S. Long (Eds.), *Testing structural equation models* (pp. 136–162). Newbury Park, CA: Sage.
- Burnay, J., Billieux, J., Blairy, S., & Larøi, F. (2015). Which psychological factors influence internet addiction? Evidence through an integrative model. *Computers in Human Behavior*, 43, 28–34. doi:10.1016/j.chb.2014.10.039
- Butcher, J. N., Dahlstrom, W. G., Graham, J. R., Tellegen, A., & Kaemmer, B. (1989). *MMPI-2: Manual for administration and scoring*. Minneapolis, MN: University of Minnesota Press.
- Buzzell, T., Foss, D., & Middleton, Z. (2006). Explaining use of online pornography: A test of self-control theory and opportunities for deviance. *Journal of Criminal Justice and Popular Culture*, 13, 96–116.
- Carpenter, B. N., Reid, R. C., Garos, S., & Najavits, L. M. (2013). Personality disorder comorbidity in treatment-seeking men with hypersexual disorder. *Sexual Addiction & Compulsivity*, 20, 79–90. doi:10.1080/10720162.2013.772873
- Carroll, J. S., Padilla-Walker, L. M., Nelson, L. J., Olson, C. D., Barry, C. M., & Madsen, S. D. (2008). Generation XXX pornography acceptance and use among emerging adults. *Journal of Adolescent Research*, 23, 6–30. doi:10.1177/0743558407306348
- Ceyhan, A. A., & Ceyhan, E. (2008). Loneliness, depression, and computer self-efficacy as predictors of problematic internet use. *CyberPsychology & Behavior*, 11, 699–701. doi:10.1089/cpb.2007.0255
- Chapple, C. L., & Johnson, K. A. (2007). Gender differences in impulsivity. *Youth Violence and Juvenile Justice*, 5, 221–234. doi:10.1177/1541204007301286
- Claes, L., Vandereycken, W., & Vertommen, H. (2005). Impulsivity-related traits in eating disorder patients. *Personality and Individual Differences*, 39, 739–749. doi:10.1016/j.paid.2005.02.022
- Comings, D. E., & Blum, K. (2000). Reward deficiency syndrome: Genetic aspects of behavioral disorders. *Progress in Brain Research*, 126, 325–341. doi:10.1016/S0079-6123(00)26022-6
- Conway, K. P., Kane, R. J., Ball, S. A., Poling, J. C., & Rounsvaille, B. J. (2003). Personality, substance of choice, and polysubstance involvement among substance dependent patients. *Drug and Alcohol Dependence*, 71, 65–75. doi:10.1016/S0376-8716(03)00068-1
- Cooper, A. (1998). Sexuality and the internet: Surfing into the new millennium. *CyberPsychology & Behavior*, 1, 187–193. doi:10.1089/cpb.1998.1.187
- Cooper, A., Delmonico, D. L., & Burg, R. (2000). Cybersex users, abusers, and compulsives: New findings and implications. *Sexual Addiction & Compulsivity*, 7, 5–29. doi:10.1080/10720160008400205
- Cross, C. P., Copping, L. T., & Campbell, A. (2011). Sex differences in impulsivity: A meta-analysis. *Psychological Bulletin*, 137, 97–130.
- Davis, C., & Carter, J. C. (2009). Compulsive overeating as an addiction disorder. A review of theory and evidence. *Appetite*, 53, 1–8. doi:10.1016/j.appet.2009.05.018

- Deacon, B. J., & Abramowitz, J. S. (2005). The Yale-Brown Obsessive-Compulsive Scale: Factor analysis, construct validity, and suggestions for refinement. *Journal of Anxiety Disorders*, 19, 573–585. doi:10.1016/j.janxdis.2004.04.009
- Deckman, T., & DeWall, C. N. (2011). Negative urgency and risky sexual behaviors: A clarification of the relationship between impulsivity and risky sexual behavior. *Personality and Individual Differences*, 51, 674–678. doi:10.1016/j.paid.2011.06.004
- Ding, W. N., Sun, J. H., Sun, Y. W., Chen, X., Zhou, Y., Zhuang, Z. G., ... Du, Y. S. (2014). Trait impulsivity and impaired prefrontal impulse inhibition function in adolescents with internet gaming addiction revealed by a Go/No-Go fMRI study. *Behavioral and Brain Functions*, 10(1), 20. doi:10.1186/1744-9081-10-20
- Egan, V., & Parmar, R. (2013). Dirty habits? online pornography use, personality, obsessiveness, and compulsivity. *Journal of Sex & Marital Therapy*, 39, 394–409. doi:10.1080/0092623X.2012.710182
- el-Guebaly, N., Mudry, T., Zohar, J., Tavares, H., & Potenza, M. N. (2012). Compulsive features in behavioural addictions: The case of pathological gambling. *Addiction*, 107, 1726–1734. doi:10.1111/j.1360-0443.2011.03546.x
- Engel, S. G., Corneliusen, S. J., Wonderlich, S. A., Crosby, R. D., Le Grange, D., Crow, S., ... Mitchell, J. E. (2005). Impulsivity and compulsivity in bulimia nervosa. *International Journal of Eating Disorders*, 38, 244–251. doi:10.1002/eat.20169
- Fillmore, M. T. (2003). Drug abuse as a problem of impaired control: Current approaches and findings. *Behavioral and Cognitive Neuroscience Reviews*, 2, 179–197. doi:10.1177/1534582303257007
- Fineberg, N. A., Chamberlain, S. R., Goudriaan, A. E., Stein, D. J., Vandershuren, L. J., Gillan, C. M., ... Denys, D. (2014). New developments in human neurocognition: Clinical, genetic, and brain imaging correlates of impulsivity and compulsivity. *CNS Spectrums*, 19, 69–89. doi:10.1017/S1092852913000801
- Finney, S. J., & DiStefano, C. (2006). Non-normal and categorical data in structural equation modeling. In G. R. Hancock & R. D. Mueller (Eds.), *Structural equation modeling: A second course* (pp. 269–314). Charlotte, NC: Information Age Publishing.
- First, M. B., Gibbon, M., Spitzer, R. L., Williams, J. B. W., & Benjamin, L. S. (1997). *SCID-II personality questionnaire*. Washington, DC: American Psychiatry Press.
- Fischer, S., Anderson, K. G., & Smith, G. T. (2004). Coping with distress by eating or drinking: Role of trait urgency and expectancies. *Psychology of Addictive Behaviors*, 18, 269–274. doi:10.1037/0893-164X.18.3.269
- Fischer, S., & Smith, G. T. (2008). Binge eating, problem drinking, and pathological gambling: Linking behavior to shared traits and social learning. *Personality and Individual Differences*, 44, 789–800. doi:10.1016/j.paid.2007.10.008
- Grant, J. E., Atmaca, M., Fineberg, N. A., Fontenelle, L. F., Matsunaga, H., Janardhan Reddy, Y. C., ... Woods, D. W. (2014). Impulse control disorders and “behavioural addictions” in the ICD-11. *World Psychiatry*, 13, 125–127. doi:10.1002/wps.20115
- Griffiths, M. (2005). A ‘components’ model of addiction within a biopsychosocial framework. *Journal of Substance Use*, 10, 191–197. doi:10.1080/14659890500114359
- Griffiths, M. D. (2014). The use of behavioural tracking methodologies in the study of online gambling. *SAGE Research Methods Cases*. doi:10.4136/978144627305013517480
- Griffiths, M. D. (2016). Compulsive sexual behaviour as a behavioural addiction: The impact of the internet and other issues. *Addiction*, 111, 2107–2108. doi:10.1111/add.13315
- Griffiths, M. D. (2017). The myth of ‘addictive personality’. *Global Journal of Addiction & Rehabilitation Medicine*, 3, 555610. doi:10.19080/GJARM.2017.03.555610
- Grubbs, J. B., Exline, J. J., Pargament, K. I., Hook, J. N., & Carlisle, R. D. (2015). Transgression as addiction: Religiosity and moral disapproval as predictors of perceived addiction to pornography. *Archives of Sexual Behavior*, 44, 125–136. doi:10.1007/s10508-013-0257-z
- Grubbs, J. B., Volk, F., Exline, J. J., & Pargament, K. I. (2015). Internet pornography use: Perceived addiction, psychological distress, and the validation of a brief measure. *Journal of Sex & Marital Therapy*, 41, 83–106. doi:10.1080/0092623X.2013.842192
- Hollander, E. (1993). Obsessive-compulsive spectrum disorders: An overview. *Psychiatric Annals*, 23, 255–358.
- Hollander, E., & Benzaquen, S. D. (1997). The obsessive-compulsive spectrum disorders. *International Review of Psychiatry*, 9, 99–110. doi:10.1080/09540269775628
- Hollander, E., & Wong, C. M. (1995). Obsessive-compulsive spectrum disorders. *Journal of Clinical Psychiatry*, 56(Suppl 4), 3–6.
- Hu, L., & Bentler, P. M. (1999). Cutoff criteria for fit indexes in covariance structure analysis: Conventional criteria versus new alternatives. *Structural Equation Modeling*, 6, 1–55. doi:10.1080/1070551990540118
- Kafka, M. P. (2010). Hypersexual disorder: A proposed diagnosis for DSM-V. *Archives of Sexual Behavior*, 39, 377–400. doi:10.1007/s10508-009-9574-7
- Kafka, M. P. (2015). DSM-IV Axis I psychopathology in males with non-paraphilic hypersexual disorder. *Current Addiction Reports*, 2, 202–206. doi:10.1007/s40429-015-0060-0
- Karila, L., Wéry, A., Weinstein, A., Cottencin, O., Petit, A., Reynaud, M., & Billieux, J. (2014). Sexual addiction or hypersexual disorder: Different terms for the same problem? A review of the literature. *Current Pharmaceutical Design*, 20, 4012–4020.
- Kerr, J. S. (1996). Two myths of addiction: The addictive personality and the issue of free choice. *Human Psychopharmacology*, 11, S9–S14.
- Kline, R. B. (2011). *Principles and practice of structural equation modeling* (3rd ed.). New York, NY: Guilford Press.
- Kraus, S. W., Krueger, R. B., Briken, P., First, M. B., Stein, D. J., Kaplan, M. S., ... Reed, G. M. (2018). Compulsive sexual behaviour disorder in the ICD-11. *World Psychiatry*, 17, 109–110. doi:10.1002/wps.20499
- Kraus, S. W., Meshberg-Cohen, S., Martino, S., Quinones, L. J., & Potenza, M. N. (2015). Treatment of compulsive pornography use with naltrexone: A case report. *American Journal of Psychiatry*, 172, 1260–1261. doi:10.1176/appi.ajp.2015.15060843
- Kraus, S. W., & Rosenberg, H. (2014). The pornography craving questionnaire: Psychometric properties. *Archives of Sexual Behavior*, 43, 451–462. doi:10.1007/s10508-013-0229-3
- Kraus, S. W., Rosenberg, H., Martino, S., Nich, C., & Potenza, M. N. (2017). The development and initial evaluation of the pornography-use avoidance self-efficacy scale. *Journal of Behavioral Addictions*, 6, 354–363. doi:10.1556/2006.6.2017.057
- Kraus, S. W., Rosenberg, H., & Tompsett, C. J. (2015). Assessment of self-efficacy to employ self-initiated pornography use-reduction strategies. *Addictive Behaviors*, 40, 115–118. doi:10.1016/j.addbeh.2014.09.012
- Kraus, S. W., Voon, V., & Potenza, M. N. (2016). Should compulsive sexual behavior be considered an addiction? *Addiction*, 111, 2097–2106. doi:10.1111/add.13297
- Krishnan-Sarin, S., Reynolds, B., Duhig, A. M., Smith, A., Liss, T., McFetridge, A., ... Potenza, M. N. (2007). Behavioral impulsivity predicts treatment outcome in a smoking cessation program for adolescent smokers. *Drug and Alcohol Dependence*, 88, 79–82. doi:10.1016/j.drugalcdep.2006.09.006
- Kuder, G. F., & Richardson, M. W. (1937). The theory of the estimation of test reliability. *Psychometrika*, 2, 151–160. doi:10.1007/BF02288391
- Leeman, R. F., & Potenza, M. N. (2012). Similarities and differences between pathological gambling and substance use disorders: A focus on impulsivity and compulsivity. *Psychopharmacology*, 219, 469–490. doi:10.1007/s00213-011-2550-7

- Levin, M. E., Lillis, J., & Hayes, S. C. (2012). When is online pornography viewing problematic among college males? Examining the moderating role of experiential avoidance. *Sexual Addiction & Compulsivity*, 19, 168–180. doi:10.1080/10720162.2012.657150
- Lewczuk, K., Szmyd, J., Skorko, M., & Gola, M. (2017). Treatment seeking for problematic pornography use among women. *Journal of Behavioral Addictions*, 6, 445–456. doi:10.1556/2006.6.2017.063
- Little, T. D., Cunningham, W. A., Shahar, G., & Widaman, K. F. (2002). To parcel or not to parcel: Exploring the question, weighing the merits. *Structural Equation Modeling*, 9, 151–173. doi:10.1207/S15328007SEM0902\_1
- Little, T. D., Rhemtulla, M., Gibson, K., & Schoemann, A. M. (2013). Why the items versus parcels controversy needn't be one. *Psychological Methods*, 18, 285–300. doi:10.1037/a0033266
- Lochner, C., Hemmings, S. M., Kinnear, C. J., Niehaus, D. J., Nel, D. G., Corfield, V. A., ... Stein, D. J. (2005). Cluster analysis of obsessive-compulsive spectrum disorders in patients with obsessive-compulsive disorder: Clinical and genetic correlates. *Comprehensive Psychiatry*, 46, 14–19. doi:10.1016/j.comppsych.2004.07.020
- Logan, G. D. (1994). On the ability to inhibit thought and action: A users' guide to the stop signal paradigm. In D. Dagenbach & T. H. Carr (Eds.), *Inhibitory processes in attention, memory, and language* (pp. 189–239). San Diego, CA: Academic Press.
- Lynam, D. R., Smith, G. T., Whiteside, S. P., & Cyders, M. A. (2006). *The UPPS-P: Assessing five personality pathways to impulsive behavior*. Technical report. West Lafayette, IN: Purdue University.
- Matsunaga, M. (2008). Item parceling in structural equation modeling: A primer. *Communication Methods and Measures*, 2, 260–293. doi:10.1080/19312450802458935
- Mick, T. M., & Hollander, E. (2006). Impulsive-compulsive sexual behavior. *CNS Spectrums*, 11, 944–955. doi:10.1017/S1092852900015133
- Miner, M. H., Romine, R. S., Raymond, N., Janssen, E., MacDonald, A., & Coleman, E. (2016). Understanding the personality and behavioral mechanisms defining hypersexuality in men who have sex with men. *Journal of Sexual Medicine*, 13, 1323–1331. doi:10.1016/j.jsxm.2016.06.015
- Modell, J. G., Glaser, F. B., Mountz, J. M., Schmaltz, S., & Cyr, L. (1992). Obsessive and compulsive characteristics of alcohol abuse and dependence: Quantification by a newly developed questionnaire. *Alcoholism: Clinical and Experimental Research*, 16, 266–271. doi:10.1111/j.1530-0277.1992.tb01374.x
- Mottram, A. J., & Fleming, M. J. (2009). Extraversion, impulsivity, and online group membership as predictors of problematic internet use. *CyberPsychology & Behavior*, 12, 319–321. doi:10.1089/cpb.2007.0170
- Mulhauser, K. R., Struthers, W. M., Hook, J. N., Pykkonen, B. A., Womack, S. D., & MacDonald, M. (2014). Performance on the Iowa gambling task in a sample of hypersexual men. *Sexual Addiction & Compulsivity*, 21, 170–183. doi:10.1080/10720162.2014.908333
- Muthén, L. K., & Muthén, B. O. (1998–2012). *Mplus user's guide* (7th ed.). Los Angeles, CA: Muthén & Muthén.
- Nunnally, J. C. (1978). Psychometric theory. In *McGraw-Hill series in psychology* (2nd ed.). New York, NY: McGraw-Hill.
- Orosz, G., Vallerand, R. J., Bóthe, B., Tóth-Király, I., & Paskuj, B. (2016). On the correlates of passion for screen-based behaviors: The case of impulsivity and the problematic and non-problematic Facebook use and TV series watching. *Personality and Individual Differences*, 101, 167–176. doi:10.1016/j.paid.2016.05.368
- Pachankis, J. E., Rendina, H. J., Ventuneac, A., Grov, C., & Parsons, J. T. (2014). The role of maladaptive cognitions in hypersexuality among highly sexually active gay and bisexual men. *Archives of Sexual Behavior*, 43, 669–683. doi:10.1007/s10508-014-0261-y
- Paul, B. (2009). Predicting Internet pornography use and arousal: The role of individual difference variables. *Journal of Sex Research*, 46, 344–357. doi:10.1080/00224490902754152
- Paul, B., & Shim, J. W. (2008). Gender, sexual affect, and motivations for internet pornography use. *International Journal Of Sexual Health*, 20, 187–199. doi:10.1080/19317610802240154
- Peter, J., & Valkenburg, P. M. (2010). Processes underlying the effects of adolescents' use of sexually explicit internet material: The role of perceived realism. *Communication Research*, 37, 375–399. doi:10.1177/0093650210362464
- Peter, J., & Valkenburg, P. M. (2011). The use of sexually explicit internet material and its antecedents: a longitudinal comparison of adolescents and adults. *Archives Of Sexual Behavior*, 40, 1015–1025. doi:10.1007/s10508-010-9644-x
- Pornhub's. 2017 Year in review (2018, January 9). Retrieved January 12, 2018, from <https://www.pornhub.com/insights/2017-year-in-review>
- Potenza, M. N. (2007). Impulsivity and compulsivity in pathological gambling and obsessive-compulsive disorder. *Revista Brasileira De Psiquiatria*, 29, 105–106. doi:10.1590/S1516-44462007000200004
- Potenza, M. N., Gola, M., Voon, V., Kor, A., & Kraus, S. W. (2017). Is excessive sexual behaviour an addictive disorder? *The Lancet Psychiatry*, 4, 663–664. doi:10.1016/S2215-0366(17)30316-4
- Reid, R. C., Bramen, J. E., Anderson, A., & Cohen, M. S. (2014). Mindfulness, emotional dysregulation, impulsivity, and stress proneness among hypersexual patients. *Journal of Clinical Psychology*, 70, 313–321. doi:10.1002/jclp.22027
- Reid, R. C., & Carpenter, B. N. (2009). Exploring relationships of psychopathology in hypersexual patients using the MMPI-2. *Journal of Sex & Marital Therapy*, 35, 294–310. doi:10.1080/00926230902851298
- Reid, R. C., Carpenter, B. N., Hook, J. N., Garos, S., Manning, J. C., Gilliland, R., ... Fong, T. (2012). Report of findings in a DSM-5 field trial for hypersexual disorder. *The Journal of Sexual Medicine*, 9, 2868–2877. doi:10.1111/j.1743-6109.2012.02936.x
- Reid, R. C., Li, D. S., Gilliland, R., Stein, J. A., & Fong, T. (2011). Reliability, validity, and psychometric development of the pornography consumption inventory in a sample of hypersexual men. *Journal of Sex & Marital Therapy*, 37, 359–385. doi:10.1080/0092623X.2011.607047
- Rogers, W. M., & Schmitt, N. (2004). Parameter recovery and model fit using multidimensional composites: A comparison of four empirical parceling algorithms. *Multivariate Behavioral Research*, 39, 379–412. doi:10.1207/S15327906MBR3903\_1
- Schermelleh-Engel, K., Moosbrugger, H., & Müller, H. (2003). Evaluating the fit of structural equation models: Tests of significance and descriptive goodness-of-fit measures. *Methods of Psychological Research Online*, 8, 23–74.
- Scherrer, J. F., Xian, H., Slutskie, W. S., Eisen, S. A., & Potenza, M. N. (2015). Associations between obsessive-compulsive classes and pathological gambling in a national cohort of male twins. *JAMA Psychiatry*, 72, 342–349. doi:10.1001/jamapsychiatry.2014.2497
- Stein, D. J., Kogan, C. S., Atmaca, M., Fineberg, N. A., Fontenelle, L. F., Grant, J. E., ... Van Den Heuvel, O. A. (2016). The classification of obsessive-compulsive and related disorders in the ICD-11. *Journal of Affective Disorders*, 190, 663–674. doi:10.1016/j.jad.2015.10.061
- Szádóczky, E., Unoka, Z., & Rózsa, S. (2004). *User's guide for the structured clinical interview for DSM-IV axis II personality disorders (SCID-II), Hungarian version*. Budapest, Hungary: OS Hungary Kft.
- Szalavitz, M. (2016). *Unbroken brain: A revolutionary new way of understanding addiction*. New York, NY: St. Martin's Press.
- Tabachnick, B. G., & Fidell, L. S. (2001). *Using multivariate statistics* (4th ed.). Boston, MA: Allyn and Bacon.
- Tolin, D. F., Abramowitz, J. S., Przeworski, A., & Foa, E. B. (2002). Thought suppression in obsessive-compulsive disorder. *Behaviour Research and Therapy*, 40, 1255–1274. doi:10.1016/S0006-3705(01)00095-X
- Tóth-Király, I., Morin, A. J., Bóthe, B., Orosz, G., & Rigó, A. (2018). Investigating the multidimensionality of need fulfillment: A bifactor exploratory structural equation modeling representation. *Structural Equation Modeling: A Multidisciplinary Journal*, 25, 267–286. doi:10.1080/10705511.2017.1374867

- Træen, B., Nilsen, T. S. R., & Stigum, H. (2006). Use of pornography in traditional media and on the internet in norway. *Journal of Sex Research*, 43, 245–254.
- Twohig, M. P., Crosby, J. M., & Cox, J. M. (2009). Viewing internet pornography: For whom is it problematic, how, and why? *Sexual Addiction & Compulsivity*, 16, 253–266. doi:10.1080/10720160903300788
- Waldeck, T. L., & Miller, L. S. (1997). Gender and impulsivity differences in licit substance use. *Journal of Substance Abuse*, 9, 269–275. doi:10.1016/S0899-3289(97)90021-3
- Walton, M. T., Cantor, J. M., & Lykins, A. D. (2017). An online assessment of personality, psychological, and sexuality trait variables associated with self-reported hypersexual behavior. *Archives of Sexual Behavior*, 46, 721–733. doi:10.1007/s10508-015-0606-1
- Wéry, A., & Billieux, J. (2017). Problematic cybersex: Conceptualization, assessment, and treatment. *Addictive Behaviors*, 64, 238–246. doi:10.1016/j.addbeh.2015.11.007
- Wetterneck, C. T., Burgess, A. J., Short, M. B., Smith, A. H., & Cervantes, M. E. (2012). The role of sexual compulsivity, impulsivity, and experiential avoidance in internet pornography use. *Psychological Record*, 62, 3–18.
- Whiteside, S. P., & Lynam, D. R. (2001). The five factor model and impulsivity: Using a structural model of personality to understand impulsivity. *Personality and Individual Differences*, 30, 669–689. doi:10.1016/S0191-8869(00)00064-7
- World Health Organization. (2017). *International statistical classification of diseases and related health problems*. (11th ed. beta version). Retrieved December 8, 2017, from <https://icd.who.int/dev11/l-m/en#/http%3a%2f%2fid.who.int%2fidc%2fentity%2f1630268048>
- Zsila, Á., Bőthe, B., Demetrovics, Z., Billieux, J., & Orosz, G. (2017). Further exploration of the SUPPS-P impulsive behavior scale's factor structure: Evidence from a large Hungarian sample. *Current Psychology*, 1–11. doi:10.1007/s12144-017-9773-7
- Zsila, Á., Orosz, G., Bőthe, B., Tóth-Király, I., Király, O., Griffiths, M., & Demetrovics, Z. (2017). An empirical study on the motivations underlying augmented reality games: The case of Pokémon go during and after Pokémon fever. *Personality and Individual Differences*. doi:10.1016/j.paid.2017.06.024