

1 **Global terrestrial distribution of penguins (*Spheniscidae*) and their conservation by**
2 **protected areas**

3

4

5 Rachel P. Hickcox^{1,2,5}, Manuel Jara^{2,3}, Laura A. K. Deacon², Lilly P. Harvey⁴ & Daniel
6 Pincheira-Donoso^{4,5}

7

8 ¹Department of Zoology, University of Otago, Dunedin 9054, New Zealand

9 ²Laboratory of Evolutionary Ecology of Adaptations, School of Life Sciences, Joseph Banks

10 Laboratories, University of Lincoln, Brayford Campus, Lincoln, LN6 7DL, Lincolnshire, United
11 Kingdom

12 ³Department of Population Health and Pathobiology, College of Veterinary Medicine, North
13 Carolina State University, Raleigh, North Carolina 27606, United States

14 ⁴School of Science and Technology, Nottingham Trent University, Clifton Campus, Nottingham,
15 NG11 8NS, United Kingdom

16 ⁵Authors for correspondence: rachel.hickcox@postgrad.otago.ac.nz;

17 daniel.pincheiradonoso@ntu.ac.uk

18

19 **ABSTRACT**

20 Establishing protected areas (PAs) ranks among the top priority actions to mitigate the global
21 scale of modern biodiversity declines. However, the distribution of biodiversity is spatially
22 asymmetric among regions and lineages, and the extent to which PAs offer effective protection
23 for species and ecosystems remains uncertain. Penguins, regarded as prime bioindicator birds of
24 the ecological health of their terrestrial and marine habitats, represent priority targets for such
25 quantitative assessments. Of the world's 18 penguin species, eleven are undergoing population
26 declines, of which ten are classified as 'Vulnerable' or 'Endangered'. Here, we employ a global-
27 scale dataset to quantify the extent to which their terrestrial breeding areas are currently
28 protected by PAs. Using quantitative methods for spatial ecology, we compared the global
29 distribution of penguin colonies, including range and population size analyses, with the
30 distribution of terrestrial PAs classified by the International Union for Conservation of Nature,
31 and generated hotspot and endemism maps worldwide. Our assessment quantitatively reveals
32 <40% of the terrestrial range of eleven penguin species is currently protected, and that range size
33 is the significant factor in determining PA protection. We also show that there are seven global
34 hotspots of penguin biodiversity where four or five penguin species breed. We suggest that
35 future penguin conservation initiatives should be implemented based on more comprehensive,
36 quantitative assessments of the multi-dimensional interactions between areas and species to
37 further the effectiveness of PA networks.

38 **Keywords:** biodiversity hotspots, IUCN, macroecology, penguins, protected areas, species
39 richness

40

41 **ACKNOWLEDGMENTS**

42 The data used are derived from public repositories. We thank UNEP World Conservation
43 Monitoring Centre, IUCN World Commission on Protected Areas (World Database on Protected
44 Areas), IUCN Global Species Programs and the Species Survival Commission, OBIS, GBIF, and
45 Birdlife International for developing the source data. Thank you to Professor Philip Seddon,
46 James Hunter, Saif Khan, and anonymous reviewers for manuscript comments and edits. M.J and
47 L.A.K.D are fully funded by the University of Lincoln, School of Life Sciences. R.P.H is fully
48 funded by the University of Otago Doctoral Scholarship. R.P.H, D.P.D designed the study.
49 R.P.H performed data collection. R.P.H, M.J performed analysis and interpretation. R.P.H wrote
50 manuscript. D.P.D. supervised project. All authors (R.P.H, M.J, L.A.K.D, L.P.H, D.P.D)
51 discussed the results and implications and edited and commented on the manuscript at all stages.
52

53 **INTRODUCTION**

54 In recent decades, direct anthropogenic threats to terrestrial wildlife, primarily habitat
55 degradation and exploitation of natural resources, and indirect anthropogenic threats, primarily
56 climate change, have become increasingly prevalent, triggering declines and extinctions of
57 biodiversity (Dirzo et al. 2014; Trathan et al. 2014; Newbold et al. 2015; Urban 2015; Ceballos
58 et al. 2015). Concerns over accelerating wildlife loss have importantly been mitigated by the
59 establishment of protected areas (PAs) – geographical space designated and managed with the
60 long-term aim to sustainably conserve biodiversity, ecosystem services, and cultural values
61 (Brooks et al. 2004; Moilanen et al. 2009; Bertzky et al. 2012). They have become the most
62 widely implemented conservation action (Gillingham et al. 2015), and as of 2018, 14.9% of
63 global terrestrial areas (including inland waters) and 7.3% of the ocean are covered by some
64 form of legal protection (UNEP-WCMC et al. 2018). However, one of the central challenges
65 faced by the PA approach is the identification of vulnerable or irreplaceable organisms and
66 geographic regions that take into account the spatial and phylogenetic asymmetry of resident
67 biodiversity (e.g., endemism, species richness, taxonomic uniqueness) and population structure
68 (e.g., range size, population size, conservation status) (Reid 1998; Myers et al. 2000; Orme et al.
69 2005; Gaston et al. 2008). Here, we implement an exhaustive global-scale approach to assess the
70 overlap between PAs and the terrestrial breeding range (i.e., observed locations of individuals or
71 colonies of penguins on land) of penguins globally as a primary step towards an integrative
72 understanding of the efficiency of the current PA network in mitigating biodiversity declines.

73 Over the last six decades, PAs have generally been considered an effective conservation
74 approach. Their goal is to encourage ecological resilience by buffering against negative pressures
75 such as climate change, sustainably manage resources, and promote mutually beneficial human-

76 ecosystem interactions (refer to Gaston et al. 2008; Secretariat of the Convention on Biological
77 Diversity 2008). They have also been designated for the protection of species and populations in
78 biodiversity hotspots, including areas with high species richness or endemism (Myers et al. 2000;
79 Thiollay 2002; Brooks et al. 2006; Trathan et al. 2014). These biodiversity hotspots represent
80 areas that are environmentally suitable and able to sustain multiple species, making the area
81 valuable and worthy of protection. Protected areas also encompass areas and organisms which
82 have been prioritized for conservation actions based on ecological attributes that affect
83 persistence, such as range size, population size, and threats such as habitat degradation (Reid
84 1998; Boersma and Parrish 1999; Pichegru et al. 2010; Bertzky et al. 2012; Dirzo et al. 2014;
85 Trathan et al. 2014; Meiri et al. 2018). Range size and population size are commonly used to
86 estimate vulnerability, rarity, and extinction risk of a species and thus supports PA designation
87 and threat classification (Ferrière et al. 2004; Höglund 2009; Chevin et al. 2010; Pimm et al.
88 2014; Venter et al. 2014; Meiri et al. 2018). For example, species with small geographic ranges
89 generally have fewer individuals and lower genetic variation compared with species of larger
90 ranges (e.g., Galapagos penguins, *Spheniscus mendiculus*). As a result, these species might not
91 be able to maintain genetic diversity and spatial persistence if a portion of their range is altered,
92 which would ultimately maximize their priority as targets for conservation (Frankham 1996;
93 Gaston 2003; Höglund 2009; Charlesworth and Charlesworth 2010; Borboroglu and Boersma
94 2013; Meiri et al. 2018). Effective protection of these restricted populations is likely to have a
95 bigger impact on overall species survival than protecting one population in a wide ranging
96 species (Mace et al. 2008; Pimm et al. 2014).

97 While the majority of PAs are nationally designated and categorized using the International
98 Union for Conservation of Nature (IUCN) system based on management objectives and legal

99 status (IUCN 2001; Dudley 2008; see Table 1 in Online Resource), alternative international,
100 regional, and national classifications are also used (e.g., World Heritage sites). The purpose of
101 PA category systems is to first acknowledge a PA, its current conservation goals, and its
102 governing organisation and then to provide stakeholders with a framework for managing,
103 reporting, and monitoring management effectiveness into the future. Different category systems
104 call for different levels of protection, each with different management approaches (e.g., restricted
105 access, public use, resource exploitation). These categories provide a standardized outline for
106 defining PAs, but there is high variability between its actual management and the broad category
107 recommendations. The category system and associated data does not indicate if a PA was created
108 to protect a specific species or if that species merely occur within a PA that was established for
109 other management objectives. The system also does not quantify the effectiveness of the PA
110 designation on a specific species Nevertheless, any organism occupying area within a PA will be
111 subject to the effects of the PAs. Therefore, it is useful as a classification tool to group similar
112 PAs by overall management objective (e.g. protect a specific species, promote sustainable
113 ecosystem use) as a baseline for further studies on efficacy. Furthermore, when assessing the
114 irreplaceability of a species and its vulnerability to population decline, it is important to consider
115 how PA classification affects the overall coverage of the PA (Pressey et al. 1994; Pressey and
116 Taffs 2001; Dudley 2008).

117 A prime example of taxonomically unique organisms encompassing critical ecological
118 features considered in conservation decisions and PAs are penguins. Penguins, broadly regarded
119 as wildlife and cultural icons, are represented in public climate change and conservation
120 movements as focal targets for protection. These unique birds, comprising of 18 species globally,
121 are primarily restricted to the southern hemisphere (the only exception being *Spheniscus*

122 *mendiculus* from the Galápagos Archipelago). Approximately two-thirds of penguin species are
123 experiencing major population declines (Borboroglu and Boersma 2013; Boersma and Rebstock
124 2014; Trathan et al. 2014; Ropert-Coudert et al. 2019), which has resulted in ten species (>50%
125 of their global diversity) currently at risk of extinction, categorised as Vulnerable or Endangered
126 by the IUCN Red List (Ellis 1999; Boersma 2008; IUCN 2018). While some species have
127 widespread distributions and high population densities, others have highly restricted ranges
128 (Figure 1, Table 1), which likely increases their vulnerability to environmental change.

129 Penguins are critically dependent on and constrained to limited areas of land for breeding and
130 associated regions of the ocean for foraging (Borboroglu and Boersma 2013). Typically, foraging
131 ranges are influenced by prey availability and other factors, while breeding occurs annually at
132 the same location (Boersma 2008). Both habitats are vital for penguin survival and pose different
133 threats that they must contend with (Ropert-Coudert et al. 2019). Anthropogenic drivers of
134 population declines for penguins include climate change, habitat loss and degradation,
135 commercial fishing and bycatch, oil spills, pollution, and tourism, whereas environmental threats
136 include invasive species competition, El Niño events, and predation (Borboroglu et al. 2008;
137 Gandini et al. 2010; Pichegru et al. 2010; Borboroglu and Boersma 2013; Trathan et al. 2014;
138 Ropert-Coudert et al. 2019). While many threats operate in the marine environment (i.e.,
139 overfishing and bycatch), terrestrial threats such as unregulated tourism, over-exploitation, and
140 habitat modification have more direct negative effects on penguin productivity and survival
141 (Trathan et al. 2014).

142 This paper focuses on the overlap between terrestrial PAs and breeding sites of penguins for
143 several reasons. Firstly, although penguins spend a disproportionate amount of time in the ocean
144 rather than on land, breeding is only possible on land and during a specific time of the year.

145 Penguins are also philopatric, returning to the same nesting areas each year and even to the same
146 nest. Without successful breeding, recruitment of new individuals and population stability is
147 impossible. Having PAs include penguin nesting sites will protect them from the aforementioned
148 terrestrial threats, limiting these pressures and increasing their overall reproductive success.
149 Therefore it is critical to analyse current conservation methods impacting penguin colonies to
150 ensure continued survival. Secondly, differences in PA management, designation categories,
151 conservation objectives, and overall ecosystem structure on land versus in the ocean highlight the
152 necessity of assessing terrestrial PAs and marine PAs (MPAs) separately. Lastly, there are more
153 terrestrial PAs globally than MPAs and data on penguin range is of higher quality and quantity
154 than marine distribution data.

155 In this paper, we provide a global analysis of the patterns of terrestrial penguin biodiversity
156 distribution and their protection under the current PA network. Therefore, we aim to address
157 whether: *(i)* the terrestrial geographic distribution of global penguin species is sufficiently
158 protected by existing terrestrial PAs or overlaps with biodiversity hotspots classified by Myers et
159 al. (2000) (hereafter called Myers' hotspots), *(ii)* endangerment, as categorized by the IUCN Red
160 List, is predominant among penguin species for which lower proportions of their ranges are
161 covered by PAs, and *(iii)* whether terrestrial hotspots of penguin biodiversity (species richness
162 and endemism) fall within existing PAs. Our findings thus focus on quantifying the extent of
163 protection for penguins, which types of PAs occur within terrestrial sites used by penguins, and
164 if factors such as range or population size are correlated to the level of protection in order to
165 identify species and areas lacking protection and inform the future implementation and
166 management of these PAs.

167

168 **METHODS**

169 **Species occurrence data**

170 We compiled a global-scale dataset of the terrestrial geographic distribution of all 18 known
171 penguin species (family Spheniscidae). We first downloaded coordinate data points for all
172 Spheniscidae species from the open-access database Global Biodiversity Information Facility
173 (GBIF 2018). This data was filtered to exclude any points without a record date or dates prior to
174 1969 (points included last 50 years only to minimize inaccuracies). Data for each species was
175 assessed and compiled individually to limit exclusion errors. We excluded records with duplicate
176 and incorrectly formatted coordinates, records north of the Equator (except for *Spheniscus*
177 *mendiculus*, whose breeding sites extend slightly over the Equator), records without a valid
178 country code, and records classified as fossil/dead specimens or vagrants (only those recorded as
179 human observation were included). We also excluded spatial records whose locality description
180 was blank, included the keywords “pelagic”, “offshore”, “at sea”, “no information”, “marine”,
181 “sea”, “ocean”, or contained ocean names only (such as “Southern Ocean”). The majority of the
182 records in this dataset are colony/breeding site coordinates. However, it does include
183 observations of vagrant penguins sited outside of breeding areas, because there is no systematic
184 way to limit these observations further. The GBIF database does not distinguish between
185 vagrants and breeding sites. Therefore, we included colony data points from Borboroglu and
186 Boersma (2013), the most recent published compilation of colony records. The GBIF points were
187 checked against Borboroglu and Boersma (2013) range maps to identify incorrect or impossible
188 records, which were then excluded from the analysis. Finally, a mask was applied to crop all
189 points to global land surfaces. Therefore, our newly curated dataset of global penguins will,
190 additionally, contribute a new resource for future penguin and bird research.

191 Data on penguin population size and IUCN Red List conservation status (hereafter
192 conservation status) were obtained from Borboroglu and Boersma (2013) and the IUCN Red List
193 (2014, 2018) as a compilation of published and unpublished data from many sources. While
194 population sizes are naturally variable, these population estimates are the most reliable to date
195 based on satellite imaging and/or long-term data collection.

196

197 **Protected Areas data**

198 We collated the spatial data for PAs from the World Database on Protected Areas
199 (WDPA; www.protectedplanet.com) . This dataset includes PAs classified by the IUCN
200 Protected Areas Categories System (henceforth referred to as IUCN PAs), the world’s most
201 inclusive and globally accepted prioritization scheme for nationally managed PAs (see Dudley
202 (2008) for category descriptions). Due to the variability of protection within and between each
203 IUCN category, we grouped all categories as “IUCN PAs”, as the intent was to quantify
204 protection as a whole. Category-specific examination of protection was out of the scope of this
205 analysis. In addition to IUCN PAs, the dataset differentiates PAs that are nationally protected but
206 not categorized (“Not Reported”, NR) and international PAs categorised as “Not Applicable”
207 (NA). Not reported and not applicable PAs were grouped as “Not Categorized” (NC) in our
208 analyses.

209 The PA distribution map was derived using the 2018 WDPA shapefiles and
210 corresponding attribute tables. Due to the ambiguity of particular records, all point records, those
211 with null latitude and longitude, those listed as “marine”, polygon records with no area
212 information, and those north of the Equator were excluded from these analyses. Some areas are

213 classified using both IUCN and other category systems simultaneously, so overlap between
214 different designation types was removed when determining the total protection for each species.

215 In addition to the above protected areas, we included Antarctic Specially Protected
216 Areas (ASPA) in our analyses (Terauds 2017, 2018). Similar to IUCN Ia or II PAs, ASPAs
217 protect mammals and seabirds (and other associated ecosystem values) by primarily limiting
218 human interference (Southwell et al. 2017). These areas are recognized by the Protocol on
219 Environmental Protection to the Antarctic Treaty (United Nations 1991) and managed by
220 respective international governments depending on location. Antarctica SPAs are the only set of
221 PAs in Antarctica that can be considered equivalent to IUCN PAs in terms of classification
222 requirements and management objectives (Coetzee et al. 2017). The ASPAs were grouped as
223 “ASPAs” in our analyses.

224

225 **Species distribution analyses**

226 In order to determine spatial overlap between penguin ranges and PAs, we first calculated range
227 size for each individual species. Due to the fragmented distribution of penguin breeding sites, the
228 area that penguins occupy (‘area of occupancy’, AOO) was calculated. The circular buffer
229 method presented in Hernández and Navarro (2007), Rivers et al. (2010), and Breiner and
230 Bergamini (2018) was modified to create ranges based upon the distance between points for each
231 species. A distance matrix between all points determined the mean value of the minimum
232 distance between points. Using this mean value as the radius, each point was buffered by this
233 distance. Overlapping circles were merged. Although these AOO ranges can include areas not
234 currently occupied by breeding penguins (e.g., area between colonies, geographic features), this

235 method best represents unrecorded colonies, potential future colonies, and areas used by
236 penguins for non-breeding purposes.

237 Next, we masked and clipped the PAs using each species' AOO to quantify the overlap of
238 each PA type (IUCN, NC, and ASPA) within all species ranges. Each type of PA was classified
239 and area was calculated and summed. Overlap between PA type was determined by dissolving
240 all PAs and calculating the difference. We performed all analyses using QGIS 3.2.1 Bonn (QGIS
241 2018).

242

243 **Species richness and endemism analyses**

244 After creating a GIS grid shapefile of global penguin distribution with the southern
245 hemisphere (3°N to 90°S) as a mask and a cell size of 1 degree (~111.12 km at the Equator)
246 projected using South Pole Lambert Azimuthal Equal Area, we constructed the distribution of
247 species richness of penguins (i.e., number of penguin species contained per single grid cell)
248 using Spatial Analysis in Macroecology (SAM) software, available at
249 <http://www.ecoevol.ufg.br/sam> (Rangel et al. 2010). We considered as hotspots of penguins
250 those grid cells in which at least four breeding species have been recorded, which represents the
251 richest 2.5% cells (Orme et al. 2005). We then determined the overlap between worldwide
252 biodiversity hotspots, as established by Myers et al. (2000), and AOO to quantify the extent to
253 which a species range within a biodiversity hotspot is protected by IUCN or NC PAs. Myers et
254 al. (2000) terrestrial biodiversity hotspots (1) “contain at least 0.5% or 1,500 of the world's
255 300,000 plant species as endemics”, (2) contain a high percentage of endemic vertebrate species
256 (mammals, birds, reptiles, and amphibians), and/or (3) have lost 70% or more of its primary

257 vegetation (Myers et al. 2000). We performed all biodiversity hotspot analyses using QGIS 3.2.1
258 Bonn (QGIS 2018).

259 Additionally, we investigated whether hotspots of penguin endemism are associated with
260 PAs. A species is endemic if it occurs only in a defined area (for penguins, endemic species are
261 usually range restricted to one island or one country). An area has high endemism if it contains
262 many range-restricted species. To determine global endemism, we first calculated the Corrected
263 Weighted Endemism (CWE) for each grid cell. CWE represents the weighted endemism (for
264 each grid cell, the sum of the reciprocal of the total number of grid cells that each species occurs
265 in) divided by species richness (the total number of species in that cell) to correct for species
266 richness correlation. In other words, CWE emphasizes areas that have species with restricted
267 distribution rather than areas with high species richness (Crisp et al. 2001). This index ranges
268 from 0.0 to 1.0, corresponding to having 0-100% of the species occurring within that cell having
269 a restricted range to that cell (Laffan and Crisp 2003). We performed all CWE analyses using the
270 Analysis and Spatial Statistics tools and SDMToolbox (CWE) of ArcGIS 10.6.1 (Brown 2014;
271 ESRI 2018).

272

273 **Quantitative analyses**

274 To address whether existing PAs are related to specific biodiversity factors, we first employed
275 Spearman Rank Correlation tests to quantify the relationship between population and range size
276 between different types of PAs. Kruskal-Wallis Rank Sum tests were performed to determine
277 whether protection levels (percentage of area covered by an IUCN, NC, of ASPA PA for each
278 species) differed among conservation statuses. We also used a Kruskal-Wallis test to evaluate

279 whether there is an association between range size/population size and conservation status. All
280 statistics were implemented in R version 3.1.2 (R Development Core Team 2019).

281

282 **RESULTS**

283 **Global species distributions**

284 Penguin species are widely distributed across four continents and occupy a global terrestrial area
285 of 629,887 km² (Figure 1, Table 1). Geographic range and population sizes vary considerably
286 across species but are not normally distributed (Kolmogorov–Smirnov $p < 0.01$; Table 2, Online
287 Resource Figure 1). There is a skewed tendency for range sizes to be small (Online Resource
288 Figure 1), with the smallest range being only 0.81 km² (*Eudyptes robustus*) and the largest being
289 135,395 km² (*Aptenodytes forsteri*). Thirteen species have ranges between 0.81 km² to 40,000
290 km². Individual species ranges can span a large portion of the Antarctic coast (*Pygoscelis*
291 *adeliae*) while others are restricted to a small island (*E. robustus*).

292

293 **Protected area coverage**

294 All penguin species are protected to some degree (Table 1, Figure 2; see Figures 2 and 3 in
295 Online Resource for maps of PAs) by at least one PA (Online Resource Table 2). Total
296 protection based on species range covered by any type of PA varies from 0.16% (*Aptenodytes*
297 *forsteri*) to 100% of a species range. For seven species, total protection is greater than 50%, and
298 three of these seven species are fully protected by IUCN and NC PAs (*E. robustus*, *Eudyptes*
299 *schlegeli*, and *S. mendiculus*; Table 1). For fourteen species, IUCN protection is less than 40%,
300 while NC PAs cover 14 species by less than 31% (Table 1, Figure 2). All Antarctic species are
301 covered to some degree by an ASPA PA, albeit a very small percentage of their range.

302 Additionally, some areas are protected simultaneously by IUCN and NC (Online Resource Table
303 3). For example, *Eudyptes chrysocome* is 22.83%, 16.95%, and 0.07% by the IUCN, NC, and
304 ASPA, respectively. However, the total combined protection is 28.01%, indicating an overlap of
305 15.54%.

306 Protected area coverage is non-normally distributed across species. Spearman's rank tests
307 revealed that there is a slightly significant relationship between total, IUCN, and ASPA PA
308 coverage and range size (Table 2). Population size and conservation status have non-significant
309 relationships with PA coverage, except for a significant correlation between ASPA protection
310 and population (Table 2).

311 Additionally, conservation status is not significantly influenced by range size (Kruskal-
312 Wallis chi-squared = 4.44, df = 3, p value = 0.22) or population (Kruskal-Wallis chi-squared =
313 7.29, df = 3, p value = 0.06). However, Endangered penguins have smaller range sizes and
314 population sizes (Online Resource Figure 4). Vulnerable and Endangered species are, in total,
315 more protected than Least Concern and Near Threatened species. Vulnerable species are most
316 protected by IUCN PAs compared with all other conservation statuses, while NC protection
317 remains similar between status levels. Compared with IUCN PAs, NC PAs cover slightly more
318 of total, global penguin range.

319

320 **Hotspots of species richness and endemism**

321 Our analyses identify seven global hotspots of penguin biodiversity where four or five penguin
322 species breed, concentrated on the sub-Antarctic islands, southern tip of South America, and
323 Antarctic Peninsula (Figure 3a-c, Online Resource Table 4). All hotspots are protected to some
324 degree, and three are fully protected by IUCN and NC PAs. Furthermore, Macquarie Island is the

325 only penguin hotspot that is simultaneously a Myers' hotspot. Approximately 6.1% of total
326 penguin range is within a Myers' hotspot, and 10.4% of that area is protected. Out of the 13
327 species whose ranges are within a Myers' hotspot, six overlap with a hotspot by more than 60%.
328 The remaining five species are entirely excluded from a Myers' hotspot. Additionally, range size
329 and population size are not significantly related with Myers' hotspot overlap and protection
330 (Table 2).

331 Globally, CWE ranges from 0.0 to 0.51 (Figure 3d). Snares Island has the highest CWE
332 of 0.51. Macquarie, Amsterdam, and St. Paul Island have a CWE greater than 0.20, while South
333 Africa, Galapagos Islands, and parts of New Zealand have CWE values ranging from 0.08 to
334 0.11 (Figure 3d). In general, penguins have a relatively low CWE.

335

336 **DISCUSSION**

337 Our study provides the first comprehensive global assessment investigating the
338 relationships between the terrestrial distribution of the world's penguin species and existing PAs.
339 Only 16.80% of the total global penguin range is protected by IUCN, NC, and ASPA PAs
340 combined, and coverage is extremely variable and unpredictable among species, with no
341 standardisation based on conservation status or population size. In addition, penguins generally
342 breed in isolated and endemic populations (Borboroglu and Boersma 2013), resulting in few
343 hotspot areas. It is more common for PAs to be implemented to protect hotspots of biodiversity
344 than to protect isolated populations of one species. Lack of protection is likely to increase species
345 risk of decline under environmental or population changes (Isik 2011; Pimm et al. 2014).
346 Previous analyses of the irreplaceability and vulnerability of penguins (Borboroglu and Boersma
347 2013; Trathan et al. 2014; Ropert-Coudert et al. 2019), combined with our findings, highlight our

348 concerns about the generality and inadequate coverage of global PAs for penguins and support
349 our advocacy for improved prioritization of sites and species. In a rapidly changing world, the
350 identification of such biodiversity patterns will allow evidence-based predictions about the
351 magnitude and impact of anthropogenic threats on species, to potentially influence decisions
352 about environmental management. Therefore, our study closes a major gap in the knowledge of
353 these global interactions experienced by penguins, one of the most charismatic groups of
354 vertebrates on Earth.

355

356 **Protection efficiency: PAs, hotspots, and ‘coldspots’**

357 PAs ensure the persistence of nature by primarily limiting the effects of humans on species and
358 habitats. However, simultaneous management by more than one organization or categorization as
359 different types of PAs highlights the overall mismanagement and non-collaborative designation
360 processes. For example, the Galápagos Islands are classified as a World Heritage site, a
361 UNESCO-MAB Biosphere Reserve, a Ramsar site, and an IUCN national park, each of which
362 has different prioritization strategies, goals, and management objectives, resulting in conflicting
363 category rankings and overall protection methods. In theory, a site with multiple protection
364 designations (typically representing additional organizations and stakeholders) could be
365 beneficial for increasing effort, sharing responsibility, or multiplying the types of conservation
366 efforts or organisms protected. It is typical for overlap to occur between national designations
367 and international designations, as seen on the Galapagos Islands. This multiple classification
368 emphasizes the ecological importance of these type of sites on a more local and global scale
369 simultaneously (Deguignet et al. 2017). However, conflicts such as uneven and ineffective use of
370 resources or logistical problems can arise that detracts from the effectiveness of management

371 efforts (Ioja et al. 2010; Deguignet et al. 2017). Understanding the overall coverage of PAs and
372 the overlap between classifications can be used to assess PA effectiveness and the disparity (both
373 positive and negative) between classification and management now and in the future.

374 Areas and species can also be protected at national scale but not be considered within the
375 WDPA database. For example, the Falkland Islands are governmentally protected but according
376 to Protected Planet, only 61 km² of land area is IUCN protected (IUCN and UNEP 2018). A
377 subsequent analysis including and differentiating areas that are locally or nationally protected
378 under different schemes (along with an analysis of effectiveness) will support the global-scale
379 overview presented here.

380 Conservation focuses on protecting areas that support the largest number of species
381 having the smallest, most threatened populations (Eken et al. 2004; Brooks et al. 2006; Akçakaya
382 et al. 2007; Dirzo et al. 2014). This is especially true for penguins - their populations are
383 generally small with relatively small breeding areas confined to coastal zones. We identified
384 areas of high penguin endemism (CWE, Figure 3d) that contain species of small ranges which
385 inhabit few other areas. This measure also quantifies areas that have both high endemism and
386 species richness. Loss of even a few populations could be potentially detrimental to entire
387 species as a whole. Additionally, the abundance of areas supporting single species of penguins
388 (as opposed to only seven hotspots of four or five species) and the protection of these ‘coldspots’
389 may be preferable if that species is endemic (Orme et al. 2005) or declining in population
390 (Geldmann et al. 2013). For penguins, rarity is a critical parameter to take into account when
391 developing conservation planning. Rarity frequently translates into not only naturally small
392 populations or range sizes (Lennon et al. 2003) but a combination of both (Mace et al. 2008).
393 Any significant population loss could result in the eventual extinction of the whole species

394 (Borboroglu and Boersma 2013; Ropert-Coudert et al. 2019). The contradiction between species
395 richness and endemism makes it difficult to determine which penguin species and areas to
396 protect in order to simultaneously maintain genetic, species, and ecosystem diversity.

397

398 **Future protection of penguins**

399 The geographic data for penguin terrestrial areas used within this study is comprehensive
400 and inclusive of known breeding areas. However, due to the limitations of using the GBIF
401 database (including the ambiguity of local, vagrant, or unusual occurrences), areas may have
402 been included in these analyses that are outside of normal breeding areas. Arguably, while this
403 may inflate the geographic range for some species, the fact that their population persistence
404 depends on these areas is a critical feature that should not be ignored. As a result of progressing
405 and increasingly destructive anthropogenic environmental change, these areas may prove key for
406 the occupation of penguins, which may lead them to be considered for protection in the near
407 future.

408 As a whole, sites for conservation should be prioritized following the identification of
409 vulnerable and irreplaceable ecosystems and species. However, in practice, prioritization tends to
410 be (primarily) geographically or taxonomically designated, with no clear systematic connection
411 (Rodrigues et al. 2004; Bertzky et al. 2012). Furthermore, protection is focused either proactively
412 or reactively, depending on management objectives (Ropert-Coudert et al. 2019). An area can be
413 prioritized in order to prevent future biodiversity loss or repair loss that has already occurred.
414 This is the case for penguins. Existing PAs often do not include species for which conservation is
415 needed the most (Eken et al. 2004). Due to the majority of penguin species being highly
416 threatened, having small ranges and population sizes, or being endemic to small regions, we

417 propose a combination of both proactive and reactive conservation strategies (similarly
418 suggested in Ropert-Coudert et al. (2019)). Additionally, the effectiveness of protection should
419 be considered for species experiencing threats or large population declines, in addition to
420 biodiversity hotspots where multiple penguin species breed (specifically the Falkland Islands,
421 Tierra del Fuego, and Southern New Zealand).

422 Finally, additional assessments of the effectiveness of marine PAs at protecting penguin
423 marine foraging areas and prey are required for the global conservation of all areas vital to
424 penguin survival. Penguins are primarily marine animals and spend most of their time at sea.
425 There is currently no assessment of global-scale marine protection for penguins, although there is
426 ongoing research regarding the threats faced while foraging (Ropert-Coudert et al. 2019). This
427 critical habitat should be equally, if not more, protected than their breeding sites.

428

429 **Conclusion**

430 Over the past three decades, the increasing global biodiversity crises arising as a result of human
431 activities have promoted exponential growth in the development of ecologically- and
432 evolutionary-based conservation approaches (Ferrière et al. 2004; Höglund 2009). These
433 methods rely primarily on PAs to maintain and increase biodiversity and population by
434 promoting processes such as migration and proliferation (e.g., improving habitat connectivity,
435 reducing fragmentation, limiting poaching) (Thomas and Gillingham 2015). However, they are
436 generally failing to protect key species (Gaston 2003). From our findings, we suggest future
437 research should focus on determining those key penguin species that require more protection
438 based upon their rarity. We also suggest protection requirements and conservation needs for each
439 individual species and population sustainability within each PA should be determined.

440 Management and policy should be assessed to distinguish between effective and non-effective
441 PAs, so that future evidence-based policy, including the global promotion of the IUCN category
442 system, can be implemented.

443 **REFERENCES**

- 444 Akçakaya HR, Mills G, Doncaster CP (2007) The role of metapopulations in conservation. In:
445 Macdonald DW, Service K (eds) Key topics in conservation biology. Blackwell Publishing,
446 Oxford, UK, pp 64–84
- 447 Bertzky B, Corrigan C, Kemsey J, et al (2012) Protected planet report 2012: Tracking progress
448 towards global targets for protected areas. UNEP-WCMC, Cambridge, UK
- 449 Boersma PD (2008) Penguins as marine sentinels. *Bioscience* 58:597–607
- 450 Boersma PD, Parrish JK (1999) Limiting abuse: marine protected areas, a limited solution. *Ecol*
451 *Econ* 31:287–304. doi: 10.1016/S0921-8009(99)00085-3
- 452 Boersma PD, Rebstock GA (2014) Climate change increases reproductive failure in Magellanic
453 penguins. *PLoS One* 9:. doi: 10.1371/
- 454 Borboroglu PG, Boersma PD (2013) Penguins: natural history and conservation. University of
455 Washington Press
- 456 Borboroglu PG, Boersma PD, Reyes L, Skewgar E (2008) Petroleum, pollution, and penguins:
457 marine conservation tools to reduce the problem. In: Hofer TN (ed) *Marine Pollution: New*
458 *Research*. Nova Science Publishers Inc., New York, USA, pp 339–356
- 459 Breiner FT, Bergamini A (2018) Improving the estimation of area of occupancy for IUCN Red
460 List assessments by using a circular buffer approach. *Biodivers Conserv* 27:2443–2448. doi:
461 10.1007/s10531-018-1555-5
- 462 Brooks TM, Bakarr MI, Boucher TIM, et al (2004) Coverage provided by the global protected
463 area system: Is it enough? *Bioscience* 54:1081–1091. doi: 10.1641/0006-
464 3568(2004)054[1081:CPBTGP]2.0.CO;2
- 465 Brooks TM, Mittermeier RA, da Fonseca GAB, et al (2006) Global biodiversity conservation

466 priorities. *Science* (80-) 313:58–61. doi: 10.1126/science.1127609

467 Brown JL (2014) SDMtoolbox: a python-based GIS toolkit for landscape genetic, biogeographic,
468 and species distribution model analyses. *Methods Ecol Evol* 5:694–700

469 Ceballos G, Ehrlich PR, Barnosky AD, et al (2015) Accelerated modern human-induced species
470 losses: Entering the sixth mass extinction. *Sci Adv* 1:e1400253–e1400253. doi:
471 10.1126/sciadv.1400253

472 Charlesworth B, Charlesworth D (2010) *Elements of evolutionary genetics*. Roberts & Co.,
473 Greenwood Village, Colorado

474 Chevin L-M, Lande R, Mace GM (2010) Adaptation, plasticity, and extinction in a changing
475 environment: towards a predictive theory. *PLoS Biol* 8:e1000357. doi:
476 10.1371/journal.pbio.1000357

477 Coetsee BWT, Convey P, Chown SL (2017) Expanding the Protected Area Network in
478 Antarctica is Urgent and Readily Achievable. *Conserv Lett* 10:670–680. doi:
479 10.1111/conl.12342

480 Crisp MD, Laffan S, Linder HP, Monro A (2001) Endemism in the Australian flora. *J Biogeogr*
481 28:183–198

482 Deguignet M, Arnell A, Juffe-Bignoli D, et al (2017) Measuring the extent of overlaps in
483 protected area designations. *PLoS One* 12:e0188681. doi: 10.1371/journal.pone.0188681

484 Dirzo R, Young HS, Galetti M, et al (2014) Defaunation in the Anthropocene. *Science* (80-)
485 345:401–406. doi: 10.1126/science.1251817

486 Dudley N (2008) *Guidelines for applying protected area management categories*. IUCN, Gland,
487 Switzerland

488 Eken G, Bennun L, Brooks TM, et al (2004) Key biodiversity areas as site conservation targets.

489 Bioscience 54:1110–1118. doi: 10.1641/0006-3568(2004)054[1110:kbaasc]2.0.co;2

490 Ellis S (1999) The penguin conservation assessment and management plan: a description of the
491 process. *Mar Ornithol* 27:163–169

492 ESRI (2018) ArcGIS Desktop: 10.2.2. 10.6.1

493 Ferrière R, Dieckmann U, Couvet D (2004) *Evolutionary conservation biology*. Cambridge
494 University Press

495 Frankham R (1996) Relationship of genetic variation to population size in wildlife. *Conserv Biol*
496 10:1500–1508. doi: 10.1046/j.1523-1739.1996.10061500.x

497 Gandini P, Frere E, Boersma PD (2010) Status and conservation of Magellanic Penguins
498 *Spheniscus magellanicus* in Patagonia, Argentina. *Bird Conserv Int* 6:307–316. doi:
499 10.1017/s0959270900001787

500 Gaston KJ (2003) *The structure and dynamics of geographic ranges*. Oxford University Press,
501 Oxford, UK

502 Gaston KJ, Jackson SF, Cantú-Salazar L, Cruz-Piñón G (2008) The ecological performance of
503 protected areas. *Annu Rev Ecol Evol Syst* 39:93–113. doi:
504 10.1146/annurev.ecolsys.39.110707.173529

505 GBIF (2018) *The Global Biodiversity Information Facility Backbone Taxonomy*.
506 <http://www.gbif.org/species>

507 Geldmann J, Barnes M, Coad L, et al (2013) Effectiveness of terrestrial protected areas in
508 reducing habitat loss and population declines. *Biol Conserv* 161:230–238. doi:
509 10.1016/J.BIOCON.2013.02.018

510 Gillingham PK, Bradbury RB, Roy DB, et al (2015) The effectiveness of protected areas in the
511 conservation of species with changing geographical ranges. *Biol J Linn Soc*

512 Hernández HM, Navarro M (2007) A new method to estimate areas of occupancy using
513 herbarium data. *Biodivers Conserv* 16:2457–2470. doi: 10.1007/s10531-006-9134-6

514 Höglund J (2009) *Evolutionary conservation genetics*. Oxford University Press

515 Iojă CI, Pătroescu M, Rozyłowicz L, et al (2010) The efficacy of Romania’s protected areas
516 network in conserving biodiversity. *Biol Conserv* 143:2468–2476. doi:
517 10.1016/J.BIOCON.2010.06.013

518 Isik K (2011) Rare and endemic species: why are they prone to extinction. *Turkish J Bot*
519 35:411–417. doi: 10.3906/bot-1012-90

520 IUCN (2001) *IUCN red list categories and criteria, version 3.1*. IUCN Species Survival
521 Commission, Gland, Switzerland and Cambridge, UK

522 IUCN (2018) *The IUCN Red List of Threatened Species*. Version 2018-1.
523 <http://www.iucnredlist.org>

524 IUCN, UNEP (2018) *The World Database on Protected Areas (WDPA)*.
525 <http://www.protectedplanet.net>

526 Laffan SW, Crisp MD (2003) Assessing endemism at multiple spatial scales, with an example
527 from the Australian vascular flora. *J Biogeogr* 30:511–520. doi: 10.1046/j.1365-
528 2699.2003.00875.x

529 Lennon JJ, Koleff P, Greenwood JJD, Gaston KJ (2003) Contribution of rarity and commonness
530 to patterns of species richness. *Ecol Lett* 7:81–87. doi: 10.1046/j.1461-0248.2004.00548.x

531 Mace GM, Collar NJ, Gaston KJ, et al (2008) Quantification of Extinction Risk: IUCN’s System
532 for Classifying Threatened Species. *Conserv Biol* 22:1424–1442. doi: 10.1111/j.1523-
533 1739.2008.01044.x

534 Meiri S, Bauer AM, Allison A, et al (2018) Extinct, obscure or imaginary: The lizard species

535 with the smallest ranges. *Divers Distrib* 24:262–273. doi: 10.1111/ddi.12678

536 Moilanen A, Wilson KA, Possingham HP (2009) *Spatial conservation prioritization : quantitative*
537 *methods and computational tools*. Oxford University Press

538 Myers N, Mittermeier RA, Mittermeier CG, et al (2000) Biodiversity hotspots for conservation
539 priorities. *Nature* 403:853–858. doi: 10.1038/35002501

540 Newbold T, Hudson LN, Hill SLL, et al (2015) Global effects of land use on local terrestrial
541 biodiversity. *Nature* 520:45–50. doi: 10.1038/nature14324

542 Orme CDL, Davies RG, Burgess M, et al (2005) Global hotspots of species richness are not
543 congruent with endemism or threat. *Nature* 436:1016–1019. doi: 10.1038/nature03850

544 Pichegru L, Grémillet D, Crawford RJM, Ryan PG (2010) Marine no-take zone rapidly benefits
545 endangered penguin. *Biol Lett* 6:498–501. doi: 10.1098/rsbl.2009.0913

546 Pimm SL, Jenkins CN, Abell R, et al (2014) The biodiversity of species and their rates of
547 extinction, distribution, and protection. *Science* (80-) 344:. doi: 10.1126/science.1246752

548 Pressey RL, Johnson IR, Wilson PD (1994) Shades of irreplaceability: towards a measure of the
549 contribution of sites to a reservation goal. *Biodivers Conserv* 3:242–262

550 Pressey RL, Taffs KH (2001) Scheduling conservation action in production landscapes: priority
551 areas in western New South Wales defined by irreplaceability and vulnerability to
552 vegetation loss. *Biol Conserv* 100:355–376

553 QGIS (2018) QGIS 3.2.1

554 R Development Core Team (2019) R: A language and environment for statistical computing

555 Rangel TF, Diniz-Filho JAF, Bini LM (2010) SAM: a comprehensive application for spatial
556 analysis in macroecology. *Ecography (Cop)* 33:46–50

557 Reid W V (1998) Biodiversity hotspots. *Trends Ecol Evol* 13:275–280

558 Rivers MC, Bachman SP, Meagher TR, et al (2010) Subpopulations, locations and
559 fragmentation: Applying IUCN red list criteria to herbarium specimen data. *Biodivers*
560 *Conserv* 19:2071–2085. doi: 10.1007/s10531-010-9826-9

561 Rodrigues ASL, Akcakaya HR, Andelman SJ, et al (2004) Global gap analysis: priority regions
562 for expanding the global protected-area network. *Bioscience* 54:1092–1100

563 Ropert-Coudert Y, Chiaradia A, Ainley D, et al (2019) Happy Feet in a hostile world? The future
564 of penguins depends on proactive management of current and predictable threats. *Front Mar*
565 *Sci*

566 Secretariat of the Convention on Biological Diversity (2008) *Protected Areas in Today's World:*
567 *Their Values and Benefits for the Welfare of the Planet.* Montreal

568 Southwell C, Emmerson L, Takahashi A, et al (2017) Large-scale population assessment informs
569 conservation management for seabirds in Antarctica and the Southern Ocean: A case study
570 of Adélie penguins. *Glob Ecol Conserv* 9:104–115. doi: 10.1016/J.GECCO.2016.12.004

571 Terauds A (2017) An update to the Antarctic Specially Protected Areas (ASPAs). In: *Aust.*
572 *Antarct. Data Cent.*
573 [https://data.aad.gov.au/metadata/records/AAS_4296_Antarctic_Specially_Protected_Areas_](https://data.aad.gov.au/metadata/records/AAS_4296_Antarctic_Specially_Protected_Areas_v2)
574 [v2](https://data.aad.gov.au/metadata/records/AAS_4296_Antarctic_Specially_Protected_Areas_v2). Accessed 11 Mar 2019

575 Terauds A (2018) Antarctic Specially Protected Areas (Points and Polygons). In: *Aust. Antarct.*
576 *Data Cent.* https://data.aad.gov.au/metadata/records/AAS_4296_Updated_ASPAs_2018.
577 Accessed 11 Mar 2019

578 Thiollay J-M (2002) Bird diversity and selection of protected areas in a large neotropical forest
579 tract. *Biodivers Conserv* 11:1377–1395. doi: 10.1023/A:1016269813160

580 Thomas CD, Gillingham PK (2015) The performance of Protected Areas for biodiversity under

581 climate change. *Biol J Linn Soc*

582 Trathan PN, Borboroglu PG, Boersma D, et al (2014) Pollution, habitat loss, fishing, and climate
583 change as critical threats to penguins. *Conserv Biol* 1–11. doi: 10.1111/cobi.12349

584 UNEP-WCMC, IUCN, NGS (2018) Protected Planet Report 2018. Cambridge, UK; Gland,
585 Switzerland; Washington, DC, USA

586 United Nations (1991) Protocol on Environmental Protection to the Antarctic Treaty. In:
587 Antarctic Treaty Consultative Meeting XVI, 7-18 Oct 1991. Bonn, Germany

588 Urban MC (2015) Accelerating extinction risk from climate change. *Science* (80-) 348:571–573.
589 doi: 10.1126/science.aaa4984

590 Venter O, Fuller RA, Segan DB, et al (2014) Targeting global protected area expansion for
591 imperiled biodiversity. *PLoS Biol* 12:e1001891. doi: 10.1371/journal.pbio.1001891

592

593 **FIGURE CAPTIONS**

594 **Fig. 1** Map of penguin nest site distribution in **(a)** Antarctica, **(b)** Australia, New Zealand, and
595 surrounding sub-Antarctic islands, **(c)** South America, and **(d)** South Africa and surrounding sub-
596 Antarctic islands. Not shown are Galapagos penguins nesting only on the Galapagos Islands.
597 Panel **a** is projected using South Pole Lambert Azimuthal Equal Area. Panels **b**, **c**, and **d** are
598 projected using the World Geodetic System 1984. Basemap from Natural Earth
599 (<http://www.naturalearthdata.com>).

600 **Fig. 2** Percent of occupancy area coverage by IUCN Protected Areas Categories System 1b-VI
601 (IUCN, black bar) and IUCN “Not Reported” and “Not Categorized” (NC, grey bar) protected
602 areas for all penguin species. Total, non-overlapping protected area percent coverage is indicated
603 by the black horizontal line. Antarctic and sub-Antarctic species indicated by *. Species are
604 categorized by IUCN Red List conservation status.

605 **Fig. 3** Map of **(a)** global penguin species richness, sub-sectioned by regions including **(b)**
606 southern South America and the Antarctic Peninsula and **(c)** Australia and New Zealand. Species
607 richness legend applicable for panels a-c, and colours represent the number of species per 1
608 degree grid cell. Map of **(d)** global penguin corrected weighted endemism ranges from 0 to 0.51
609 (1 being the highest possible) per 1 degree grid cell. All maps are projected using the World
610 Geodetic System 1984. Basemap from Natural Earth (<http://www.naturalearthdata.com>).

611


612 **Table 1** Summary table of all penguin species, including IUCN Red List conservation status,
613 population size from Borboroglu and Boersma (2013), and area of occupancy (range size).
614 Included is percent of occupancy area coverage by IUCN Protected Areas Categories System 1b-
615 VI (IUCN), IUCN “Not Reported” and “Not Categorized” Protected Areas (NC), and Antarctic

616 Specially Protected Areas (ASPAs), Myers' biodiversity hotspots percent coverage of each
617 species range, and total protection of those biodiversity areas. Refer to Table 3 in Online
618 Resource for complete PA coverage data.

619 **Table 2** Summary of population and range size Spearman Rank tests and IUCN Red List
620 conservation status Kruskal-Wallis test ($df = 3$, denoted with †) for protected area coverage by
621 IUCN Protected Areas Categories System 1b-VI (IUCN), IUCN "Not Reported" and "Not
622 Categorized" Protected Areas (NC), and Antarctic Specially Protected Areas (ASPAs). Same tests
623 done for Myers' biodiversity hotspots. Coverage represents the percent of penguin ranges
624 covered by a biodiversity hotspot, and Protection represents the total percent protection of these
625 hotspots.


626

627



628


629 Fig. 1 Map of penguin nest site distribution in (a) Antarctica, (b) Australia, New Zealand, and surrounding sub-Antarctic islands, (c) South America,
 630 and (d) South Africa and surrounding sub-Antarctic islands. Not shown are Galapagos penguins nesting only on the Galapagos Islands. Panel a is
 631 projected using South Pole Lambert Azimuthal Equal Area. Panels b, c, and d are projected using the World Geodetic System 1984. Basemap from
 632 Natural Earth (<http://www.naturalearthdata.com>).



633

634 Fig. 2 Percent of occupancy area coverage by IUCN Protected Areas Categories System 1b-VI
 635 (IUCN, black bar) and IUCN “Not Reported” and “Not Categorized” (NC, grey bar) protected
 636 areas for all penguin species. Total, non-overlapping protected area percent coverage is indicated
 637 by the black horizontal line. Antarctic and sub-Antarctic species indicated by *. Species are
 638 categorized by IUCN Red List conservation status.

639



640
 641
 642
 643
 644
 645
 646
 647
 648

Fig. 3 Map of (a) global penguin species richness, sub-sectioned by regions including (b) southern South America and the Antarctic Peninsula and (c) Australia and New Zealand. Species richness legend applicable for panels a-c, and colours represent the number of species per 1 degree grid cell. Map of (d) global penguin corrected weighted endemism ranges from 0 to 0.51 (1 being the highest possible) per 1 degree grid cell. All maps are projected using the World Geodetic System 1984. Basemap from Natural Earth (<http://www.naturalearthdata.com>).

649 Table 1

Species	Common Name	Status *	Population Size	Occurrence Area (km ²)	Protection Level (%)				Biodiversity Hotspot (%)	
					IUCN	NC ‡	ASPA ‡	Total	Coverage	Protection
<i>Aptenodytes forsteri</i>	emperor	NT	595000	135395.63	0.11	0.00	0.09	0.16	0.00	0.00
<i>Aptenodytes patagonicus</i>	King	LC	3200000	12855.37	27.06	27.95	0.03	30.18	0.69	100
<i>Eudyptes chrysocome</i>	southern rockhopper	VU	2460000	131371.72	22.83	16.95	0.07	28.01	0.62	100
<i>Eudyptes chrysolophus</i>	macaroni	VU	12600000	92703.12	8.73	12.48	0.16	18.47	0.10	100
<i>Eudyptes moseleyi</i>	Northern rockhopper	EN	530000	238.36	25.34	58.75	n/a	58.75	0.00	0.00
<i>Eudyptes pachyrhynchus</i>	Fiordland-crested	VU	6000	782.70	97.21	72.83	n/a	97.21	100	97.21
<i>Eudyptes robustus</i>	Snares	VU	62000	0.81	100	0.00	n/a	100	0.00	0.00
<i>Eudyptes schlegeli</i>	royal	NT	1700000	123.05	100	0.00	n/a	100	100	100
<i>Eudyptes sclateri</i>	Erect-crested	EN	140000	21.50	98.23	0.00	n/a	98.23	96.93	100
<i>Eudyptula minor</i>	little	LC	469760	12455.67	36.97	14.96	n/a	39.17	24.49	29.83
<i>Megadyptes antipodes</i>	yellow-eyed	EN	3400	773.80	33.56	14.21	n/a	33.12	100	100
<i>Pygoscelis adeliae</i>	adelie	LC	7580000	104087.96	0.06	0.00	0.29	0.30	0.00	0.00
<i>Pygoscelis antarcticus</i>	chinstrap	LC	8000000	33972.38	3.55	5.10	0.48	4.99	0.26	100
<i>Pygoscelis papua</i>	gentoo	LC	774000	9872.58	14.05	16.07	10.90	17.36	0.00	0.00
<i>Spheniscus demersus</i>	African	EN	52000	10392.15	21.77	92.16	n/a	80.10	62.92	100
<i>Spheniscus humboldti</i>	Humboldt	VU	32000	7926.59	9.64	5.13	n/a	11.16	59.45	5.37
<i>Spheniscus magellanicus</i>	Magellanic	NT	2600000	75092.42	19.68	30.61	n/a	36.12	25.93	10.98
<i>Spheniscus mendiculus</i>	Galapagos	EN	1200	1821.34	0.00	100	n/a	100	100	100

650

651 * LC Least Concern, NT Near Threatened, VU Vulnerable, EN Endangered

652 ‡ NC Not categorized, ASPA Antarctic Specially Protected Areas

653

654 Table 2

	PREDICTOR	RESPONSE	R_s	P
PROTECTED AREA A COVERAGE	Range size	Total	0.65	0.004*
		IUCN	0.62	0.007*
		NC	0.46	0.05
		ASPA	0.67	0.002*
	Population	Total	0.30	0.22
		IUCN	0.46	0.05
		NC	0.21	0.40
		ASPA	0.71	0.001*
	Conservation Status†	Total	$\chi^2 = 1.19$	0.76
		IUCN	$\chi^2 = 3.46$	0.33
		NC	$\chi^2 = 0.91$	0.52
		ASPA	$\chi^2 = 7.09$	0.07
BIODIVERSITY HOTSPOTS	Range size	Coverage	0.09	0.73
		Protection	0.08	0.74
	Population	Coverage	-0.30	0.22
		Protection	-0.32	0.19
	Conservation Status†	Coverage	$\chi^2 = 1.10$	0.78
		Protection	$\chi^2 = 1.34$	0.72

655 * significant p-value

656 †Kruskal-Wallis test

657