



Severely restricting energy intake for 24 h does not affect markers of bone metabolism at rest or in response to re-feeding

David J. Clayton^{1,2} · Lewis J. James¹ · Craig Sale² · Iain Templeman³ · James A. Betts³ · Ian Varley²

Received: 13 August 2019 / Accepted: 17 January 2020
© The Author(s) 2020

Abstract

Purpose Intermittent energy restriction commonly refers to ad libitum energy intake punctuated with 24 h periods of severe energy restriction. This can improve markers of metabolic health but the effects on bone metabolism are unknown. This study assessed how 24 h severe energy restriction and subsequent refeeding affected markers of bone turnover.

Methods In a randomised order, 16 lean men and women completed 2, 48 h trials over 3 days. On day 1, participants consumed a 24 h diet providing 100% [EB: 9.27 (1.43) MJ] or 25% [ER: 2.33 (0.34) MJ] of estimated energy requirements. On day 2, participants consumed a standardised breakfast (08:00), followed by an ad libitum lunch (12:00) and dinner (19:30). Participants then fasted overnight, returning on day 3. Plasma concentrations of C-terminal telopeptide of type I collagen (CTX), procollagen type 1 N-terminal propeptide (P1NP) and parathyroid hormone (PTH) were assessed as indices of bone metabolism after an overnight fast on days 1–3, and for 4 h after breakfast on day 2.

Results There were no differences between trials in fasting concentrations of CTX, P1NP or PTH on days 1–3 ($P > 0.512$). During both trials, consuming breakfast reduced CTX between 1 and 4 h ($P < 0.001$) and PTH between 1 and 2 h ($P < 0.05$), but did not affect P1NP ($P = 0.773$). Postprandial responses for CTX ($P = 0.157$), P1NP ($P = 0.148$) and PTH ($P = 0.575$) were not different between trials. Ad libitum energy intake on day 2 was greater on ER [12.62 (2.46) MJ] than EB [11.91 (2.49) MJ].

Conclusions Twenty-four hour severe energy restriction does not affect markers of bone metabolism.

Keywords Intermittent energy restriction · Intermittent fasting · Bone · Bone metabolism · Weight management

Introduction

Net energy balance over time will dictate changes in body mass, with a chronic positive energy balance increasing both tissue mass and the risk of several chronic diseases [1]. For most people, weight gain tends to occur during

early to middle adulthood (18–49 years of age), with modest yearly weight increments eventually leading to a substantial excess of adipose tissue and increased risk of obesity-related chronic diseases [2]. This indicates strategies to help facilitate attaining and/or sustaining a healthy proportion of body fat are required.

Imposing a daily limit on energy intake, such as reducing habitual energy intake by 20–50%, is a commonly used method of achieving a healthy weight [3]. This method does not, however, appear to be effective for long-term maintenance of a healthy weight in most people [4], with the requirement for daily adherence a likely barrier to success [5]. One alternative method is intermittent energy restriction, which nominally involves alternating between periods of severely reduced energy intake (by ~ 75%) with periods of adequate or ad libitum energy intake [6]. Intermittent energy restriction can achieve similar weight loss and reduction in risk markers of obesity-related diseases compared to continuous energy restriction [7], suggesting

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s00394-020-02186-4>) contains supplementary material, which is available to authorized users.

✉ David J. Clayton
david.clayton@ntu.ac.uk

¹ National Centre for Sport and Exercise Medicine, School of Sport, Exercise and Health Sciences, Loughborough University, Loughborough, Leicestershire LE11 3TU, UK

² School of Science and Technology, Nottingham Trent University, Nottingham NG11 8NS, UK

³ Department for Health, University of Bath, Bath BA2 7AY, UK

this may be a viable alternative weight management strategy.

It is unknown whether severe energy restriction and the resultant acute perturbation of energy balance influences bone remodelling (i.e. osteoclastic bone resorption and osteoblastic bone formation). Bone mineral density is dictated by the balance between bone formation and resorption that occurs over time, which is a reflection of the physiological and mechanical environment [8]. Energy restriction is a risk factor for stress fracture incidence [9, 10], bone metabolism is markedly disrupted by 3–5 days of energy restriction (i.e. 10–30 kcal kg LBM⁻¹ day⁻¹), as evidenced by reductions in plasma markers of bone formation but without a corresponding decrease in bone resorption markers [11, 12]. Whilst these studies clearly demonstrate the potential for sustained moderate–severe energy restriction to impair bone health, typically, intermittent energy restriction involves shorter periods (1–2 days) of limited energy intake (~10–15 kcal kg LBM⁻¹ day⁻¹) but interspersed with refeeding periods [6]. Such cyclic energy restriction and replacement over serial days, therefore, represent a different model for which the daily kinetics of bone metabolism have never been assessed. One recent study has reported that the accumulated effect of 6 months of alternating daily between 25 and 125% of energy requirements did not alter post-absorptive markers of bone resorption (CTX, osteocalcin) or formation (bone alkaline phosphatase) at follow-up relative to standard continuous energy restriction or control [13]. However, this study employed a self-selected diet in the final 3 months of the intervention, with food diaries revealing only ~500 kcal difference between ‘fed’ and ‘fasting’ days in the alternate-day fasting condition. As such, the acute day-to-day variance in bone metabolism between the fasted days and subsequent refeeding days remains unknown.

Feeding is an essential stimulus for bone tissue accretion, as it causes an immediate postprandial reduction in markers of bone resorption and increases in markers of bone formation [14–16]. The majority of previous studies have assessed markers of bone metabolism in the fasted state, but the magnitude and duration of the response in bone metabolic markers after food intake may have a critical impact on bone remodelling. Recent studies have observed that the effects of short-term (1–2 days) severe energy restriction (consuming 10–15 kcal kg LBM⁻¹ day⁻¹) are not readily apparent in the fasted-state, but become pronounced upon refeeding. When comparing severe energy restriction diets to energy balanced maintenance control diets, previous studies have shown alterations in postprandial glucose, insulin, fatty acids, glucagon-like peptide-1, peptide-P and ghrelin after 24–48 h of severe energy restriction [17–20], with several of these hormones/substrates also postulated to influence bone remodelling [15].

To understand the efficacy and possible bone health effects of intermittent energy restriction diets, it is important to determine how a short period of severe energy restriction affects markers of bone remodelling upon refeeding. This will elucidate how intermittent energy restriction might affect bone health. Accordingly, this experiment investigated how markers of bone turnover respond temporally to an acute episode of severe energy restriction and refeeding, compared to an energy balanced control trial. This study was performed in young, lean individuals to investigate the effect of this eating pattern as a means to prevent weight gain.

Methods

Participants

This is a secondary analysis of a study that compared 24 h diets of 25% (ER) and 100% (EB) of estimated energy requirements (EER) on appetite regulation and energy intake [17]. This trial is registered at <https://www.clinicaltrials.gov.uk> as NCT02696772. Eighteen men ($n = 10$) and women ($n = 8$) were recruited and provided written informed consent to take part in the original study. Two men were removed from this analysis due to issues with obtaining blood samples during 1 trial (Table 1). Participants were healthy, weight stable (self-reported) and recreationally active (3–10 h week⁻¹).

Study design

Height, mass and body fat percentage were determined during a 1-day preliminary trial. Participants then completed two 3-day experimental trials administered in a randomised, crossover, and counterbalanced order. Trials were separated by at least 14 days for men and by exactly one menstrual cycle for women, who were tested during the post-menstruation follicular phase (5–12 days after the onset of menstruation) [21].

Table 1 Participant baseline characteristics

	Male ($n = 8$)	Female ($n = 8$)	Combined ($n = 16$)
Age (years)	23 (2)	21 (2)	22 (2)
Weight (kg)	75.6 (6.6)	63.8 (8.6)	69.7 (9.6)
Height (m)	1.80 (0.04)	1.61 (0.05)	1.70 (0.11)
BMI (kg m ⁻²)	23.4 (2.2)	24.5 (2.3)	24.0 (2.2)
Body fat (%)	14.3 (3.4)	27.3 (5.0)	20.8 (7.9)

Values are mean (1SD)

Protocol

During the 2 days immediately preceding the first experimental trial, participants recorded dietary intake and habitual physical activity, which was subsequently repeated in the 2 days immediately preceding the second experimental trial. Alcohol consumption and moderate/vigorous exercise were not permitted during this time or during the experimental trials. For each trial, participants arrived at the laboratory at 07:30 on three consecutive mornings having fasted at least 10 h, and post-void nude body mass was measured.

On day 1, a blood sample was collected via venepuncture (– 24 h), after which (08:30) participants left the laboratory with a 24 h diet providing either 25% or 100% of EER (resting metabolic rate estimated from Mifflin et al. [22] multiplied by 1.4) and instructions on when to consume each item. During EB, 100% of EER [9.27 (1.43) MJ] was distributed into four meals: 20% (of total food energy on EB) at 08:00 (cereal, milk and orange juice), 30% at 12:00 (white bread, chicken, mayonnaise, salad and cookies), 10% at 16:00 (yogurt and cereal bar), and 40% at 19:00 (pasta, chicken, Bolognese sauce, olive oil and cookies). During ER, 25% of EER [2.33 (0.34) MJ] was divided between two meals: 34% (of total food energy on ER) at 12:00 (chicken and salad) and 66% at 19:00 (pasta, chicken and Bolognese sauce), with a bolus of water consumed at 08:00 on ER, isovolume to the morning meal provided at 08:00 during EB. Additional water intake on both trials was prescribed at 35 mL kg⁻¹ body mass [2516 (283) mL] and was evenly distributed throughout the day. Diets were tailored to participants' preferences. The ER diet was created by primarily removing high-carbohydrate and high-fat foods from the EB diet to preserve protein content and ensure similar

foods were provided on day 1 for both trials. Lean body mass (LBM) was derived from body mass and body fat percentages determined from skinfold measurements collected at baseline [23]. It was estimated that the EB diet provided 40 (2) kcal kg LBM⁻¹ and the ER diet provided 10 (1) kcal kg LBM⁻¹. Participants were asked to perform minimal physical activity when outside the laboratory, including the strict avoidance of any structured exercise.

On day 2, participants arrived at 07:30, and remained in the laboratory until 20:00, with all food consumed in the laboratory. A fasted blood sample was collected (0 h; 08:00) from an indwelling cannula, after which participants consumed a standardised breakfast consisting of cereal, semi-skimmed milk, white bread, jam and butter, providing 25% of EER [2.44 (0.35) MJ; 93 (14) g carbohydrate; 16 (2) g protein; 15(2) g fat, 3 (0) g fibre]. Further blood samples were collected 1, 2, and 4 h after commencement of this meal. Participants were provided ad libitum lunch (12:00; 4 h) and dinner (19:00; 11 h) meals, as well as a standardised mid-afternoon snack [16:00; 8 h; 0.86 (0.12) MJ], as described previously [17]. Participants left the laboratory after dinner 20:00 (12 h), but were not permitted to consume any additional food or drink (other than plain water). Participants returned to the laboratory at 07:30, the following morning (day 3) and a final blood sample was collected via venepuncture (08:00; 24 h; Fig. 1).

Blood sample collection and analysis

C-terminal telopeptide of type I collagen (CTX) and procollagen type 1 N-terminal propeptide (PINP) were selected as markers of bone resorption and formation, as recommended by the International Osteoporosis Foundation and




Fig. 1 Schematic representation of study design

the International Federation of Clinical Chemistry [24]. Parathyroid hormone (PTH) was also measured as a marker of calcium metabolism. For each sample, blood was collected from an antecubital vein after at least 30 min of supine rest, and was dispensed into pre-chilled tubes containing EDTA (1.75 mg mL^{-1}). Plasma was separated by centrifuge (15 min; 1750g ; $4 \text{ }^{\circ}\text{C}$) and the supernatant was stored at $-20 \text{ }^{\circ}\text{C}$ for 24 h, before being transferred to $-80 \text{ }^{\circ}\text{C}$ for later analysis. Samples collected at -24 , 0 , 1 , 2 , 4 and 24 h were analysed for concentrations of total P1NP (Bioassay Technology Laboratory, Shanghai, China; $\text{CV} \leq 12.5\%$), CTX (Immunodiagnostic Systems, West Bolden, UK; $\text{CV} \leq 5.2\%$) and PTH (DiaMetra, Milan, Italy; $\text{CV} \leq 10.3\%$) by ELISA.

Statistical analysis

Data were analysed using SPSS 24.0 (SPSS Inc, Chicago, USA). All data were checked for normality using a Shapiro–Wilk test. Repeated measures ANOVA was used to evaluate main effects of time, trial and time-by-trial interactions, followed where necessary, by Holm–Bonferroni adjusted post hoc paired *t* tests (normally distributed data) or Wilcoxon signed-ranks tests (non-normally distributed data). Total area under the curve (AUC) was calculated in response to the standard breakfast (0–4 h) using the trapezoidal method and were analysed using a *t* test or Wilcoxon signed-ranks test, as appropriate. Sex was also entered as a between-participant factor in repeated measures to test for sex-by-time-by-trial and sex-by-trial interaction effects. Data sets were determined to be significantly different when $P < 0.05$. Data are presented as mean (1SD) unless stated otherwise.

Results

Body mass

There were main sex-by time ($P < 0.05$) and sex-by-time-by-trial ($P < 0.05$) effects, whereby body mass loss during ER

between day 1 and day 2 was greater in males than females ($P < 0.001$). However, these differences were due to the larger body mass of males compared to females, as when presented as a percentage change in body mass, no differences were noted ($P = 0.524$) and, therefore, male and female data are presented together. There were time ($P < 0.001$) and interaction ($P < 0.001$) effects for body mass. Body mass was not different between trials on day 1 ($P = 0.059$) or day 3 ($P > 1.00$), but was lower during ER on day 2 ($P < 0.01$). Body mass decreased from day 1 to day 2 during both trials, but to a greater extent during ER ($P < 0.001$; Table 2).

CTX and P1NP

There were no sex-by-time-by-trial effects for CTX ($P = 0.188$) or P1NP ($P = 0.426$), so male and female data are presented together. There were no differences between trials at -24 h (baseline) in plasma CTX ($P = 0.532$) or P1NP ($P = 0.548$) concentrations.

There was a time ($P < 0.001$), but no trial ($P = 0.489$) or interaction ($P = 0.157$) effects for CTX. Compared to baseline, plasma CTX concentrations were not different at 0 h ($P = 0.512$). Plasma CTX concentrations decreased after consuming breakfast, and were lower than baseline between 1 and 4 h, ($P < 0.001$) but were not different at 24 h ($P = 0.110$). There was no difference in postprandial AUC between trials for CTX ($P = 0.916$; Fig. 2a).

There were no time ($P = 0.773$), trial ($P = 0.584$) or interaction ($P = 0.148$) effects for P1NP (Fig. 2b). There was no difference in postprandial AUC between trials for P1NP ($P = 0.661$; Fig. 2b).

PTH

There were no sex-by-time-by-trial ($P = 0.471$) effects for PTH, so male and female data are presented together. There were no differences between trials in fasting plasma PTH ($P = 0.573$) concentrations at -24 h (baseline).

There was a time effect ($P < 0.001$), but no trial ($P = 0.566$) or interaction ($P = 0.575$) effects for PTH.

Table 2 Morning body mass measurements during each day of each experimental trial

	Energy balance (EB)			Energy restriction (ER)		
	Day 1 (-24 h)	Day 2 (0 h)	Day 3 (24 h)	Day 1 (-24 h)	Day 2 (0 h)	Day 3 (24 h)
Body mass (kg)	69.2 (9.6)	68.9 (9.6)*	68.9 (9.6)*	69.5 (9.8)	68.4 (9.5) ^{†,*}	68.9 (9.7)*
Δ from day 1 (kg)	–	-0.3 (0.3)	-0.3 (0.5)	–	-1.1 (0.4) [†]	-0.6 (0.4) [†]

Values are mean (1SD)

EB energy balance trial, ER energy restriction trial

[†]Significant difference to EB at corresponding time point

*Significant difference to -24 h during the same trial ($P < 0.05$)




Fig. 2 Top panel shows C-terminal telopeptide of type I collagen (CTX) (a), middle panel shows procollagen N-terminal propeptide (P1NP) (b) and bottom panel shows parathyroid hormone (PTH) (c). From left to right, concentrations at each time-point for all participants ($n=16$), concentrations at each time-point for male ($n=8$) and female ($n=8$) participants, total area under the curve for all par-

ticipants after consuming the standardised breakfast (0–4 h; 08:00–12:00). Hatched box indicates when breakfast was consumed. Data are mean with error bars representing standard error of the mean. Asterisk indicates the value is significantly different to baseline (–24 h)

Compared to baseline, plasma PTH concentrations were similar at 0 h ($P=1.00$). Plasma PTH concentrations decreased after consuming breakfast and were lower than baseline at 1 h ($P<0.001$) and 2 h ($P<0.05$), but not 4 h or 24 h ($P=1.00$). There was no difference in postprandial AUC between trials for PTH ($P=0.678$; Fig. 2c).

Energy intake

There was a between-participant main effect of sex, with ad libitum energy intake greater in males compared to females ($P<0.001$), but there were no sex-by-trial

($P=0.614$) or sex-by-time-by-trial ($P=0.086$) interaction effects for energy intake, so male and female data are presented together.

Energy provided in the pre-prepared diets on day 1 was 6.94 (1.00) MJ lower on ER compared to EB (Table 2). On day 2, total ad libitum energy intake (i.e. lunch and dinner) was greater during ER compared to EB [EB: 8.62 (2.14) MJ; ER: 9.33 (2.14) MJ; $P<0.05$]. The greater energy intake on day 2 [0.71 (1.27) MJ] was sufficient to replace ~11% of the energy deficit created on day 1 (Table 3).

Table 3 Total and relative energy and macronutrient intake during each day of the experimental trial

	Day 1		Day 2		Average	
	EB	ER	EB	ER	EB	ER
Protein (g)	97 (15)	60 (9) [†]	93 (20)	97 (18)	95 (16)	78 (12) [†]
Protein (g kg BM ⁻¹)	1.39 (0.11)	0.87 (0.08) [†]	1.34 (0.26)	1.41 (0.21)	1.37 (0.17)	1.14 (0.13) [†]
Carbohydrate (g)	293 (44)	56 (8) [†]	402 (91)	419 (98)	347 (64)	237 (52) [†]
Carbohydrate (g kg BM ⁻¹)	4.21 (0.11)	0.81 (0.05) [†]	5.77 (1.11)	6.09 (1.19)	4.99 (0.70)	3.45 (0.61) [†]
Fat (g)	69 (10)	9 (1) [†]	90 (23)	99 (21) [†]	80 (15)	54 (11) [†]
Fat (g kg BM ⁻¹)	0.99 (0.06)	0.13 (0.01) [†]	1.30 (0.06)	1.46 (0.30) [†]	1.14 (0.16)	0.79 (0.15) [†]
Fibre (g)	11 (2)	3 (1) [†]	21 (5)	22 (5)	16 (3)	13 (3) [†]
Fibre (g kg BM ⁻¹)	0.16 (0.01)	0.05 (0.00) [†]	0.31 (0.06)	0.32 (0.06)	0.23 (0.04)	0.18 (0.03) [†]
Energy (MJ)	9.23 (1.34)	2.33 (0.34) [†]	11.91 (2.49)	12.62 (2.46) [†]	10.59 (1.83)	7.48 (1.36) [†]
Energy (kJ kg BM ⁻¹)	133 (9)	34 (2) [†]	171 (30)	184 (29) [†]	157 (19)	109 (15) [†]

Values are mean (1SD) and are presented in grams (g), grams per kilogram of body mass (g kg BM⁻¹), megajoules (MJ) and kilojoules per kilogram of body mass (kJ kg BM⁻¹)

EB energy balance trial, ER energy restriction trial

[†]Significantly different from EB ($P < 0.05$)

Discussion

The present study shows that an acute episode of severe energy restriction has no effect on markers of bone metabolism in the fasted state, or for 4 h after consuming a high-carbohydrate breakfast. Although short periods of severe energy restriction (such as intermittent fasting) might not, therefore, have a deleterious effect on bone health, longer term studies utilising radiological scanning to elicit a more direct assessment of bone health are warranted.

This study assessed the effect of an acute 24 h period of severe energy restriction on markers of bone formation and resorption in fasted and postprandial states, finding that a diet providing ~ 10 kcal kg LBM⁻¹ had no immediate effects on markers of bone metabolism. In the fasted state, studies of longer periods of continuous energy restriction have observed a dose–response decrease in markers of bone formation (PICP) when energy availability was restricted to 10, 20 or 30 kcal kg LBM⁻¹ day⁻¹ for 5 days, but N-terminal telopeptide (NTX; a marker of bone resorption) was only increased in the 10 kcal kg LBM⁻¹ day⁻¹ condition [12]. Similarly, Papageorgiou et al. [11] showed a decrease in a marker of bone formation (PINP), but no change in a marker of bone resorption (CTX) after restricting energy availability to 15 kcal kg LBM⁻¹ day⁻¹ for 3 days. In combination with the current study, these studies suggest that the length of energy restriction or the total energy deficit created is important when considering the effect on bone formation and resorption. It may also be that a threshold in duration or magnitude exists, or a combination of these two factors, before energy restriction elicits a deleterious effect on bone metabolism, although this remains to be systematically investigated.

The current study also found there was no effect on markers of bone metabolism to feeding the following day. This is important as feeding is critical to the bone remodelling process [8]. As the current study only assessed the postprandial response to the first meal consumed after energy restriction, it is possible that delayed changes occurring in the subsequent hours or days were not captured. Studies that assess bone metabolism in response to multiple meals over multiple days would be required to elucidate this. The current data indicate that diets involving short periods of severe energy restriction alternated with longer periods of adequate energy intake might not detrimentally affect bone health. Whilst further research is required to determine the chronic effects, intermittent energy restriction diets may aid bone health to a greater extent than continuous energy restriction diets, which have consistently been shown to reduce bone health [25].

Consistent with the results of the current study, Barnosky et al. [13] found markers of bone metabolism in the fasted state were unchanged after 6-month alternate-day modified fasting or continuous energy restriction. Although surprisingly, Barnosky et al. [13] also reported no change in bone mineral content or BMD, in any condition, despite participants losing approximately 8% body mass. These measurements of bone remodelling were a secondary analysis of a larger study [26], which also found that when participants were permitted to self-select their diet, the disparity between ‘fasting’ and ‘fed’ days was less than 500 kcal by study endpoint. It is, therefore, debatable whether this study truly assessed the effects of intermittent severe energy restriction dieting. Nevertheless, weight-loss achieved via continuous energy restriction has been consistently shown to lead to reduced BMD [25]. Issues with dual-energy X-ray absorptiometry (DXA) in participants losing weight [8, 27], as

well as difficulty controlling exercise [28] and diet composition (e.g. calcium intake) [29], over a 6-month intervention period involving 3-months of self-selected food intake, might help to explain why these results differ from other diet-induced weight-loss studies.

The present study is the first to report how bone metabolic markers respond after energy restriction in the postprandial state, finding that 24 h of severe energy restriction did not affect markers of bone resorption or bone formation in response to feeding. Consuming a meal of individual or mixed macronutrient content causes a rapid suppression of bone resorption markers [14–16]. Feeding, therefore, acutely influences markers of bone metabolism, but whether this response is affected after a dietary intervention is rarely considered. Given the importance of feeding to bone metabolism, an impaired postprandial bone metabolic response could be indicative of a deleterious effect on BMD. Consistent with most acute studies, this study showed CTX (marker of bone resorption) was strongly suppressed after feeding, with no change in PINP (marker of bone formation). Importantly, the postprandial bone metabolic response was similar, whether participants had consumed 25% (~ 10 kcal kg LBM⁻¹) or 100% (~ 40 kcal kg LBM⁻¹) of their energy requirements the previous day, indicating that a single 24 h period of severe energy restriction does not interfere with the subsequent postprandial bone remodelling response. How bone metabolism responds during the energy restriction period itself remains unknown. It seems likely that energy restriction of this severity would have a negative impact on bone metabolism, but it would be interesting to explore in future studies what the net effect is of 24 h severe energy restriction and subsequent ad libitum refeeding on bone remodelling.

PTH is recognised as a key factor in bone remodelling, with elevated concentrations increasing bone turnover [30]. Results of the present study demonstrate that 24 h of severe energy restriction does not affect PTH concentrations, indicating that PTH does not change in response to acute perturbations in energy balance. Whilst no difference was shown in PTH, the original study on which this secondary analysis was performed showed that several factors associated with bone metabolism were affected by 24 h of severe energy restriction. After consuming a standardised meal, Clayton et al. [17] showed elevated plasma glucose, suppressed plasma acylated ghrelin and a tendency ($P=0.06$) for elevated plasma insulin concentrations, after 24 h severe energy restriction compared to 24 h of adequate energy intake. A similar study involving a short period of severe energy restriction (< 10% EER for 48 h) with subsequent refeeding similarly showed these changes, as well as elevated peptide P, glucagon-like peptide-1 and total ghrelin [18]. Despite several of these endocrine changes being linked to bone metabolism [15], no postprandial changes in markers

of bone metabolism in response to acute energy restriction were observed in the current study.

Mechanistic evidence explaining the involvement of endocrine factors on bone remodelling is derived primarily from studies that have identified corresponding receptors expressed on osteoblasts and/or osteoclasts [31]. However, in vivo evidence showing a clear effect of these endocrine factors on bone metabolism is lacking. For example, diabetes and insulin resistance are associated with increased fracture risk [32], but glucose-clamp studies infusing insulin across the physiological range showed no change in bone formation or resorption, suggesting insulin per se does not directly impact bone metabolism [33]. The results of the current study indicate that insulin and other aforementioned endocrine factors are not involved in the immediate post-prandial bone metabolic response following severe energy restriction. However, the time-course response for changes in markers of bone metabolism is not fully established. Although no differences were noted in fasting concentrations on days 1 and 2, it is possible that the observed postprandial changes in associated endocrine factors precede changes in bone metabolism. Therefore, this might not have been captured by the current study. Studies with longer monitoring periods, for example, over multiple restriction-repletion cycles and in response to several meals, are required to fully establish whether an acute episode of energy restriction affects bone remodelling. In addition, future studies should aim to investigate the cause of altered bone metabolism through dietary manipulation, by assessing fluctuations in hormones (e.g. incretin hormones and those associated with bone regulatory pathways such as Wnt signalling) that have a suggested association with bone metabolism.

Alongside diet, exercise is considered to be important for achieving successful long-term weight management [34]. Weight-bearing exercise is typically considered to have an anabolic effect on bone initiated via an increase in bone loading [35]. However, the extent to which exercise increases BMD is likely to be, at least partially, mediated by energy availability. A recent study [36] showed that restricting energy intake after a morning high-intensity exercise session to ~ 10 kcal kg BM⁻¹ resulted in greater concentrations of CTX in comparison to consuming a diet providing ~ 52 kcal kg BM⁻¹. This study also showed that energy provided specifically by carbohydrate, rather than overall energy availability was important in attenuating bone resorption after exercise. These results suggest the undertaking of exercise during intermittent energy restriction, particularly when implemented with carbohydrate restriction, may have a detrimental effect on bone metabolism.

Traditionally, intermittent fasting has been considered a means of achieving weight loss rather than weight maintenance. It is unclear from the current study whether acute severe energy restriction would prompt a similar bone

metabolic response in an overweight or obese population. Due to relatively greater bone loading caused by carrying additional weight, it might be expected that bone turnover would be increased in obese individuals, causing the accrual of bone mass if the loading stimulus is of an appropriate magnitude [25]. In line with Hammond et al. [36], it could be speculated that reduced energy and/or carbohydrate intake in this population may impair optimal bone remodeling leading to an increase in bone resorption. Future studies should aim to investigate the effects of severe energy restriction in overweight and obese individuals, as well as people at increased risk of fragility fracture, such as postmenopausal women.

In conclusion, 24 h of severe energy restriction had no effect on markers of bone resorption or bone formation. The acute nature of the intervention may explain why bone metabolism was unchanged. This suggests that short-term severe energy restriction as a method of weight control might not negatively affect bone health, but further studies are required to determine how repeated episodes to severe energy restriction influence long-term bone accrual.

Acknowledgements This report is an independent research by the National Institute for Health Research. The views expressed in this publication are those of the authors and not necessarily those of the NHS, the National Institute for Health Research or the Department of Health.

Author contributions The authors would like to thank Kirsty Burrell, Georgina Mynott, Mark Creese and Nicola Skidmore for their assistance with data collection. LJJ and DJC designed and carried out the clinical trial. Data analysis and interpretation were performed by DJC and IV, with assistance from LJJ, JAB and IT. DJC and IV wrote the manuscript, with assistance from LJJ, JAB, IT and CS.

Compliance with ethical standards

Conflict of interest LJJ has received funding for research from and/or acted as a consultant for Decathlon SA, the Collagen Research Institute, PepsiCo, Inc., Volac International Ltd, British Summer Fruits, Lucozade Ribena Suntory and Progressive Sports Technologies. LJJ has also received funding to attend conferences from GSSI and Danone Nutricia. In all cases, no personal payments were received by LJJ. CS has received research funding or support from the following sources in relation to bone health research: UK Ministry of Defence, GlaxoSmithKline, PepsiCo, English Institute of Sport and the Dairy Council. JAB has received research funding from BBSRC, GlaxoSmithKline, Lucozade Ribena Suntory, Kellogg's, Nestlé and PepsiCo and is a scientific advisor to the International Life Sciences Institute (ILSI). All other authors have no conflict of interest.

Ethical approval Participants gave full written consent prior to their inclusion in the study. The study was approved by the Loughborough University Ethics Approvals (Human Participants) Sub-Committee (Ref no: R13-P149) and has, therefore, been conducted in accordance with the 1964 Declaration of Helsinki and its later amendments. The research trial was registered at <https://www.clinicaltrials.gov.uk> as NCT02696772.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bray GA (2004) Medical consequences of obesity. *J Clin Endocrinol Metab* 89:2583–2589. <https://doi.org/10.1210/jc.2004-0535>
2. Ostbye T, Malhotra R, Landerman LR (2011) Body mass trajectories through adulthood: results from the National Longitudinal Survey of Youth Cohort (1981–2006). *Int J Epidemiol* 40:240–250. <https://doi.org/10.1093/ije/dyq142>
3. Omodei D, Fontana L (2011) Calorie restriction and prevention of age-associated chronic disease. *FEBS Lett* 585:1537–1542. <https://doi.org/10.1016/j.febslet.2011.03.015>
4. Greenberg I, Stampfer MJ, Schwarzfuchs D, Shai I, DIRECT Group (2009) Adherence and success in long-term weight loss diets: the dietary intervention randomized controlled trial (DIRECT). *J Am Coll Nutr* 28:159–168
5. Harvie M, Wright C, Pegington M et al (2013) The effect of intermittent energy and carbohydrate restriction v. daily energy restriction on weight loss and metabolic disease risk markers in overweight women. *Br J Nutr* 110:1537–1547. <https://doi.org/10.1017/S0007114513000792>
6. Longo VD, Mattson MP (2014) Fasting: molecular mechanisms and clinical applications. *Cell Metab* 19:181–192. <https://doi.org/10.1016/j.cmet.2013.12.008>
7. Schubel R, Nattenmuller J, Sookthai D et al (2018) Effects of intermittent and continuous calorie restriction on body weight and metabolism over 50 wk: a randomised controlled trial. *Am J Clin Nutr* 108:933–945. <https://doi.org/10.1093/ajcn/nqy196>
8. Harper C, Pattinson AL, Fernando HA, Zibellini J, Seimon RV, Sainsbury A (2016) Effects of obesity treatments on bone mineral density, bone turnover and fracture risk in adults with overweight or obesity. *Horm Mol Biol Clin Invest* 28:133–149. <https://doi.org/10.1515/hmbci-2016-0025>
9. Bennell K, Matheson G, Meeuwisse W, Brukner P (1999) Risk factors for stress fractures. *Sports Med* 28:91–122. <https://doi.org/10.2165/00007256-199928020-00004>
10. Ackerman KE, Holtzman B, Cooper KM, Flynn EF, Bruinvels G, Tenforde AS, Popp KL, Simpkin AJ, Parziale AL (2019) Low energy availability surrogates correlate with health and performance consequences of relative energy deficiency in sport. *Br J Sports Med* 53:628–633. <https://doi.org/10.1136/bjsports-2017-098958>
11. Papageorgiou M, Martin D, Colgan H, Cooper S, Greeves JP, Tang JCY, Frazer WD, Elliott-Sale KJ, Sale C (2018) Bone metabolic responses to low energy availability achieved by diet or exercise in active eumenorrheic women. *Bone* 114:181–188. <https://doi.org/10.1016/j.bone.2018.06.016>
12. Ihle R, Loucks AB (2004) Does-response relationship between energy availability and bone turnover in young

- exercising women. *J Bone Miner Res* 19:1231–1240. <https://doi.org/10.1359/JBMR.040410>
13. Barnosky A, Kroeger CM, Trepanowski JF, Klempel MC, Bhutani S, Hoddy KK, Gabel K, Shapses SA, Varady KA (2017) Effect of alternate day fasting on markers of bone metabolism: an exploratory analysis of a 6-month randomized controlled trial. *Nutr Healthy Aging* 4:255–263. <https://doi.org/10.3233/NHA-170031>
 14. Scott JPR, Sale C, Greeves JP, Casey A, Dutton J, Fraser WD (2012) Effect of fasting versus feeding on the bone metabolic response to running. *Bone* 51:990–999. <https://doi.org/10.1016/j.bone.2012.08.128>
 15. Clowes JA, Hannon RA, Yap TS, Hoyle NR, Blumsohn A, Eastell R (2002) Effect of feeding on bone turnover markers and its impact on biological variability of measurements. *Bone* 30:886–890
 16. Henriksen DB, Alexandersen P, Bjarnason NH, Vilsbøll T, Hartmann B, Eg Henriksen E, Byrjalsen I, Krarup T, Holst JH, Christiansen C (2003) Role of gastrointestinal hormones in postprandial reduction of bone resorption. *J Bone Miner Res* 18:2180–2189. <https://doi.org/10.1359/jbmr.2003.18.12.2180>
 17. Clayton DJ, Burrell K, Mynott G, Creese M, Skidmore N, Stensel DJ, James LJ (2016) Effect of 24-h severe energy restriction on appetite regulation and ad libitum energy intake in lean men and women. *Am J Clin Nutr* 104:1545–1553. <https://doi.org/10.3945/ajcn.116.136937>
 18. O'Connor KL, Scisco JL, Smith TJ, Young AJ, Montain SJ, Price LL, Lieberman HR, Karl JP (2016) Altered appetite-mediating hormone concentrations precede compensatory overeating after severe, short-term energy deprivation in healthy adults. *J Nutr* 146:209–217. <https://doi.org/10.3945/jn.115.217976>
 19. Antoni R, Johnston KL, Collins AL, Robertson MD (2016) Investigation into the acute effects of total and partial energy restriction on postprandial metabolism among overweight/obese participants. *Br J Nutr* 115:951–959. <https://doi.org/10.1017/S0007114515005346>
 20. Clayton DJ, Biddle J, Maher T, Funnell MP, Sargeant JA, King JA, Hulston CJ, Stensel DJ, James LJ (2018) 24-h severe energy restriction impairs postprandial glycaemic control in young, lean males. *Br J Nutr* 120:1107–1116. <https://doi.org/10.1017/S0007114518002568>
 21. Vescovi JD, VanHeest JL, De Souza MJ (2008) Short-term response of bone turnover to low-dose oral contraceptives in exercising women with hypothalamic amenorrhea. *Contraception* 77:97–104. <https://doi.org/10.1016/j.contraception.2007.10.007>
 22. Mifflin MD, St Jeor ST, Hill LA, Scott BJ, Daugherty SA, Koh YO (1991) A new predictive equation in healthy individuals for resting energy. *Am J Clin Nutr* 51:241–247. <https://doi.org/10.1093/ajcn/51.2.241>
 23. Durnin JVG, Womersley J (1974) Body fat assessed from total body density and its estimation from skinfold thickness: measurements on 481 men and women aged from 16 to 72 years. *Br J Nutr* 32:77–97. <https://doi.org/10.1079/bjn19740060>
 24. Vasikaran S, Cooper C, Eastell R, Griesmacher A, Morris HA, Trenti T, Kanis JA (2011) International Osteoporosis Foundation and International Federation of Clinical Chemistry and Laboratory Medicine position on bone marker standards in osteoporosis. *Clin Chem Lab Med* 49:1271–1274. <https://doi.org/10.1515/CCLM.2011.602>
 25. Zibellini J, Seimon RV, Lee CMY, Gibson AA, Hsu MSH, Shapses SA, Nguyen TV, Sainsbury A (2015) Does diet-induced weight loss lead to bone loss in overweight or obese adults? A systematic review and meta-analysis of clinical trials. *J Bone Miner Res* 30:2168–2178. <https://doi.org/10.1002/jbmr.2564>
 26. Trepanowski JF, Kroeger CM, Barnosky A, Klempel MC, Bhutani S, Hoddy KK, Gabel K, Freels S, Rigdon J, Rood J, Ravussin E, Varady KA (2017) Effect of Alternate-day fasting on weight loss, weight maintenance, and cardioprotection among metabolically healthy obese adults: a randomized clinical trial. *JAMA Intern Med* 177:930–938. <https://doi.org/10.1001/jamainternmed.2017.0936>
 27. Yu EW (2014) Bone metabolism after bariatric surgery. *J Bone Miner Res* 29:1507–1518. <https://doi.org/10.1002/jbmr.2226>
 28. Wilks DC, Winwood K, Gilliver S, Kwiet A, Chatfield M, Michaelis I, Sun L, Ferretti JL, Sargeant AJ, Felsenberg D (2009) Bone mass and geometry of the tibia and the radius of master sprinters, middle and long distance runners, race-walkers and sedentary control participants: a pQCT study. *Bone* 45:91–97. <https://doi.org/10.1016/j.bone.2009.03.660>
 29. Zhu K, Prince RL (2012) Calcium and bone. *Clin Biochem* 45:936–942. <https://doi.org/10.1016/j.clinbiochem.2012.05.006>
 30. Poole KES, Reeve J (2005) Parathyroid hormone—a bone anabolic and catabolic agent. *Curr Opin Pharmacol* 5:612–617. <https://doi.org/10.1016/j.coph.2005.07.004>
 31. Walsh JS, Eastell R (2009) Effects of feeding on bone metabolism. *Clin Rev Bone Miner Metab* 7:230–239. <https://doi.org/10.1007/s12018-009-9052-8>
 32. Tonks KT, White CP, Center JR, Samocha-Bonet D, Greenfield JR (2017) Bone turnover is suppressed in insulin resistance, independent of adiposity. *J Clin Endocrinol Metab* 102:1112–1121. <https://doi.org/10.1210/jc.2016-3282>
 33. Basu R, Peterson J, Rizza R, Khosla S (2011) Effects of physiological variations in circulating insulin levels on bone turnover in humans. *J Clin Endocrinol Metab* 96:1450–1455. <https://doi.org/10.1210/jc.2010-2877>
 34. Franz MJ, VanWormer JJ, Crain AL, Boucher JL, Histon T, Caplan W, Bowman JD, Pronk NP (2007) Weight-loss outcomes: a systematic review and meta-analysis of weight-loss clinical trials with a minimum 1-year follow-up. *J Am Diet Assoc* 107:1755–1767. <https://doi.org/10.1016/j.jada.2007.07.017>
 35. Ozcivici E, Luu YK, Adler BA, Qin Y-X, Rubin J, Judex S, Rubin CT (2010) Mechanical signals as anabolic agents in bone. *Nat Rev Rheumatol* 6:50–59. <https://doi.org/10.1038/nrrheum.2009.239>
 36. Hammond KM, Sale C, Fraser W, Tang J, Shepherd SO, Strauss JA, Close GL, Cocks M, Louis J, Pugh J, Stewart C, Sharples AP, Morton JP (2019) Post-exercise carbohydrate and energy availability induce independent effects on skeletal muscle cell signalling and bone turnover: implications for training adaptation. *J Physiol* 597:4779–4796. <https://doi.org/10.1113/JP278209>