

**Cumulative Lifetime Stress Exposure, Depression, Anxiety, and Well-Being in Elite  
Athletes: A Mixed-Method Study**

Ella McLoughlin<sup>1</sup>, David Fletcher<sup>1</sup>, George M. Slavich<sup>2</sup>, Rachel Arnold<sup>3</sup>, and Lee J. Moore<sup>3</sup>

<sup>1</sup>School of Sport, Exercise and Health Sciences, Loughborough University

<sup>2</sup>Cousins Center for Psychoneuroimmunology and Department of Psychiatry and Biobehavioral  
Sciences, University of California, Los Angeles

<sup>3</sup>Department for Health, University of Bath

**Author Note**

Ella McLoughlin is now at the Department for Health, University of Bath.

Declaration of interest: none.

Correspondence concerning this article should be addressed to Ella McLoughlin,  
Department for Health, University of Bath, Claverton Down, Bath, BA2 7AY, United Kingdom.

E-mail: em2050@bath.ac.uk

1 **Abstract**

2 Research suggests that elite athletes are at increased risk of poor mental health, partly due  
3 to the intense demands associated with top-level sport. Despite growing interest in the topic, the  
4 factors that influence the mental health and well-being of elite athletes remain unclear. From a  
5 theoretical perspective, the accumulation of stress and adversity experienced over the life course  
6 may be an important factor. To investigate this possibility, we employed a mixed-method design  
7 to: (a) examine whether cumulative lifetime stress predicted depression, anxiety, and well-being  
8 in elite athletes; and (b) help explain why cumulative lifetime stress exposure might have resulted  
9 in poor mental health and well-being. Ninety-five elite athletes ( $M_{age} = 29.81, SD = 10.88$ )  
10 completed the Stress and Adversity Inventory, Patient Health Questionnaire, Generalized Anxiety  
11 Disorder 7-item scale, and the Scales of General Well-Being. Hierarchical regression  
12 analyses revealed that total count and severity of lifetime stressor exposure significantly predicted  
13 greater depression ( $\beta = .42, p < .001; \beta = .46, p < .001$ ) and anxiety symptoms ( $\beta = .34, p = .003;$   
14  $\beta = .28, p = .018$ ), and worse well-being ( $\beta = -.42, p < .001; \beta = -.30, p = .015$ ). Semi-structured  
15 interviews were then conducted with six athletes. Thematic analysis revealed that cumulative  
16 lifetime stress exposure fostered poor mental health and well-being by promoting maladaptive  
17 long-term coping strategies, increasing susceptibility to future stress, and limiting interpersonal  
18 relationships. These findings can help practitioners identify, and intervene accordingly with, elite  
19 athletes at risk of experiencing stress-related mental health problems.

20 *Keywords:* adversity, allostatic load, life stress, pressure, sport



1 stressors that a person has experienced across their lifespan (Lam, Shields, Trainor, Slavich, &  
2 Yonelinas, 2019). The integrative model is of particular relevance as it introduces important  
3 conceptual dimensions of cumulative lifetime stress, whereby stressors are distinguished by their  
4 timing (i.e., early life vs. adulthood), type (i.e., acute life events vs. chronic difficulties), primary  
5 life domain (e.g., housing, education, work), and social-psychological characteristic (e.g.,  
6 interpersonal loss, humiliation, entrapment). This conceptual differentiation is important, given  
7 that different types of stressors have varying effects on health and well-being (Epel et al., 2018).  
8 Well-being, in turn, is defined as a multidimensional and complex construct with two distinct  
9 perspectives: hedonic and eudaimonic (Neil, McFarlane, & Smith, 2017). Whereas the hedonic  
10 perspective focuses on the subjective experience of happiness and life satisfaction, the eudaimonic  
11 perspective focuses on psychological functioning and self-realization (Ryan & Deci, 2001).  
12 Moreover, mental health is defined by the World Health Organization (2004) as “a state of well-  
13 being in which the individual realises his or her own abilities, can cope with the normal stresses  
14 of life, can work productively and fruitfully, and is able to make a contribution to his or her  
15 community” (p. 12). Pursuant with the predictions of the integrative model, it is important to  
16 identify the specific types of stressors that are particularly harmful for health (Epel et al., 2018).

17 Broadly speaking, the effect of cumulative lifetime stress exposure has recently been  
18 examined in relation to a variety of psychological, biological, and clinical outcomes (e.g., Olvera  
19 Alvarez et al., 2019; Pegg et al., 2019). The consensus from this body of work is that as the  
20 incidence of acute life events and chronic difficulties increases, so too does the risk for poor health  
21 outcomes (Slavich & Shields, 2018). For example, greater lifetime stress exposure has been  
22 associated with greater depression and anxiety in adolescents (e.g., Slavich, Stewart, Esposito,  
23 Shields, & Auerbach, 2019; Toussaint, Shields, Dorn, & Slavich, 2016), with cumulative stressor

1 severity being a relatively stronger predictor than the total count of stressors experienced (Slavich  
2 et al., 2019). In addition, chronic difficulties have been found to be a marginally stronger predictor  
3 of depression and anxiety, than acute life events (Slavich et al., 2019). Although this quantitative  
4 population-based research is insightful, combining this approach with qualitative methods may  
5 help researchers better understand precisely how cumulative lifetime stress affects mental health  
6 and well-being (i.e., underlying mechanisms; Greene, Caracelli, & Graham, 1989).

7         In addition to the aforementioned findings, emerging research within a sport context  
8 suggests that experiencing moderate amounts of adversity can lead to enhanced psychological  
9 resilience (e.g., Fletcher, 2019; Fletcher & Sarkar, 2012), psychosocial growth (e.g., Howells &  
10 Fletcher, 2015), and more optimal sports performance (e.g., Moore, Young, Freeman, & Sarkar,  
11 2017). For example, Hardy et al. (2017) reported that early life non-sport adversity was a consistent  
12 feature in Olympic and world champions lives, particularly when experienced closely to a positive  
13 sport-related event. Despite these emergent findings, researchers in sport have predominantly  
14 examined how the *frequency* of adverse life events is associated with performance-related  
15 outcomes (e.g., Moore et al., 2017). This differs from examining the combined and cumulative  
16 effect of stressors occurring across an athletes' lifespan (Fletcher, 2019). Accordingly, additional  
17 research is required that adopts a more multidimensional and life course perspective, where  
18 researchers identify and examine the underlying dimensions of a stressful exposure (e.g.,  
19 frequency, intensity, timing, duration, and severity), in order to better understand how different  
20 aspects of lifetime stress exposure are related to mental health and well-being in elite sport.

21         The purpose of the present study was to examine how cumulative lifetime stress exposure  
22 was associated with symptoms of depression and anxiety, as well as well-being, among elite  
23 athletes. Given that the majority of research on this topic outside of sport has found linear

1 relationships between lifetime stress exposure and mental health and well-being (e.g., Slavich &  
2 Shields, 2018), it was hypothesized that greater cumulative lifetime stress exposure (both total  
3 stressor count and severity) would predict poorer mental health and well-being, but that these  
4 associations would differ by stressor timing, type, life domain, and social-psychological  
5 characteristic (e.g., Slavich et al., 2019). This study also collected qualitative data to help explain  
6 why cumulative lifetime stress exposure might have resulted in poor mental health and well-being.

## 7 **Method**

### 8 **Study Design**

9 This study was underpinned by methodological pluralism (i.e., drawing on both positivism  
10 and interpretive epistemologies) as opposed to methodological puritanism (i.e., affinity to a single  
11 paradigm; Onwuegbuzie & Leech, 2005). This approach enabled a more comprehensive  
12 understanding of phenomena, whereby the weaknesses of one approach were addressed by the  
13 strengths of the other (Johnson & Onwuegbuzie, 2004). Driven by this approach, the underlying  
14 theme of this research was pragmatism (Teddlie & Tashakkori, 2011).

15 Consistent with this research paradigm, a mixed-method explanatory sequential (two-  
16 phase) design was employed, with qualitative data (phase two) used to partly explain quantitative  
17 results (phase one). As such, integration of methods occurred across chronological phases of the  
18 study, where the collection and analysis of quantitative data (i.e., phase one) sought to inform the  
19 direction of the qualitative phase (i.e., phase two). Consistent with this approach, the weighting of  
20 each method was unequal. Specifically, since the quantitative data informed the qualitative method,  
21 greater attention was focused on the quantitative data (Creswell & Plano-Clark, 2011).

### 22 **Participants**

23 Phase one was completed by 95 participants between 18 and 60 years old (58 females, 37

1 males;  $M_{\text{age}} = 29.81$ ,  $SD = 10.88$ ) from a variety of sports (e.g., athletics, swimming, triathlon,  
2 netball, football, and hockey). All participants were classified as elite athletes because they had  
3 competed, or were currently competing, at an international or professional level (Swann, Moran,  
4 & Piggott, 2015). An *a priori* power calculation using G\*Power software revealed that a minimum  
5 of 60 participants was required given a medium effect size of 0.30, an alpha of 0.05, and power of  
6 0.80. During phase two, criterion-based purposeful sampling identified participants that were in  
7 the top 10% of lifetime stressor count ( $M = 38.60$ ,  $SD = 13.26$ ) and severity ( $M = 97.30$ ,  $SD =$   
8  $30.50$ ). Six participants (five females, one male) between 28 and 36 years old ( $M_{\text{age}} = 32.17$ ,  $SD =$   
9  $3.19$ ) were interviewed. These participants were from a range of sports, including athletics,  
10 triathlon, ultrarunning, and powerlifting. The sample size for phase two offered sufficient  
11 information power given the quality of data garnered from in-depth interviews (Malterud, Siersma,  
12 & Guassora, 2016). Specifically, the quality of the data was assessed via the communication  
13 between the researcher and the participant. Due to the explanatory nature of the qualitative data,  
14 semi-structured interviews had a specific aim, which promoted a clear and focused dialogue. Next,  
15 the homogeneity of the sample provided sufficient information power, given that participants had  
16 experienced the phenomena in question (i.e., lifetime stress exposure), thus, belonging to the  
17 specified target group. Finally, the mean duration of interviews suggested that participants  
18 provided in-depth and rich detail relating to lifetime stress and health (Malterud et al., 2016).

## 19 **Procedure**

20 Following institutional ethical approval, participants were recruited using the researchers'  
21 existing contacts and social media (e.g., Twitter). After providing informed consent, participants  
22 completed the online questionnaire, which took approximately 30 minutes. Once the quantitative  
23 data had been analyzed, participants in the top 10% of self-reported cumulative lifetime stressor

1 count and severity were emailed to arrange an in-depth interview at a convenient time. Six  
2 participants agreed to be interviewed and completed semi-structured interviews on Skype ( $M_{duration}$   
3 = 57.90 minutes,  $SD = 12.85$ ; range = 40-76 minutes). A semi-structured interview format was  
4 adopted as this was considered the most effective and convenient method for gathering in-depth  
5 information (Kvale & Brinkmann, 2009). These interviews were informed by an interview guide  
6 that contained questions relating to lifetime stress and health, and comprised of four sections,  
7 including: introductory questions (e.g., “how did you first get involved in your sport?”); questions  
8 relating to the stressors experienced over the life course (e.g., “can you tell me about an event or  
9 period during your childhood that you found particularly difficult?”); questions relating to how  
10 participants reacted to the culmination of stressors over the life course (e.g., “how did you respond  
11 to these stressors and adversities?”); and questions identifying the impact of lifetime stress on  
12 mental health and well-being (e.g., “can you please explain how these stressors and adversities  
13 affected your mental health and well-being?”). The interviews were recorded with audio  
14 equipment and manually transcribed.

## 15 **Measures**

### 16 ***Cumulative Lifetime Stressor Exposure***

17 Cumulative lifetime stress exposure was assessed using the Stress and Adversity Inventory  
18 for Adults (Adult STRAIN; Slavich & Shields, 2018). The Adult STRAIN is an online stress-  
19 assessment system that enquires about 55 major stressors (e.g., job loss, death of a relative). For  
20 every stressor that is endorsed, follow-up questions are asked that assess the severity (1 = *not at*  
21 *all* to 5 = *extremely*), frequency (1 to 5 or more times), timing (1 = *ongoing* to 7 = *over 5 years*),  
22 and duration (an open box where participants indicate years and/or months) of that stressor  
23 (Slavich & Shields, 2018). Stressors that are assessed with the Adult STRAIN span two time

1 periods (early life vs. adulthood), and can be categorized into two stressor types (acute life events  
2 vs. chronic difficulties), 12 major life domains (housing, education, work, health, marital/partner,  
3 reproduction, financial, legal, other relationships, death, life-threatening situations, and  
4 possessions), and five social-psychological characteristics (interpersonal loss, physical danger,  
5 humiliation, entrapment, and role change/disruption). The two main variables used in this study  
6 were: (1) total count of lifetime stressors, which was calculated by summing the number of  
7 stressors encountered by participants (range = 0-159), and (2) total cumulative severity of lifetime  
8 stressors, which was calculated by summing the perceived severity of the stressors experienced by  
9 participants (range = 0-275). The Adult STRAIN has very good concurrent ( $r_s = .15 - .62$ ) and  
10 discriminant validity, with excellent test-retest reliability ( $r_s = .90 - .95$ ). Furthermore, the  
11 predictive validity of the Adult STRAIN has previously been demonstrated in relation to various  
12 outcomes, including mental and physical health complaints, sleep quality, and cognitive function  
13 among others (e.g., Cazassa et al., 2020; Slavich & Shields, 2018; Sturmbauer et al., 2020).

#### 14 ***Depression***

15 The Patient Health Questionnaire (PHQ-9) was used to assess symptoms of depression over  
16 the past two weeks. The PHQ-9 consists of nine items (e.g., little interest or pleasure in doing  
17 things), with each item scored on a scale ranging from 0 (*not at all*) to 3 (*nearly every day*). Higher  
18 scores indicated greater symptoms (range = 0-27), with scores of 5, 10, 15, and 20 representing  
19 mild, moderate, moderately severe, and severe depression, respectively (Kroenke, Spitzer, &  
20 Williams, 2001). The PHQ-9 has good internal consistency ( $\alpha = .86 - .89$ ) and test-retest reliability  
21 ( $r = .84$ ), as well as good construct and criterion validity (Kroenke et al., 2001).

#### 22 ***Anxiety***

23 The Generalized Anxiety Disorder (GAD-7) scale was used to assess symptoms of anxiety

1 over the past two weeks. The GAD-7 consists of seven items (e.g., feeling nervous, anxious, or on  
2 edge), with each item scored on a scale ranging from 0 (*not at all*) to 3 (*nearly every day*). Higher  
3 scores indicated greater symptoms (range = 0-21), with scores of 5, 10, and 15 representing mild,  
4 moderate, and severe anxiety, respectively. The GAD-7 has good internal consistency ( $\alpha = .89 -$   
5  $.92$ ) and test-retest reliability ( $r_s = .83$ ), as well as convergent, construct, criterion, and factorial  
6 validity (Löwe et al., 2008; Spitzer, Kroenke, Williams, & Löwe, 2006).

### 7 ***Well-being***

8 The Scales of General Well-Being (SGWB) was used to assess multiple indicators of well-  
9 being (e.g., happiness, vitality, self-acceptance; Longo, Coyne, & Joseph, 2018). The SGWB  
10 consists of 14 items (e.g., I have a purpose), with each item scored on a scale ranging from 1 (*not*  
11 *at all true*) to 5 (*very true*). Higher scores indicated better well-being (range = 14-70), with scores  
12 above 40 representing 'good' well-being. The SGWB has demonstrated good internal consistency  
13 ( $\alpha = .86$ ) and construct validity ( $r_s = .67 - .88$ ; Longo et al., 2018).

### 14 **Data Analysis**

15 Quantitative data were analyzed using SPSS version 25.0. Prior to any analyses, outliers  
16 ( $z$ -scores greater or less than 3.29) were removed from the dataset (six values for total lifetime  
17 stressor count, seven values for total lifetime stressor severity, four values for PHQ-9, and three  
18 values for GAD-7). Following these outlier analyses, visual inspection confirmed that all data were  
19 normally distributed. First, to examine if total lifetime stressor count or severity predicted  
20 depression, anxiety, and well-being, a series of hierarchical linear regression analyses were  
21 conducted. Specifically, depression, anxiety, and well-being were entered into separate models as  
22 dependent variables, while in each model, independent variables were entered at step 1 (total

1 lifetime stressor count or severity), and *a priori* covariates were entered at step 2 (age and gender)<sup>1</sup>.  
2 Next, a series of hierarchical linear regression analyses were conducted to examine if the different  
3 stressor types (acute life events vs. chronic difficulties), time periods (early-life vs. adulthood), life  
4 domains (e.g., work, health, death), and social-psychological characteristics (e.g., physical danger,  
5 humiliation, entrapment) significantly predicted depression, anxiety, and well-being, above and  
6 beyond age and gender. Three life domains (education, legal, and reproduction) were excluded  
7 from analyses, as very few participants reported experiencing these stressors.

8         Qualitative data were analyzed using a confirmatory approach to thematic analysis because  
9 the quantitative findings guided the ideas the researcher assessed via semi-structured interviews  
10 (Guest, MacQueen, & Namey, 2012). The six-step procedure described by Braun and Clarke  
11 (2006) was adopted. Analysis began by listening to interview recordings, and reading and re-  
12 reading each transcript, prior to the development of preliminary codes. The initial list of codes was  
13 then reduced through grouping to identify overarching themes that were relevant throughout the  
14 entire dataset. Themes were then refined, defined and named, before the results were written. As  
15 the qualitative component was subordinate to the quantitative element, themes were identified at  
16 a semantic level to help explain the quantitative results (Braun & Clarke, 2006).

### 17 **Methodological Integrity**

18         Approaches were considered that sought to enhance the methodological integrity of the  
19 qualitative data via its two constituents: fidelity and utility (Levitt, Motulsky, Wertz, Morrow, &  
20 Ponterotto, 2017). Fidelity to the subject matter was established using perspective management in

---

<sup>1</sup> Curvilinear or quadratic relationships between lifetime stress exposure (i.e., total count and severity of lifetime stress) and outcomes (i.e., depression, anxiety, and well-being) were also assessed via curve estimation regression analyses. The results revealed no significant curvilinear or quadratic relationships between any of the lifetime stress exposure and outcome variables.

1 data collection. To elaborate, the researcher engaged in a process of bracketing to identify one's  
2 implicit assumptions and biases (Morrow, 2005). Bracketing was achieved through the use of a  
3 reflexive journal where the researcher's influence on the data collection process was identified. As  
4 a result, the use of a reflexive journal sought to limit the influence of the researcher's implicit  
5 assumptions and biases to obtain a clearer representation of phenomena (Levitt et al., 2017).  
6 Furthermore, utility in achieving goals was achieved by collecting data that provided rich grounds  
7 for insightful analyses (Levitt et al., 2017). To elaborate, the methods adopted within the present  
8 study identified participants that had experienced the phenomena in question (i.e., lifetime stress  
9 exposure), resulting in enhanced insight to be derived from the data (Levitt et al., 2017).

## 10 **Results**

### 11 **Quantitative Findings**

#### 12 *Descriptive Statistics*

13 Participants experienced an average of 13 stressors over their lifetime (range = 0-54),  
14 including an average of eight acute life events (range = 0-43), five chronic difficulties (range = 0-  
15 26), and three early (range = 0-22) and ten adulthood (range = 0-48) adversities. The mean  
16 cumulative lifetime stressor severity was 31.38 ( $SD = 27.70$ ). On average, participants reported  
17 relatively low symptoms of depression ( $M = 4.15$ ,  $SD = 3.62$ ) and anxiety ( $M = 3.49$ ,  $SD = 3.04$ ),  
18 and relatively high levels of well-being ( $M = 53.67$ ,  $SD = 9.58$ ). Overall, 1.3%, 2.6%, 3.8%, and  
19 24% of participants met the criteria for severe, moderately severe, moderate, and mild depression,  
20 respectively. In addition, 1.2%, 3.8%, and 21.8% of participants met the criteria for severe,  
21 moderate and mild anxiety, respectively. No participants met the criteria for both severe depression  
22 and anxiety. Bivariate correlation analyses revealed that most of the STRAIN variables were  
23 moderately associated with poorer mental health and well-being (Table 1).

### 1 *Lifetime Stressor Count, Mental Health, and Well-Being*

2 Cumulative lifetime stressor count significantly predicted symptoms of depression ( $\beta = .42$ ,  
3  $p < .001$ ) and anxiety ( $\beta = .34$ ,  $p = .003$ ), and well-being ( $\beta = -.42$ ,  $p < .001$ ), above and beyond  
4 age and gender (Table 2). Examining stressor type revealed that chronic difficulties and acute life  
5 events both significantly predicted greater symptoms of depression ( $\beta = .44$ ,  $p < .001$ ;  $\beta = .30$ ,  $p$   
6  $= .007$ ) and anxiety ( $\beta = .24$ ,  $p = .040$ ;  $\beta = .24$ ,  $p = .039$ ), and well-being ( $\beta = -.35$ ,  $p = .004$ ;  $\beta =$   
7  $-.28$ ,  $p = .018$ ), above and beyond age and gender. With respect to stressor exposure timing, total  
8 count of early adversities significantly predicted symptoms of depression above and beyond age  
9 and gender ( $\beta = .25$ ,  $p = .030$ ), but did not significantly predict symptoms of anxiety ( $\beta = .22$ ,  $p$   
10  $= .065$ ) or well-being ( $\beta = -.16$ ,  $p = .180$ ). Total count of adulthood adversities, in contrast,  
11 significantly predicted symptoms of depression ( $\beta = .43$ ,  $p < .001$ ) and anxiety ( $\beta = .32$ ,  $p = .006$ ),  
12 as well as well-being ( $\beta = -.40$ ,  $p = .001$ ), above and beyond age and gender.

13 In terms of stressor characteristics, the total count of stressors that significantly predicted  
14 symptoms of depression across most of the life domains and social-psychological characteristics,  
15 while controlling for age and gender, including: work ( $\beta = .34$ ,  $p = .002$ ), marital/partner ( $\beta = .29$ ,  
16  $p = .010$ ), financial ( $\beta = .28$ ,  $p = .011$ ), other relationships ( $\beta = .29$ ,  $p = .007$ ), entrapment ( $\beta = .40$ ,  
17  $p < .001$ ), role change/disruption ( $\beta = .33$ ,  $p = .002$ ), humiliation ( $\beta = .30$ ,  $p = .006$ ), and  
18 interpersonal loss ( $\beta = .28$ ,  $p = .011$ ). Housing, health, death, life-threatening situations,  
19 possessions, and physical danger were not significantly associated with symptoms of depression  
20 (Figure 1a). The stressor characteristics that significantly predicted symptoms of anxiety, while  
21 controlling for age and gender, included: role change/disruption ( $\beta = .26$ ,  $p = .025$ ), life-threatening  
22 situations ( $\beta = .26$ ,  $p = .022$ ), other relationships ( $\beta = .24$ ,  $p = .034$ ), and humiliation ( $\beta = .31$ ,  $p$   
23  $= .005$ ). Housing, work, health, marital/partner, financial, death, entrapment, possessions,

1 interpersonal loss, and physical danger were not associated with symptoms of anxiety (Figure 1b).  
2 Finally, the stressor characteristics that significantly predicted well-being, while controlling for  
3 age and gender, included: health ( $\beta = -.33, p = .005$ ), role change ( $\beta = -.36, p = .002$ ),  
4 marital/partner ( $\beta = -.26, p = .024$ ), other relationships ( $\beta = -.25, p = .027$ ), life-threatening  
5 situations ( $\beta = -.25, p = .032$ ), entrapment ( $\beta = -.25, p = .034$ ), and physical danger ( $\beta = -.35, p$   
6  $= .005$ ). Housing, work, financial, death, possessions, interpersonal loss, and humiliation were not  
7 associated with well-being (Figure 1c).

### 8 ***Lifetime Stressor Severity, Mental Health, and Well-Being***

9 In terms of stressor severity, total severity of lifetime stressors significantly predicted  
10 symptoms of depression ( $\beta = .46, p < .001$ ) and anxiety ( $\beta = .28, p = .018$ ), as well as well-being  
11 ( $\beta = -.30, p = .015$ ), above and beyond age and gender (Table 2). Examining stressors types  
12 revealed that severity of acute life events significantly predicted symptoms of depression ( $\beta = .39,$   
13  $p = .001$ ) and anxiety ( $\beta = .36, p = .003$ ), in addition to well-being ( $\beta = -.30, p = .019$ ), above and  
14 beyond age and gender. In contrast, total severity of chronic difficulties significantly predicted  
15 symptoms of depression ( $\beta = .43, p < .001$ ) and well-being ( $\beta = -.30, p = .010$ ), above and beyond  
16 age and gender, but not symptoms of anxiety ( $\beta = .21, p = .063$ ). With respect to stress exposure  
17 timing, total severity of early adversities significantly predicted symptoms of anxiety above and  
18 beyond age and gender ( $\beta = .32, p = .010$ ), but not symptoms of depression ( $\beta = .21, p = .083$ ) or  
19 well-being ( $\beta = -.20, p = .105$ ). In contrast, total severity of adulthood adversities significantly  
20 predicted symptoms of depression ( $\beta = .52, p < .001$ ) and anxiety ( $\beta = .24, p = .045$ ), as well as  
21 well-being ( $\beta = -.38, p = .002$ ), above and beyond age and gender.

22 In terms of stressor characteristics, the total severity of stressors that significantly predicted  
23 symptoms of depression across the different life domains and social-psychological characteristics,

1 while controlling for age and gender, included: housing ( $\beta = .30, p = .008$ ), marital/partner ( $\beta = .42,$   
2  $p < .001$ ), other relationships ( $\beta = .28, p = .012$ ), work ( $\beta = .30, p = .008$ ), health ( $\beta = .27, p = .020$ ),  
3 financial ( $\beta = .30, p = .006$ ), humiliation ( $\beta = .41, p < .001$ ), entrapment ( $\beta = .40, p < .001$ ),  
4 interpersonal loss ( $\beta = .32, p = .004$ ), physical danger ( $\beta = .29, p = .010$ ), and role change/disruption  
5 ( $\beta = .27, p = .015$ ). Death, life-threatening situations, and possessions were not associated with  
6 symptoms of depression (Figure 1a). The stressor characteristics that significantly predicted  
7 symptoms of anxiety, while controlling for age and gender, included: housing ( $\beta = .24, p = .043$ ),  
8 marital/partner ( $\beta = .24, p = .029$ ), other relationships ( $\beta = .34, p = .002$ ), life-threatening situations  
9 ( $\beta = .28, p = .018$ ), interpersonal loss ( $\beta = .28, p = .015$ ), humiliation ( $\beta = .35, p = .002$ ), physical  
10 danger ( $\beta = .30, p = .012$ ), entrapment ( $\beta = .24, p = .033$ ), and role change/disruption ( $\beta = .25, p$   
11  $= .027$ ). Work, health, financial, death, and possessions were not associated with symptoms of  
12 anxiety (Figure 1b). Finally, the stressor characteristics that significantly predicted well-being,  
13 while controlling for age and gender, included: work ( $\beta = -.30, p = .014$ ), marital/partner ( $\beta = -.46,$   
14  $p < .001$ ), life-threatening situations ( $\beta = -.27, p = .024$ ), interpersonal loss ( $\beta = -.30, p = .011$ ),  
15 physical danger ( $\beta = -.37, p = .002$ ), humiliation ( $\beta = -.38, p = .001$ ), and entrapment ( $\beta = -.34, p$   
16  $= .004$ ). Housing, health, financial, other relationships, death, possessions, and role  
17 change/disruption were not associated with well-being (Figure 1c).

## 18 **Qualitative Findings**

19 The qualitative data collected in this study was used to help explain why cumulative  
20 lifetime stress exposure (count and severity) might have resulted in poor mental health and well-  
21 being among elite athletes. Three themes were identified: (1) maladaptive long-term coping  
22 strategies, (2) increased susceptibility to future stress, and (3) interpersonal relationship difficulties.  
23 Pseudonyms are used throughout to protect participants identities.

### 1 *Maladaptive Long-Term Coping Strategies*

2           Three participants explained how experiencing negative childhood adversities led to the  
3 development of maladaptive coping styles. This notion was exemplified by Mary: “I lost my  
4 mother at 12... I didn’t really deal with it very well; I threw myself into anything other than being  
5 at home. I trained intensely”. Despite participants attributing the success of their athletic career to  
6 the stressors and adversities they experienced, they also identified the long-standing mental health  
7 issues associated with their coping approach, as described by Elizabeth: “It got to a point where  
8 nothing was good enough and I was never satisfied. I was overtraining, I was at burnout, I was  
9 self-harming. I was damaging myself because I couldn’t cope”. As such, participants used sport as  
10 a short-term palliative coping strategy, which ultimately led to maladaptive coping in the longer  
11 term. This was supported by Sarah: “[sport] became the only thing I had... In the short term,  
12 [experiencing stress] had a fantastic effect on my sport but in the long-term, I was emotionally  
13 stunted and crippled”. As such, an exclusive focus on sport following adversity led to the  
14 development of a constrained identity, this was further supported by Elizabeth: “Running was all  
15 I had and all I focused on... It made me push myself and I don’t think I would’ve got to the level  
16 that I did if I hadn’t been through everything that I experienced”. Thus, this theme revealed how  
17 the elite athletes initially developed positively after experiencing stress and adversity, but, also,  
18 how it led to maladaptive coping strategies that negatively impacted their mental health and well-  
19 being in the long-term.

### 20 *Increased Susceptibility to Future Stress*

21           Three participants indicated that encountering adverse experiences in developmentally  
22 sensitive periods may have acted as a catalyst for experiencing subsequent stress. For example,  
23 Daniel stated: “They [the stressors] just fed off one another... If something was going on outside

1 of sport, my sport performance would decrease, which meant I got deselected... The continuous  
2 and constant nature of stressors just fed into mental health problems... It was just this downward  
3 spiral that I couldn't get out of". As such, a history of high stressor exposure increased the  
4 likelihood of experiencing more frequent stressors, leading to the development of mental health  
5 disorders and reduced levels of well-being. This was further supported by Sophie: "It is that kind  
6 of helplessness feeling... I have never felt like I could fix it or make it better, [experiencing stress]  
7 just wears you down and you just cannot get away from it". Consequently, the accumulation of  
8 stress and adversity was found to promote maladaptive stress responses (e.g., negative emotions  
9 such anxiety or worry), which was further supported by Sophie: "I was a nightmare. I worried  
10 about everything. I was very, very insecure... Because so many other bad things had happened, I  
11 always thought the worst all the time. And even now, I catastrophize everything". This notion was  
12 exemplified by Sarah: "What happened in childhood made me more susceptible to the stressors I  
13 experienced as an adult... It made me just shut down and I couldn't cope". Thus, this theme  
14 revealed that exposure to greater lifetime stressors led to an increased susceptibility to future stress,  
15 via the promotion of maladaptive stress responses, likely explaining the development of negative  
16 health-related outcomes (e.g., depression).

### 17 *Interpersonal Relationship Difficulties*

18 Finally, four participants demonstrated difficulty in establishing trusting and intimate  
19 relationships with others as a result of experiencing prior stress and adversity. This was  
20 demonstrated by Jessica, who said that: "When I moved up to seniors at 16, there was a lot of  
21 people who were making advances on me and being inappropriate... I ended up being in an abusive  
22 marriage and nearly losing my life... It totally warped my relationships". Furthermore, the  
23 detrimental consequences associated with experiencing a high degree of lifetime stress without

1 adequate support was illustrated by Mary: “I felt abandoned, I had to fend for myself... I shy away  
2 from intimate relationships and meaningful friendships... it has dramatically affected my life and  
3 my ability to be a mother”. Fragmented relationships with others resulted in participants reporting  
4 lower social support and increased distance from others. This was exemplified by Sophie, who  
5 noted that: “It is probably a feeling of insecurity and you just think, in particular with my husband,  
6 well he won’t stay because people don’t”. This notion was also illustrated by Elizabeth, who stated  
7 that: “I think the stressors that I have experienced made it very difficult for me to interact with  
8 other people... I can thank running for a lot of things, but I can’t cope in a normal environment”.  
9 Consequently, limited interpersonal relationships with others, coupled with the presence of major  
10 stressors and adversities over the life course, affected participants mental health and well-being.  
11 For example, Elizabeth stated: “My social circle was incredibly small, I had no friends outside of  
12 sport, I had no support structure... I felt abandoned... It resulted in a long-term anxiety problem  
13 and has took a long time to even start to recover from that”. Therefore, this theme revealed that  
14 the difficulties associated with establishing interpersonal relationships with others partly explained  
15 the link between cumulative lifetime stress exposure, mental health, and well-being.

## 16 **Discussion**

17 Despite considerable interest in stress, relatively little is known about how cumulative  
18 lifetime stress is associated with mental health and well-being in an elite sporting context. This  
19 study aimed to address this issue by collecting both quantitative and qualitative data from elite  
20 athletes and in doing so, provide the first insight into how lifetime stress exposure relates to mental  
21 health and well-being. The results revealed significant associations between cumulative lifetime  
22 stress exposure and symptoms of depression and anxiety, as well as well-being. These associations  
23 were similar for both lifetime stressor count and severity, and robust when controlling for age and

1 gender, which are well-known predictors of poor mental health (Harkness et al., 2010). These  
2 findings are also consistent with the integrative model of lifespan stress and health (Epel et al.,  
3 2018), and suggest that greater lifetime stress is associated with poorer mental health and well-  
4 being but that these effects differ by the specific types of stressors experienced.

5         Associations between different stressor types and athletes' mental health status revealed  
6 that the total count and severity of chronic difficulties was a marginally stronger predictor of  
7 depression, when compared with the total count and severity of acute life events. Contextualized  
8 within extant literature and theory, these findings are congruent with the integrative model (Epel  
9 et al., 2018), suggesting that chronic stress may be a stronger predictor of depression than acute  
10 stress (Slavich et al., 2019). In contrast, when comparing these associations with symptoms of  
11 anxiety and well-being, results for the total count and severity of chronic difficulties were nearly  
12 identical to those observed for the count and severity of acute life events. An interesting lack of  
13 association, however, was observed between the total severity of chronic difficulties and symptoms  
14 of anxiety. This is in contrast to prior research, which has demonstrated that chronic difficulties  
15 were a marginally stronger predictor of psychiatric disorders (i.e., depression and anxiety) than  
16 acute life events (e.g., Slavich et al., 2019). One potential explanation for this finding could involve  
17 the low levels of anxiety reported by participants in the present study. Alternatively, chronic stress  
18 may act as a motivator for success, rather than as an anxiety-producing barrier, in elite athletes.

19         The present study revealed significant relationships between recent adulthood life events  
20 and symptoms of depression and anxiety, as well as well-being. Congruent with prior research  
21 (e.g., Lam et al., 2019), associations between lifetime stress exposure and health-related outcomes  
22 were consistent and strongest for stressors occurring in adulthood. This finding suggests that  
23 experiencing greater and more severe recent life events are predictive of mental ill-health and

1 lower levels of well-being among elite athletes. In contrast, exposure to early adversity was found  
2 to be predictive of symptoms of depression and anxiety; however, these associations were not as  
3 consistent or as strongly related to mental health problems and levels of well-being. This finding  
4 was interesting as the association between childhood adversity and poor mental health is nearly  
5 ubiquitous in the psychology literature (e.g., Hughes et al., 2017). One possible explanation for  
6 this finding may be that exposure to early adversity may interact and exacerbate the effects of  
7 recent stressors, leading to the development of mental ill-health among elite athletes (Turney, 2014).

8 Consistent with the stressor characteristic perspective (e.g., Slavich et al., 2019), stressors  
9 categorized by primary life domains and social-psychological characteristics were found to be  
10 differentially associated with symptoms of depression and anxiety, as well as well-being. Most  
11 notably, the stressor count indices that most consistently and significantly predicted mental health  
12 and well-being were role change (e.g., starting a new job) and other relationships (e.g., parental or  
13 non-intimate relationship problems). In contrast, the stressor severity indices that most consistently  
14 and significantly predicted mental health and well-being were interpersonal loss (e.g., close friend  
15 moves away), physical danger (e.g., being robbed at gun point), entrapment (e.g., having to care  
16 for a disabled parent), humiliation (e.g., being cheated on by a romantic partner) and  
17 marital/partner (e.g., ongoing marital discord). A growing body of work suggests that major life  
18 stressors that include interpersonal loss and humiliation are particularly harmful for health (e.g.,  
19 Kendler et al., 2003; Slavich, 2020). As such, this study extends existing work by highlighting the  
20 specific life domains and social-psychological characteristics that might make athletes more  
21 susceptible to poor mental health and well-being.

22 This study also used qualitative methods to help explain why cumulative lifetime stress  
23 exposure might have resulted in poor mental health and well-being, which added meaning to the

1 quantitative results. First, participants revealed how adversity-related experiences led to improved  
2 sporting performance in the short-term, but, also to some maladaptive coping strategies in the long-  
3 term. For example, the qualitative data suggested that the exclusive focus on sport following  
4 adversity led to the development of a constrained identity. Although intuitively appealing for  
5 sporting performance, research has shown that athletic identity foreclosure can limit adaptive  
6 responses to stress, resulting in psychological difficulties (e.g., Giannone, Haney, Kealy, &  
7 Ogradniczuk, 2017). The data, therefore, offer some support to the potential antithetical  
8 relationship between well-being and performance among athletes at the highest level (Fletcher,  
9 2019). Second, consistent with prior research, the accumulation of stressors during childhood was  
10 found to render participants more susceptible to stress throughout adulthood (e.g., Pearlin,  
11 Schieman, Fazio, & Meersman, 2005). This finding suggests that the accumulation of stressors  
12 over the life course may limit coping resources to deal with the demands of a stressful situation.  
13 One theoretical framework that offers a possible explanation for this result is the biopsychosocial  
14 model of challenge and threat (Blascovich, 2008a), which suggests that when coping resources are  
15 limited, individuals will typically appraise a stressful situation as more of a threat. When frequently  
16 experienced, repeated threat appraisals have been linked to deleterious health consequences (e.g.,  
17 depression; Blascovich, 2008b). Finally, this study found that participants who experienced greater  
18 and more severe lifetime stressors demonstrated difficulty in establishing and maintaining  
19 interpersonal relationships. Congruent with the interpersonal theory of psychopathology, the  
20 presence of stressors can have detrimental effects on health through the disruption of interpersonal  
21 relationships (Hammen & Rudolph, 1996). Thus, limited interpersonal relationships with others  
22 can increase the cumulative effects of stress (McEwen & Gianaros, 2010).

23         Several strengths and weaknesses of this study should be noted. In terms of strengths, this

1 is the first mixed methods study to examine cumulative lifetime stress exposure and mental health  
2 and well-being in elite athletes. Indeed, the adoption of both quantitative and qualitative  
3 methodologies provided a more comprehensive understanding of phenomena. Furthermore, this  
4 study examined stress-health links in elite athletes, which is noteworthy considering the increasing  
5 risk of mental ill-health in this population (Gorczyński et al., 2017). Finally, this study assessed  
6 participants' exposure to different types of stressors over the entire lifespan, which has rarely been  
7 done (Slavich & Shields, 2018). Consequently, the results provide meaningful information to  
8 researchers and practitioners who are interested in better understanding the risk factors that are  
9 associated with the mental health and well-being of the athletes they work with.

10         With regard to limitations, first, the exclusive focus on non-sporting stressors and  
11 adversities (e.g., bereavement), and the absence of sport-related stressors and adversities (e.g.,  
12 injury), could be seen as a potential limitation. Indeed, it is possible that stressors unique to the  
13 elite sporting environment could have different effects on athletes' mental health (Rice et al., 2016).  
14 Therefore, future research should examine the role of both sporting and non-sporting stressors to  
15 garner a more comprehensive understanding of cumulative lifetime stress exposure on athlete  
16 mental health and well-being. Second, a cross-sectional study design was used, which prevents  
17 causal inferences from being drawn from the data. Future research should therefore assess lifetime  
18 stress and mental health using longitudinal designs. Third, although adequately powered, the study  
19 used a sample size that is relatively small owing to challenges associated with collecting data from  
20 elite athletes. Consequently, future research should attempt to gather data from larger samples.

21         From an applied perspective, the results of this study have important implications for  
22 practitioners and organizations that are committed to better supporting mental health and well-  
23 being. Based on these findings, it is suggested that practitioners working with sport performers

1 should be aware of the competing demands placed on elite athletes and begin assessing their stress  
2 levels (i.e., historical and current stress exposure) to help identify athletes who may be most  
3 susceptible to developing mental health issues (Fletcher et al., 2006). Notably, the quantitative  
4 findings emphasize the importance of identifying athletes who have been exposed to stressors that  
5 are either chronic in nature or have occurred in adulthood. As a result, the implementation of a  
6 stress audit within elite sport could provide practitioners with important information and enable  
7 coaches to address stressors that increases individuals' risk for mental health problems and  
8 diminished well-being (Reardon et al., 2019). With regard to the qualitative findings, coaches  
9 should promote effective coping by improving athletes' self-awareness using systematic self-  
10 reflection (Tamminen & Holt, 2012). In doing so, athletes could become better able to assess the  
11 demands of stressful situations and identify the resources that they have to effectively cope  
12 (Blascovich, 2008a). Furthermore, the ability to cope with stressful situations can be bolstered by  
13 developing strong interpersonal relationships with others. Specifically, sport psychologists should  
14 help athletes identify various sources of social support and facilitate the development of a  
15 supportive environment (e.g., improved communication and empathy with others in the sporting  
16 organization; Freeman, 2020). Finally, sporting organizations must support elite athletes by  
17 viewing mental health as an organizational priority (Reardon et al., 2019).

18 To conclude, this study examined cumulative lifetime stress exposure and mental health  
19 and well-being in elite athletes, using a mixed-method approach. The findings demonstrated that  
20 greater and more severe lifetime stress was associated with poorer mental health and well-being.  
21 Moreover, follow-up qualitative data suggested that maladaptive long-term coping strategies,  
22 increased susceptibility to future stress, and difficulties establishing interpersonal relationships,  
23 may help explain the link between lifetime stress and poorer mental health and well-being.

## References

- 1  
2 Arnold, R., Fletcher, D., & Daniels, K. (2016). Demographic differences in sport performers'  
3 experiences of organizational stressors. *Scandinavian Journal of Medicine and Science in*  
4 *Sports*, 26(3), 348-358. doi: 10.1111/sms.12439
- 5 Blascovich, J. (2008a). Challenge and threat. In A. J. Elliot (Ed.), *Handbook of approach and*  
6 *avoidance motivation* (pp. 431-445). New York, NY: Psychology Press.
- 7 Blascovich, J. (2008b). Challenge, threat, and health. In J. Y. Shah & W. L. Gardner (Eds.),  
8 *Handbook of motivation science* (pp. 481–493). New York, NY: The Guilford Press.
- 9 Braun, V., & Clarke, V. (2006). Using thematic analysis in psychology. *Qualitative Research in*  
10 *Psychology*, 3(2), 77-101. doi: 10.1191/1478088706qp063oa
- 11 Cazassa, M., Oliveira, M., Spahr, C., Shields, G., & Slavich, G. (2020). The Stress and Adversity  
12 Inventory for Adults (Adult STRAIN) in Brazilian Portuguese: Initial validation and links  
13 with executive function, sleep, and mental and physical health. *Frontiers in Psychology*, 10,  
14 30383. doi: 10.3389/fpsyg.2019.03083
- 15 Creswell, J., & Plano-Clark, V. (2011). *Designing and conducting mixed method research* (2nd  
16 ed.). Thousand Oaks, CA: Sage.
- 17 Epel, E. S., Crosswell, A. D., Mayer, S. E., Prather, A. A., Slavich, G. M., Puterman, E., &  
18 Mendes, W. B. (2018). More than a feeling: A unified view of stress measurement for  
19 population science. *Frontiers in Neuroendocrinology*, 49, 146-169. doi:  
20 10.1016/j.yfrne.2018.03.001
- 21 Fletcher, D. (2019). Psychosocial resilience and adversarial growth in sport and performance. In  
22 E. O. Acevedo (ed.), *The Oxford Encyclopedia of Sport, Exercise, and Performance*  
23 *Psychology* (pp. 731-756). New York, NY: Oxford University Press.

- 1 Fletcher, D., & Sarkar, M. (2012). A grounded theory of psychological resilience in Olympic  
2 champions. *Psychology of Sport and Exercise, 13*(5), 669-678. doi:  
3 10.1016/j.psychsport.2012.04.007
- 4 Fletcher, D., Hanton, S., & Mellalieu, S. D. (2006). An organizational stress review: Conceptual  
5 and theoretical issues in competitive sport. In S. Hanton and S. D Mellalieu (Eds.),  
6 *Literature reviews in sport psychology* (pp. 321-374). Hauppauge, NY: Nova Science  
7 Publishers.
- 8 Foskett, R., & Longstaff, F. (2018). The mental health of elite athletes in the United Kingdom.  
9 *Journal of Science and Medicine in Sport, 21*(8), 765-770. doi:  
10 10.1016/j.jsams.2017.11.016
- 11 Freeman, P. (2020). Social Support. In R. Arnold & D. Fletcher (Eds.), *Stress, well-being, and*  
12 *performance in sport*. Oxford, UK: Taylor & Francis.
- 13 Giannone, Z., Haney, C., Kealy, D., & Ogrodniczuk, J. (2017). Athletic identity and psychiatric  
14 symptoms following retirement from varsity sports. *International Journal of Social*  
15 *Psychiatry, 63*(7), 598-601. doi: 10.1177/0020764017724184
- 16 Gorczynski, P., Coyle, M., & Gibson, K. (2017). Depressive symptoms in high-performance  
17 athletes and non-athletes: A comparative meta-analysis. *British Journal of Sports*  
18 *Medicine, 51*(18), 1348-1354. doi: 10.1136/bjsports-2016-096455
- 19 Greene, J., Caracelli, V., & Graham, W. (1989). Toward a conceptual framework for mixed  
20 method evaluation designs. *Educational Evaluation and Policy Analysis, 11*(3), 255-274.  
21 doi: 10.3102/01623737011003255
- 22 Guest, G., MacQueen, K., & Namey, E. (2012). *Applied thematic analysis*. Los Angeles, LA: Sage  
23 Publications.

- 1 Hammen, C., & Rudolph, K. D. (1996). Childhood depression. In E. J. Mash & R. A. Barkley  
2 (Eds.), *Child psychopathology* (pp. 153-195). New York, NY: Guilford Press.
- 3 Hardy, L., Barlow, M., Evans, L., Rees, T., Woodman, T., & Warr, C. (2017). Great British  
4 medalists: Psychosocial biographies of super-elite and elite athletes from Olympic sports.  
5 In V. Walsh, M. Wilson, & B. Parkin (Eds.), *Sport and the brain: The science of preparing,  
6 enduring and winning Part A* (pp. 1-119). London, UK: Academic Press.
- 7 Harkness, K., Alavi, N., Monroe, S., Slavich, G., Gotlib, I., & Bagby, R. (2010). Gender  
8 differences in life events prior to onset of major depressive disorder: The moderating  
9 effect of age. *Journal of Abnormal Psychology, 119*(4), 791-803. doi: 10.1037/a0020629
- 10 Howells, K., & Fletcher, D. (2015). Sink or swim: Adversity- and growth-related experiences in  
11 Olympic swimming champions. *Psychology of Sport and Exercise, 16*, 37-48. doi:  
12 10.1016/j.psychsport.2014.08.004
- 13 Hughes, K., Bellis, M., Hardcastle, K., Sethi, D., Butchart, A., Mikton, C., ... Dunne, M. P.  
14 (2017). The effect of multiple adverse childhood experiences on health: A systematic  
15 review and meta-analysis. *The Lancet Public Health, 2*(8), 356-366. doi: 10.1016/s2468-  
16 2667(17)30118-4
- 17 Johnson, R., & Onwuegbuzie, A. (2004). Mixed methods research: A research paradigm  
18 whose time has come. *Educational Researcher, 33*(7), 14-26. doi:  
19 10.3102/0013189x033007014
- 20 Kendler, K., Hettema, J., Butera, F., Gardner, C., & Prescott, C. (2003). Life event dimensions  
21 of loss, humiliation, entrapment, and danger in the prediction of onsets of major  
22 depression and generalized anxiety. *Archives of General Psychiatry, 60*(8), 789-796.  
23 doi: 10.1001/archpsyc.60.8.789

- 1 Kroenke, K., Spitzer, R. L., & Williams, J. B. (2001). The PHQ-9: Validity of a brief depression  
2 severity measure. *Journal of General Internal Medicine*, *16*(9), 606-613. doi:  
3 10.1046/j.1525-1497.2001.016009606.x
- 4 Kvale, S., & Brinkmann, S. (Eds). (2009). *InterViews: Learning the craft of qualitative*  
5 *research interviewing*. Los Angeles, CA: Sage.
- 6 Lam, J. C. W., Shields, G. S., Trainor, B. C., Slavich, G. M., & Yonelinas, A. P. (2019). Greater  
7 lifetime stress exposure predicts blunted cortisol but heightened DHEA responses to acute  
8 stress. *Stress and Health*, *35*(1), 15-26. doi: 10.1002/smi.2835
- 9 Levitt, H., Motulsky, S., Wertz, F., Morrow, S., & Ponterotto, J. (2017). Recommendations  
10 for designing and reviewing qualitative research in psychology: Promoting  
11 methodological integrity. *Qualitative Psychology*, *4*(1), 2–22. doi: 10.1037/qup0000082
- 12 Longo, Y., Coyne, I., & Joseph, S. (2018). Development of the short version of the Scales of  
13 General Well-Being: The 14-item SGWB. *Personality and Individual Differences*, *124*,  
14 31-34. doi: 10.1016/j.paid.2017.11.042
- 15 Löwe, B., Decker, O., Müller, S., Brähler, E., Schellberg, D., Herzog, W., & Herzberg, P.  
16 (2008). Validation and standardization of the generalized anxiety disorder screener  
17 (GAD-7) in the general population. *Medical Care*, *46*(3), 266-274. doi:  
18 10.1097/mlr.0b013e318160d093
- 19 Malterud, K., Siersma, V., & Guassora, A. (2016). Sample size in qualitative interview studies.  
20 *Qualitative Health Research*, *26*(13), 1753-1760. doi: 10.1177/1049732315617444
- 21 McEwen, B. S., & Gianaros, P. (2010). Central role of the brain in stress and adaptation. *Annals*  
22 *of the New York Academy of Sciences*, *1186*(1), 190–222. doi: 10.1111/j.1749-  
23 6632.2009.05331.x.

- 1 Moesch, K., Kenttä, G., Kleinert, J., Quignon-Fleuret, C., Cecil, S., & Bertollo, M. (2018).  
2 FEPSAC position statement: Mental health disorders in elite athletes and models of  
3 service provision. *Psychology of Sport and Exercise*, 38, 61-71. doi:  
4 10.1016/j.psychsport.2018.05.013
- 5 Moore, L., Young, T., Freeman, P., & Sarkar, M. (2017). Adverse life events, cardiovascular  
6 responses, and sports performance under pressure. *Scandinavian Journal of Medicine &  
7 Science in Sports*, 28(1), 340-347. doi: 10.1111/sms.12928
- 8 Morrow, S. L. (2005). Quality and trustworthiness in qualitative research in counseling  
9 psychology. *Journal of Counseling Psychology*, 52(2), 250-260. doi: 10.1037/0022-  
10 0167.52.2.250
- 11 Neil, R., McFarlane, H. M., & Smith, A. P. (2017). Well-being in sport organizations. In C. R.  
12 D. Wagstaff (Ed.), *The organizational psychology of sport: Key issues and practical  
13 applications* (p. 101–119). New York, NY: Routledge.
- 14 Olvera Alvarez, H., Provencio-Vasquez, E., Slavich, G., Laurent, J., Browning, M., & McKee-  
15 Lopez, G. et al. (2019). Stress and health in nursing students. *Nursing Research*, 68(6),  
16 453-463. doi: 10.1097/nnr.0000000000000383
- 17 Onwuegbuzie, A., & Leech, N. (2005). Taking the “Q” out of research: Teaching research  
18 methodology courses without the divide between quantitative and qualitative  
19 paradigms. *Quality & Quantity*, 39(3), 267-295. doi: 10.1007/s11135-004-1670-0
- 20 Pearlín, L., Schieman, S., Fazio, E., & Meersman, S. (2005). Stress, health, and the life course:  
21 Some conceptual perspectives. *Journal of Health and Social Behavior*, 46(2), 205-219.  
22 doi: 10.1177/002214650504600206
- 23 Pegg, S., Ethridge, P., Shields, G. S., Slavich, G. M., Weinberg, A., & Kujawa, A. (2019). Blunted social

- 1 reward responsiveness moderates the effect of lifetime social stress exposure on depressive  
2 symptoms. *Frontiers in Behavioral Neuroscience*, *13*, 178. doi: 10.3389/fnbeh.2019.00178
- 3 Reardon, C., Hainline, B., Aron, C., Baron, D., Baum, A., Bindra, A., ... Engebretsen, L. (2019).  
4 Mental health in elite athletes: International Olympic Committee consensus statement  
5 (2019). *British Journal of Sports Medicine*, *53*(11), 667-699. doi: 10.1136/bjsports-2019-  
6 100715
- 7 Rice, S. M., Purcell, R., De Silva, S., Mawren, D., McGorry, P. D., & Parker, A. G. (2016). The  
8 mental health of elite athletes: A narrative systematic review. *Sports Medicine*, *46*(9), 1333-  
9 1353. doi: 10.1007/s40279-016-0492-2
- 10 Ryan, R. M., & Deci, E. L. (2001). On happiness and human potential: A review of research on  
11 hedonic and eudaimonic well-being. *Annual Review of Psychology*, *52*(1), 141-166. doi:  
12 10.1146/annurev.psych.52.1.141
- 13 Slavich, G. M. (2020). Social safety theory: A biologically based evolutionary perspective on life  
14 stress, health, and behavior. *Annual Review of Clinical Psychology*, *16*(1), 265-295. doi:  
15 10.1146/annurev-clinpsy-032816-045159
- 16 Slavich, G. M., & Shields, G. S. (2018). Assessing lifetime stress exposure using the Stress and  
17 Adversity Inventory for Adults (Adult STRAIN). *Psychosomatic Medicine*, *80*(1), 17-27.  
18 doi: 10.1097/psy.0000000000000534
- 19 Slavich, G. M., Stewart, J. G., Esposito, E. C., Shields, G. S., & Auerbach, R. P. (2019). The Stress  
20 and Adversity Inventory for Adolescents (Adolescent STRAIN): Associations with mental  
21 and physical health, risky behaviors, and psychiatric diagnoses in youth seeking treatment.  
22 *Journal of Child Psychology and Psychiatry*, *60*(9), 998-1009. doi: 10.1111/jcpp.13038
- 23 Spitzer, R., Kroenke, K., Williams, J., & Löwe, B. (2006). A brief measure for assessing

- 1           generalized anxiety disorder. *Archives of Internal Medicine*, *166*(10), 1092. doi:  
2           10.1001/archinte.166.10.1092
- 3 Sturmbauer, S., Shields, G., Hetzel, E., Rohleder, N., & Slavich, G. (2019). The Stress and  
4           Adversity Inventory for Adults (Adult STRAIN) in German: An overview and initial  
5           validation. *PLOS ONE*, *14*(5), e0216419. doi: 10.1371/journal.pone.0216419
- 6 Swann, C., Moran, A., & Piggott, D. (2015). Defining elite athletes: Issues in the study of expert  
7           performance in sport psychology. *Psychology of Sport and Exercise*, *16*, 3-14. doi:  
8           10.1016/j.psychsport.2014.07.004
- 9 Tamminen, K. A., & Holt, N. L. (2012). Adolescent athletes' learning about coping and the roles  
10          of parents and coaches. *Psychology of Sport & Exercise*, *13*(1), 69-79. doi:  
11          10.1016/j.psychsport.2011.07.006
- 12 Teddlie, C., & Tashakkori, A. (2011). Mixed methods research: contemporary issues in an  
13          emerging field. In N. Denzin, & Y. Lincoln (Eds.), *The SAGE handbook of qualitative*  
14          *research*. (pp. 285-299). London: Sage.
- 15 Toussaint, L., Shields, G. S., Dorn, G., & Slavich, G. M. (2016). Effects of lifetime stress  
16          exposure on mental and physical health in young adulthood: How stress degrades and  
17          forgiveness protects health. *Journal of Health Psychology*, *21*(6), 1004-1014.  
18          doi: 10.1177/1359105314544132
- 19 Turney, K. (2014). Stress proliferation across generations? Examining the relationship between  
20          parental incarceration and childhood health. *Journal of Health and Social Behavior*, *55*(3),  
21          302-319. doi: 10.1177/0022146514544173
- 22 World Health Organization. (2004). Promoting mental health: Concepts, emerging evidence,  
23          practice: Summary report. Geneva: World Health Organization.

## Tables

**Table 1.** Means, standard deviations, and intercorrelations for main study variables.

	Mean	<i>SD</i>	1	2	3
1. Depression	4.15	3.63	-		
2. Anxiety	3.49	2.30		-	
3. Well-being	53.67	9.58			-
4. Total count of lifetime stressors	12.86	10.81	.41***	.33**	-.41***
5. Total severity of lifetime stressors	31.48	27.70	.43***	.23	-.26*
6. Count of acute life events	7.76	7.28	.30*	.22	.28*
7. Count of chronic difficulties	5.11	4.66	.44***	.24*	-.32**
8. Count of early adversity	2.77	3.25	.30**	.29*	-.16
9. Count of adulthood adversity	9.71	8.80	.39***	.27*	-.36**
10. Severity of acute life events	16.12	13.71	.34**	.29*	-.24*
11. Severity of chronic difficulties	15.36	15.68	.43***	.21	-.30**
12. Severity of early adversity	7.80	9.13	.30*	.39***	-.20
13. Severity of adulthood adversity	23.68	22.41	.42***	.15	-.31**

\*  $p < .05$ ; \*\*  $p < .01$ ; \*\*\*  $p < .001$ .

**Table 2.** Hierarchical regression models examining if lifetime stress (LTS) count and severity predicted (a) symptoms of depression, (b) symptoms of anxiety, and (c) well-being (model 1), adjusting for age and gender (model 2).

(a)	<b>Depression</b>												
	Model 1			Model 2			Model 1			Model 2			
	B	SE B	$\beta$	B	SE B	$\beta$	B	SE B	$\beta$	B	SE B	$\beta$	
LTS Count	0.17 (0.08 to 0.27)	0.05	0.41***	0.18 (0.09 to 0.26)	0.04	0.42***	LTS Severity	0.71 (0.04 to 0.11)	0.02	.043***	0.08 (0.04 to 0.11)	0.02	0.46***
Age				-0.02 (-0.06 to 0.03)	0.02	-0.08	Age				-0.03 (-0.08 to 0.01)	0.02	-0.16
Gender				1.67 (0.62 to 2.76)	0.53	0.34**	Gender				1.42 (0.38 to 2.46)	0.52	0.30**
* $p < .05$ ; ** $p < .01$ , *** $p < .001$ , two-tailed													
(b)	<b>Anxiety</b>												
	Model 1			Model 2			Model 1			Model 2			
	B	SE B	$\beta$	B	SE B	$\beta$	B	SE B	$\beta$	B	SE B	$\beta$	
LTS Count	0.14 (0.04 to 0.23)	0.05	0.33**	0.14 (0.05 to 0.24)	0.05	0.34**	LTS Severity	0.04 (0.00 to 0.07)	0.02	.23	0.04 (0.01 to 0.81)	0.02	0.28*
Age				-0.03 (-0.08 to 0.02)	0.02	-0.16	Age				-0.04 (-0.09 to 0.05)	0.02	-0.22
Gender				1.10 (-0.04 to 2.21)	0.56	0.23	Gender				0.90 (-0.20 to 2.00)	0.55	0.20
* $p < .05$ ; ** $p < .01$ , *** $p < .001$ , two-tailed													
(c)	<b>Well-Being</b>												
	Model 1			Model 2			Model 1			Model 2			
	B	SE B	$\beta$	B	SE B	$\beta$	B	SE B	$\beta$	B	SE B	$\beta$	
LTS Count	-0.65 (-0.10 to -0.31)	0.17	-0.41***	-0.67 (-1.02 to -0.32)	0.18	-0.42***	LTS Severity	-0.16 (-0.30 to -0.02)	0.07	-.26*	-0.18 (-0.33 to -0.04)	0.74	-0.30*
Age				0.09 (-0.10 to 0.28)	0.10	0.11	Age				0.13 (-0.06 to 0.32)	0.10	0.17
Gender				0.55 (-3.96 to 5.05)	2.26	0.03	Gender				1.60 (-2.85 to 6.03)	2.23	0.09
* $p < .05$ ; ** $p < .01$ , *** $p < .001$ , two-tailed													

**Figure 1.** Associations between total count and severity of lifetime stress exposure and (a) depression, (b) anxiety, and (c) well-being, categorized by stressor timing, type, primary life domain, and core social-psychological characteristic. Error bars represent 95% confidence intervals.





