


# BMJ Open Prevalence of adverse cardiometabolic health markers in UK undergraduate university students: an observational cohort study

Matthew J Savage <sup>1,2,3</sup>, Eleanor L Procter,<sup>4</sup> Philip Hennis,<sup>4</sup> Alfie G Price,<sup>4</sup> Daniele Magistro,<sup>4</sup> Ruth M James<sup>2,3,4</sup>

**To cite:** Savage MJ, Procter EL, Hennis P, *et al.* Prevalence of adverse cardiometabolic health markers in UK undergraduate university students: an observational cohort study. *BMJ Open* 2025;**15**:e089771. doi:10.1136/bmjopen-2024-089771

► Prepublication history and additional supplemental material for this paper are available online. To view these files, please visit the journal online (<https://doi.org/10.1136/bmjopen-2024-089771>).

Received 10 June 2024

Accepted 07 March 2025



© Author(s) (or their employer(s)) 2025. Re-use permitted under CC BY-NC. No commercial re-use. See rights and permissions. Published by BMJ Group.

<sup>1</sup>SHAPE Research Group, School of Science and Technology, Nottingham Trent University, Nottingham, UK

<sup>2</sup>University of Leicester Diabetes Research Centre, Leicester, UK

<sup>3</sup>NIHR Leicester Biomedical Research Centre, Leicester, UK

<sup>4</sup>Nottingham Trent University, Nottingham, UK

## Correspondence to

Dr Ruth M James;  
rmj20@leicester.ac.uk

## ABSTRACT

**Objectives** The prevalence of adverse cardiometabolic health markers has increased substantially in UK young adults, and university students now make up a significant proportion of this population. Their health-related behaviours are poorer than age-matched normative data, and students' anthropometric outcomes deteriorate during their university career. The influence of university on cardiometabolic health markers is unclear, and men and students of Minoritised Ethnicity are often under-represented in student health research. This study aimed to determine the prevalence of adverse cardiometabolic health markers in undergraduate university students and assess differences between genders, ethnic groups and year of study.

**Design** Observational cohort study.

**Setting** A higher education institution in Nottingham, UK.

**Participants** Three independent cohorts of undergraduate university students (total n=1,299) completed five physiological tests and provided demographic information. One-way ANOVAs assessed differences between year of study and ethnic groups, and paired samples t-tests assessed differences between genders.

**Main outcome measures** Body mass index (BMI), waist circumference, waist to hip ratio (WHR), waist to height ratio (WHtR), blood pressure (BP) and glycated haemoglobin concentrations (HbA1c).

**Results** 34.5% had overweight or obesity, 7.6% had a 'very high' waist circumference, 11.0% had a high WHR, 25.5% had a high WHtR, 12.7% were classified as hypertensive and 3.0% had an HbA1c  $\geq 42$  mmol/mol, indicating impaired glucose regulation. Differences between year groups were present for diastolic BP and HbA1c ( $p < 0.01$ ). Gender and ethnic group differences ( $p < 0.05$ ) were present for all variables other than BMI (gender) and diastolic BP (gender and ethnic group).

**Conclusion** Overall, these data demonstrate the prevalence of adverse cardiometabolic health markers in UK undergraduate university students, highlighting differences between year groups, genders and ethnic groups. These findings should be considered when developing strategies to promote healthy lifestyles in higher education.

## INTRODUCTION

In 2021, the UK saw 2.9 million students enrolled in higher education.<sup>1</sup> University

## STRENGTHS AND LIMITATIONS OF THIS STUDY

- ⇒ The current study included an extensive, diverse sample which incorporated substantial numbers of underrepresented subpopulations in health-based research (ie, men and students of minoritised ethnicity).
- ⇒ Data were collected following the COVID-19 pandemic. University stakeholders can therefore use these results to inform health-based initiatives within a post-COVID-19 society.
- ⇒ A limitation of the study is the between-subjects design, meaning longitudinal changes cannot be inferred. Furthermore, biochemical measures were not included in the study but would be informative in enhancing the understanding of students' cardiometabolic health.

students currently constitute a considerable proportion of the nation's young adult demographic, with half of school leavers opting to pursue higher education.<sup>2</sup> Worryingly, substantial proportions of university students are reported to have problematic dietary habits,<sup>3 4</sup> be physically inactive,<sup>3</sup> engage in extensive periods of sedentary behaviour<sup>4 5</sup> and partake in high levels of binge drinking.<sup>4</sup> Ultimately, these poor behaviours can have negative consequences for cardiometabolic health, with UK students experiencing considerable increases in body mass ( $+>0.5$  kg),<sup>6</sup> waist circumference ( $+1.9$  cm)<sup>7</sup> and body fat percentage ( $+0.8\%$ )<sup>8</sup> through their university career. Furthermore, the prevalence of students (outside of the UK) presenting with at least one risk factor for cardiometabolic syndrome (MetS) ranges from 0.3% to 13%.<sup>9 10</sup> However, within the UK, higher education encompasses unique cultural, social and educational factors (ie, behavioural habits, style of tuition and opportunities for socialising)<sup>11–15</sup> that can contribute to the health status of university students. As such,

it is unclear whether previous findings are translatable to the UK student population.

Moreover, the context of student health has changed substantially in recent years with significant alterations to the delivery of education and typical daily living routines due to the COVID-19 pandemic.<sup>16 17</sup> Indeed, throughout this period, university students developed considerably poorer movement, nutrition and sleeping behaviours<sup>18–20</sup> leading to adverse trends in anthropometric outcomes.<sup>21</sup> Furthermore, there is evidence to suggest that markers of health and related behaviours remain poorer than prior to the pandemic following the removal of restrictions.<sup>22</sup> As such, it is reasonable to suggest that prepandemic data are no longer representative of the current student health landscape and that up-to-date baseline data are required to develop health promotion policy and practices, and longitudinal data collection is needed to assess the efficacy of these.

Additionally, the diversity of the UK student population has increased considerably in recent years.<sup>1</sup> From a health perspective, this may be important given that gender and ethnicity play a substantial role in the development of adverse cardiometabolic health markers in the general population.<sup>23 24</sup> Indeed, MetS is 0.5–1.5 times more prevalent in minoritised ethnic groups,<sup>25</sup> and men observe different anthropometric outcomes to that of women.<sup>6 26</sup> However, men and minoritised ethnicity students are often under-represented within these studies, meaning results cannot be generalised to the wider student population.

Ultimately, to inform the development of effective strategies to improve health in university students, studies must strive to include participant samples that are reflective of the current student demographic. Therefore, the aim of the present study was to identify the prevalence of adverse cardiometabolic health markers in UK university students and assess gender and ethnic disparities. It was hypothesised that students who are men and students of minoritised ethnicity would have poorer cardiometabolic health outcomes.

## METHODS

### Participants and setting

University students completed a battery of five physiological tests in the first 3 weeks of term one, in one of three academic years (2021–22 *n*=347; 2022–23 *n*=422; or 2023–24 *n*=530). Using opportunistic sampling in communal spaces on campus, 1299 undergraduate students at a single institution in the midlands of England completed the testing and formed the cohort analysed in this study. All participants provided informed consent prior to taking part in the study, and ethical approval was granted by the School of Science and Technology Human Ethics Committee of Nottingham Trent University (application ID: 19/20–76). The study is reported in line with the Strengthening the Reporting of Observational Studies in Epidemiology (STROBE) guidelines for observational studies.<sup>27</sup>

### Physiology testing battery

The battery of testing included measurements of height, body mass, waist and hip circumferences to enable the calculation of body mass index (BMI), waist to hip ratio (WHR) and waist to height ratio (WHtR). Height and body mass were measured without shoes or heavy clothing using a stadiometer (Seca, Hamburg, Germany) and a digital weighing scale (Seca, Hamburg, Germany). Waist and hip circumferences were assessed using a flexible measuring tape and in accordance with International Society for the Advancement of Kinanthropometry (ISAK) guidelines.<sup>28</sup> Systolic and diastolic BP were determined using commercially available arm and wrist BP cuffs (Omron, Japan). In 2021–22 and 2022–23, BP was assessed from the upper right arm, whereas in 2023–24, BP was measured from the right wrist in an attempt to improve participant experience. Within both methods, BP was determined in accordance with manufacturer guidelines. Agreement between arm and wrist BP has previously been shown to be good,<sup>29</sup> and data from our own laboratory showed a strong correlation (*r*=0.717) between the two methods for systolic BP in undergraduate students. Glycated haemoglobin concentration (HbA1c) was determined using a fingertip blood sample analysed using the Quo-Test HbA1c analyser (EKF Diagnostics, UK).

### Data analysis

Descriptive data are reported as mean±1 SD and percentages are presented to demonstrate the proportion of participants classified into ‘risk’ categories for BMI,<sup>30</sup> waist circumference,<sup>31</sup> WHR,<sup>31</sup> WHtR,<sup>32</sup> BP<sup>33</sup> and HbA1c<sup>34</sup> based on well-established global guidelines. To characterise students’ current physical health status, data were combined to form a single cross-sectional dataset. For the purposes of analysis in this study, participants were then categorised by year at university (first, second and third year), gender (men and women) and ethnicity (White, Asian, Black and other minoritised ethnicities). If participants did not state a year group, gender or ethnicity, they were included in overall population data but excluded from analysis between year, gender or ethnic groups.

A one-way Analysis of Variance (ANOVA) was conducted to assess differences between year of study as well as ethnic groups. This test was used despite variables not meeting the assumption of normal distribution, as currently there is not a widely accepted non-parametric alternative for sample sizes substantially larger than *n*=30.<sup>35 36</sup> Group variance was assessed through Levene’s test, and homogeneity among variables was determined when the significance value was greater than 0.05 (*p*>0.05). To test the assumption of sphericity, Mauchly’s test was employed, assuming sphericity when the test yielded a non-significant result (*p*>0.05). In cases of violation, the Greenhouse–Geisser correction was applied. Furthermore, for evaluating effect sizes, partial eta squared ( $\eta^2$ ) was used, with effect magnitudes categorised as follows: small (0.02–0.12), medium (0.13–0.25) and large (>0.26).<sup>37</sup> Post hoc tests were performed using the Bonferroni correction to

**Table 1** Participant information data presented as n (%)

	Total (n=1299)	2021–22 (n=347)	2022–23 (n=422)	2023–24 (n=530)
Age (years)				
18–21	836 (64.4)	230 (66.3)	264 (62.3)	342 (64.5)
22–25	297 (22.9)	95 (27.4)	97 (23.0)	105 (19.8)
26–35	120 (9.2)	18 (5.2)	43 (10.2)	59 (11.1)
35+	46 (3.5)	4 (1.2)	18 (4.3)	24 (4.5)
Gender				
Women	425 (32.7)	26 (7.5)	199 (47.2)	200 (37.7)
Men	589 (45.3)	40 (11.5)	222 (52.6)	327 (61.7)
Not specified	285 (21.9)	281 (81.0)	1 (0.2)	3 (0.6)
Ethnicity				
White	497 (38.2)	46 (13.3)	214 (50.7)	237 (44.7)
Asian	233 (17.9)	8 (2.3)	90 (21.3)	135 (25.5)
Black	202 (15.6)	9 (2.6)	82 (19.4)	111 (20.9)
Other	77 (5.9)	3 (0.9)	33 (7.8)	41 (7.7)
Not specified	290 (22.3)	281 (81.0)	3 (0.7)	6 (1.1)
Year of study				
First Year	623 (48.0)	122 (35.2)	216 (51.2)	285 (53.8)
Second Year	357 (27.5)	146 (42.1)	87 (20.6)	124 (23.4)
Third Year	319 (24.6)	79 (22.8)	119 (28.2)	121 (22.8)

determine significant differences between year of study and ethnic groups ( $p < 0.05$ ). Gender differences were examined using an independent samples t-test. Cohen's  $d$  ( $d$ ) was used to gauge effect sizes, categorised as trivial ( $< 0.2$ ), small ( $\geq 0.2$ ), medium ( $\geq 0.5$ ) or large ( $\geq 0.8$ ) effects.<sup>38</sup> Significance was established at  $p < 0.05$ , and the analysis was conducted using IBM SPSS Statistics (SPSS V.28.0, Chicago, IL).

### Patient and public involvement

The study did not include the involvement of participants or the general public in its design. Nonetheless, all participants were informed about the research objectives, and their informed consent was obtained. Participants completed the testing voluntarily, and their input was not solicited for interpreting or writing the results.

## RESULTS

The sociodemographic characteristics of the 1299 participants included in the study are presented in [table 1](#).


### Anthropometric measures

The mean pooled data for body mass, BMI, waist circumference, WHR and WHtR are shown in [table 2](#). The prevalence of students at 'high risk' for adverse cardiometabolic health markers is displayed in online supplemental table 1. Having a BMI  $\geq 25.0$  was the most common risk factor for poor cardiometabolic health, followed by having a high WHtR ( $\geq 0.5$ ).

When separated by gender, the prevalence of overweight and obesity was highest in men; however, the prevalence of very high waist circumference, WHR and WHtR was highest in women (online supplemental table 1). When separated by ethnicity, the prevalence of overweight and obesity and very high waist circumference was highest in Black students, whereas the prevalence of high WHR and WHtR was highest in Asian students (online supplemental table 1). Across the year of study, the prevalence of overweight and obesity, and very high waist

**Table 2** Pooled cross-sectional data for all variables

	Mean $\pm$ SD
Anthropometry	
Height (m) (n=1289)	171.3 $\pm$ 12.6
Body mass (kg) (n=1284)	72.0 $\pm$ 14.6
Body mass index (kg/m <sup>2</sup> ) (n=1282)	24.3 $\pm$ 4.5
Waist circumference (cm) (n=1286)	80.2 $\pm$ 10.8
WHR (n=1281)	0.8 $\pm$ 0.1
WHtR (n=1284)	0.5 $\pm$ 0.1
Blood pressure	
Systolic (mm Hg) (n=1290)	120.0 $\pm$ 13.9
Diastolic (mm Hg) (n=1290)	74.8 $\pm$ 10.1
HbA1c (mmol/mol) (n=926)	34.0 $\pm$ 4.0
HbA1c, glycated haemoglobin concentrations; WHR, waist to hip ratio; WHtR, waist to height ratio.	



**Figure 1** Displays the differences between gender for the reported variables. \*\*\* $p < 0.001$ ; \*\* $p < 0.01$  compared with women. BMI, body mass index; BP, blood pressure; HbA1c, glycated haemoglobin concentrations; WHR, waist to hip ratio; WHtR, waist to height ratio.

circumference was highest in third year students, while the prevalence of high WHR and high WHtR was highest in first year students (online supplemental table 1).

A main effect of gender was observed for waist circumference and WHR. Overall, men had a greater waist circumference ( $t = -9.10$ ;  $p < 0.001$ ) and a larger WHR than women ( $t = -10.59$ ;  $p < 0.001$ ) with medium effect sizes ( $d = 0.6$  and  $d = 0.7$ , respectively) (figure 1B,C). No gender differences were observed for BMI or WHtR ( $p > 0.05$ ) (figure 1A,D).

A main effect of ethnicity was observed for BMI ( $p < 0.001$ ), waist circumference ( $p = 0.01$ ), WHR ( $p < 0.001$ ) and WHtR ( $p < 0.001$ ) with small effect sizes ( $\eta^2 = 0.02$ ,  $\eta^2 = 0.01$ ,  $\eta^2 = 0.02$  and  $\eta^2 = 0.02$  respectively). Overall, Black students and students of other minoritised ethnicities had a higher BMI than White students ( $p < 0.05$ ) and Asian students ( $p < 0.05$ ) (figure 2A). Black students also had a higher waist circumference than Asian students ( $p = 0.02$ ) (figure 2B). White students had a lower WHR than Asian students ( $p = 0.04$ ) and students of other minoritised ethnicities ( $p < 0.01$ ) (figure 2C). White students also had a lower WHtR than all other ethnic groups ( $p < 0.05$ ) (figure 2D).

No main effect of year of study was observed for any anthropometric variable ( $p > 0.05$ ) (figure 3A–D).

### Blood pressure

The mean systolic and diastolic BP in 1290 students is shown in table 2, and of these, 12.7% were classified as


being hypertensive (online supplemental table 1). The prevalence of hypertension was higher in men compared with women (online supplemental table 1). When separated by ethnicity, the prevalence of hypertension was highest in White students (online supplemental table 1). Across year groups, the prevalence of hypertension was highest in third year students (online supplemental table 1).

Men had higher systolic BP than women ( $t = -10.52$ ;  $p < 0.001$ ) with a medium effect size ( $d = -0.7$ ), but no difference between genders was found in diastolic BP ( $p = 0.47$ ) (figure 1E,F). Additionally, a main effect of ethnicity was observed for systolic BP ( $p < 0.001$ ) with a small effect size ( $\eta^2 = 0.05$ ). Asian students had lower systolic BP than all other ethnic groups ( $p < 0.01$ ) and Black students had lower systolic BP than White students ( $p < 0.001$ ) (figure 2E). No main effect of ethnicity was found for systolic blood pressure ( $p = 0.06$ ) (figure 2F). A main effect of year of study was found for diastolic blood pressure ( $p = 0.03$ ) with small effect size ( $\eta^2 = 0.01$ ). Post hoc tests revealed third year students had higher diastolic blood pressure than second year students ( $p = 0.01$ ) (figure 3F). No main effect of year of study was found for systolic blood pressure ( $p > 0.05$ ) (figure 3E).

### HbA1c

The mean HbA1c in 926 students was  $34.0 \pm 4.0$  mmol/mol (table 2). Of these, 3.0% were classified as 'high risk' for developing type 2 diabetes (online supplemental table





**Figure 2** Displays the differences between ethnic groups for the reported variables <sup>aaa</sup> $p < 0.001$  compared with White and Asian students; <sup>aa</sup> $p < 0.01$  compared with White and Asian students; <sup>b</sup> $p < 0.05$  compared with Asian students; <sup>ccc</sup> $p < 0.001$  compared with White students; <sup>c</sup> $p < 0.05$  compared with White students; <sup>dd</sup> $p < 0.01$  compared with White students; <sup>d</sup> $p < 0.05$  compared with White students; <sup>eee</sup> $p < 0.001$  compared with Asian students; <sup>ee</sup> $p < 0.01$  compared with Asian students; <sup>eee</sup> $p < 0.001$  compared with Black students; <sup>ggg</sup> $p < 0.001$  compared with White students. BMI, body mass index; BP, blood pressure; HbA1c, glycated haemoglobin concentrations; WHR, waist to hip ratio; WHtR, waist to height ratio.

1). The prevalence of poor glycaemic control was similar between men and women (online supplemental table 1) and highest in Black students (online supplemental table 1). When separated by year of study, the prevalence of impaired glucose control was highest in first-year students (online supplemental table 1).

Men had a higher HbA1c compared with women ( $t = -3.14$ ;  $p < 0.001$ ) with a small effect size ( $d = -0.23$ ) (figure 1G). Furthermore, a main effect of ethnicity was observed for HbA1c ( $p < 0.001$ ) with a small effect size ( $\eta^2 = 0.06$ ). White students had lower HbA1c compared with Asian ( $p < 0.001$ ) and Black students ( $p < 0.001$ ) (figure 2G). A main effect of year of study was observed with a small effect size ( $p < 0.001$ ;  $\eta^2 = 0.02$ ), specifically first year students had a higher HbA1c than second year students ( $p = 0.001$ ) and third year students ( $p = 0.02$ ) (figure 3G).

### Multiple risk factors

The prevalence of multiple adverse cardiometabolic health markers is presented in online supplemental table 1. A greater proportion of men had one or two adverse cardiometabolic health markers compared with women


(online supplemental table 1). Additionally, the prevalence of multiple adverse cardiometabolic health markers was similar between ethnicities and year groups (online supplemental table 1).

### DISCUSSION

The results of the present study provide comprehensive up-to-date baseline data surrounding the prevalence of adverse cardiometabolic health markers in UK university students. Additionally, the findings demonstrate that year of study, gender and ethnicity all play a substantial role in determining markers.

### Comparisons to UK population data

In general, average health data collected for all markers in the current study falls within healthy ranges.<sup>30–32 34 39</sup> Encouragingly, the prevalence of overweight or obesity and very high waist circumference among students was somewhat lower than UK age-matched normative data (34.5% vs 37.0%–7.6% vs 16.0%, respectively).<sup>40</sup> However, the prevalence of hypertension (9.0%) was slightly higher in students than that of UK young adults,<sup>40</sup>



**Figure 3** Displays the differences between year groups of study for the reported variables <sup>ff</sup> $p < 0.01$  compared with second year students and <sup>gg</sup> $p < 0.01$  compared with second year and third year students. BMI, body mass index; BP, blood pressure; HbA1c, glycated haemoglobin concentrations; WHR, waist to hip ratio; WHtR, waist to height ratio.

as was the prevalence of impaired glucose regulation (1.59–1.77%).<sup>41</sup> Comparisons cannot be made for WHR and WHtR due to a lack of population level data. Overall, these data suggest that university life does not substantially impact the health of young adults in comparison to non-students. While this may be interpreted as a positive finding, it should be noted that university campuses are a relatively controllable environment within which key stakeholders are able to provide access to facilities, services and information in a single setting to promote health.<sup>42</sup> Therefore, the comparability of poor health markers to that of the wider UK general population may be considered as disappointing.<sup>40 41</sup> Additionally, higher education institutions should be cognisant of the substantial negative trends in health-related behaviours of students in recent years,<sup>18–20</sup> meaning that it is likely the prevalence of poor cardiometabolic health markers will increase. It is therefore critical that university stakeholders use these data as a baseline to aid in the development of effective surveillance systems and, where appropriate, effective intervention strategies to ensure that optimising student health remains a priority for successive generations.

### Gender

In line with previous literature, the current study demonstrates that, on average, men had higher values

for waist circumference, WHR, systolic BP and [HbA1c] compared with women.<sup>43–46</sup> Additionally, the prevalence of overweight and obesity and hypertension was also greater in men than women, whereas the prevalence of very high waist circumference, WHR and WHtR was higher among women than men. These differences may be explained by genetic factors including differences in the distribution of fat and muscle mass between men and women as well as differences in hormonal concentrations (ie, upregulation of testosterone and downregulation of oestrogen in men compared with women) that mean men are at greater risk of developing adverse markers of body composition and health (eg, BP and HbA1c).<sup>47 48</sup> However, previous literature in students has also demonstrated behavioural differences between men and women that could impact these health-related variables. Specifically, men have been shown to engage in poorer eating behaviours<sup>4</sup> and greater levels of alcohol drinking in comparison to women,<sup>4 49</sup> who develop poorer movement<sup>3 4 50</sup> and sleep habits,<sup>4 51</sup> as well as having poorer indicators of mental health.<sup>4 52</sup> These data add to existing literature by further demonstrating the existence of disparities in health markers between genders in university students. Higher education institutions should therefore adopt a gender-centred approach

when developing future initiatives to improve students' health.

### Ethnicity

Data from the current study area are consistent with previous literature, indicating that minoritised ethnic groups experience a greater prevalence of adverse health markers in comparison to White Europeans.<sup>24</sup> Specifically, these findings highlight that Black and Asian students are at the greatest risk of developing poor anthropometric outcomes and impaired glycaemic control. This could be because they experience greater barriers to conducting physical activity,<sup>53</sup> develop poorer nutrition behaviours<sup>54</sup> and have poorer sleep habits<sup>55</sup> compared with their White counterparts. In contrast, adverse BP findings were more prevalent in White students compared with minoritised ethnic students, providing conflicting evidence to previous literature.<sup>56</sup> Although behavioural factors are important, a plethora of genetic, socioeconomic, religious, cultural, environmental and social differences also likely play a role in determining these health markers. As such, identifying the complex system within which these factors interact is far beyond the scope of the current study. Nonetheless, the present study provides novel information surrounding nuanced variations between students from different ethnic backgrounds.

### Year of study

Findings from the current study show a main effect for year of study on body mass, with an overall trend towards higher body mass from first year to third year. Additionally, third-year students had higher diastolic BP than second-year students. These findings are supported by data indicating that the prevalence of overweight and obesity, very high waist circumference and hypertension were highest among third-year students. Taken together, these findings align with previous literature indicating that students observe substantial weight gain throughout their university career<sup>6 57</sup> as well as increased BP during periods of critical assessment.<sup>58</sup> Previously, this has been attributed to the development of poor behavioural habits and psychological markers while at university.<sup>57 59</sup> However, no effect of year of study was observed for any other anthropometric outcome in the current data. Hence, this may be attributed to the typical age of university enrolment often coinciding with the latter phases of adolescence during which physical maturation may still be occurring due to neurological development.<sup>60</sup> Alternatively, this could be owing to positive health behaviours leading to increased skeletal muscle mass.<sup>61</sup>

Mean [HbA1C] was higher in first year students compared with second and third year students, and the prevalence of impaired glycaemic control, high WHR and high WHtR was greatest in first year students. These data may be considered surprising given that evidence suggests behavioural outcomes become poorer during the transition to, and across the first year of university.<sup>7 62</sup> However, the context of health behaviours in early adulthood

has shifted dramatically in recent years with technological advancements in conjunction with the COVID-19 pandemic promoting poorer behaviours such as reduced physical activity, increased screen-time, increased snacking and emotional eating that have all become habitual.<sup>20 63–65</sup>

It is therefore possible that attending university provided students with an opportunity to engage in a positive lifestyle shift which incorporated more social opportunities to engage in healthful movement behaviours and less reliance on food as a means of combating boredom or improving emotional states. Additionally, despite a plethora of literature demonstrating that health status and behaviours in youth influence health later in life,<sup>66</sup> it is plausible that the window during which these poor behaviours are developed while at university (~3 years) is not chronic enough to begin negatively influencing physiological markers of health in young people. However, there is currently not enough robust longitudinal data to quantify whether this is an accurate assumption. Regardless of the reason, the findings of the current study provide vital insight into how markers of cardiometabolic health may be altered across the timeline of an undergraduate degree. However, there is a distinct need for need for further longitudinal studies that incorporate substantial sample sizes in order to adequately assess the impact of university life on health and behaviours. This will ultimately provide vital insight into health and lifestyle factors that require targeted initiatives to aid higher education institutions in producing healthy and economically active alumni that will positively contribute to the wider UK society.

### Strengths and limitations

A major strength of the current study is the extensive, diverse sample of UK undergraduate students. This included large numbers of men and students of minoritised ethnicities who have previously been underrepresented in the research area.<sup>6 50</sup> As such, these findings provide vital, novel insight into the health status of different gender and ethnic groups of students. Furthermore, data were collected during the years following the COVID-19 pandemic. These results can therefore be used as a current baseline by key stakeholders to inform future decisions surrounding the development of initiatives to improve the health of university students and track changes in a post-COVID-19 society.

The between-subjects nature of the study design means that longitudinal changes cannot be inferred, but the large sample size within each year of study provides much-needed insight into potential differences across the timeline of a typical undergraduate degree. Nevertheless, future longitudinal studies with large student numbers should be developed to gain an in-depth understanding of trajectories in students' cardiometabolic health throughout an undergraduate degree programme. While the current study has excellent representation within students of minoritised ethnicity, the more complex distinction between ethnicity and culture, including



potential differences for international students, was not within the scope of this study and should be investigated in the future. Finally, the current study did not include biochemical markers (ie, HDL-cholesterol, LDL-cholesterol, total cholesterol and serum triglyceride) that have been shown to be important in the screening of poor metabolic health in university students.<sup>10</sup> However, the processes required to collect such data would require participants to be fasted, which could substantially reduce the study sample size.

## Conclusion

The current study provides up-to-date data surrounding markers of cardiometabolic health in university students. On average, data falls within healthy ranges and the prevalence of adverse outcomes is largely similar to that of age-matched young adults in the UK. While encouraging, the relatively controllable environment in which many undergraduate students reside provides university stakeholders with opportunities to optimise health in their student populations. The current data demonstrate that gender and ethnicity play a key role in determining markers of cardiometabolic health, and therefore these factors must be considered when developing and implementing interventions. Furthermore, researchers in student health should work together to collect representative data within this population, to further elucidate differences between student populations, and to develop, test and implement successful health interventions for students.

**Contributors** MJS, RMJ and PH were responsible for study conceptualisation, data collection, data analysis and drafting of the manuscript. ELP and AGP contributed to data collection and drafting of the manuscript. DM contributed to drafting of the manuscript. All authors have read and agreed to the published version of the manuscript. MJS acted as the guarantor.

**Funding** The authors have not declared a specific grant for this research from any funding agency in the public, commercial or not-for-profit sectors.

**Competing interests** None declared.

**Patient and public involvement** Patients and/or the public were not involved in the design, conduct, reporting or dissemination plans of this research.

**Patient consent for publication** Not applicable.

**Ethics approval** This study involves human participants and was granted by the School of Science and Technology Human Ethics Committee of Nottingham Trent University (application ID: 19/20-76). Participants gave informed consent to participate in the study before taking part.

**Provenance and peer review** Not commissioned; externally peer reviewed.

**Data availability statement** Data are available upon reasonable request. The datasets used and/or analysed during the current study are available from the corresponding author on reasonable request.

**Supplemental material** This content has been supplied by the author(s). It has not been vetted by BMJ Publishing Group Limited (BMJ) and may not have been peer-reviewed. Any opinions or recommendations discussed are solely those of the author(s) and are not endorsed by BMJ. BMJ disclaims all liability and responsibility arising from any reliance placed on the content. Where the content includes any translated material, BMJ does not warrant the accuracy and reliability of the translations (including but not limited to local regulations, clinical guidelines, terminology, drug names and drug dosages), and is not responsible for any error and/or omissions arising from translation and adaptation or otherwise.

**Open access** This is an open access article distributed in accordance with the Creative Commons Attribution Non Commercial (CC BY-NC 4.0) license, which permits others to distribute, remix, adapt, build upon this work non-commercially,

and license their derivative works on different terms, provided the original work is properly cited, appropriate credit is given, any changes made indicated, and the use is non-commercial. See: <http://creativecommons.org/licenses/by-nc/4.0/>.

## ORCID iD

Matthew J Savage <http://orcid.org/0000-0003-2922-3681>

## REFERENCES

- 1 HESA. Who's studying in HE? | HESA, 2024. Available: <https://www.hesa.ac.uk/data-and-analysis/students/whos-in-he>
- 2 Bolton P. Higher education student numbers, 2024. Available: <https://commonslibrary.parliament.uk/research-briefings/cbp-7857/>
- 3 Aceijas C, Waldhäusl S, Lambert N, et al. Determinants of health-related lifestyles among university students. *Perspect Public Health* 2017;137:227–36.
- 4 Savage MJ, Procter EL, Hennis PJ, et al. Characterising the activity, lifestyle behaviours and health outcomes of UK university students: An observational cohort study. *BMC Public Health* 2024;24:3501.
- 5 Deliens T, Deforche B, De Bourdeaudhuij I, et al. Determinants of physical activity and sedentary behaviour in university students: a qualitative study using focus group discussions. *BMC Public Health* 2015;15:201.
- 6 Vadeboncoeur C, Foster C, Townsend N. Freshman 15 in England: a longitudinal evaluation of first year university student's weight change. *BMC Obes* 2016;3:45.
- 7 Gibson A-M, Shaw J, Hewitt A, et al. A longitudinal examination of students' health behaviours during their first year at university. *J Further Higher Educat* 2018;42:36–45.
- 8 Deliens T, Clarys P, Van Hecke L, et al. Changes in weight and body composition during the first semester at university. A prospective explanatory study. *Appetite* 2013;65:111–6.
- 9 Mattsson N, Rönnemaa T, Juonala M, et al. Childhood predictors of the metabolic syndrome in adulthood. The Cardiovascular Risk in Young Finns Study. *Ann Med* 2008;40:542–52.
- 10 Yahia N, Brown CA, Snyder E, et al. Prevalence of metabolic syndrome and its individual components among midwestern university students. *J Community Health* 2017;42:674–87.
- 11 Bennett G, O'Hara C, Bardón LA, et al. Comparison of meal patterns across common racial groups in the UK and the USA, examining associations with weight status and diet quality: a secondary analysis of NDNS and NHANES datasets. *J Racial and Ethnic Health Disparities* 2025;12:499–512.
- 12 Gallagher D, Gilmore A. Social integration and the role of student societies in higher education: an exploratory study in the UK. *J of Philanthropy and Mktg* 2013;18:275–86.
- 13 Hibell B, Andersson B, Bjarnason T, et al. Alcohol and other drug use among students in 35 European countries. 2003.
- 14 Holton M. Adapting relationships with place: Investigating the evolving place attachment and 'sense of place' of UK higher education students during a period of intense transition. *Geoforum* 2015;59:21–9.
- 15 Mai LW. A comparative study between UK and US: the student satisfaction in higher education and its influential factors. *Journal of Marketing Management* 2005;21:859–78.
- 16 Cao W, Fang Z, Hou G, et al. The psychological impact of the COVID-19 epidemic on college students in China. *Psychiatry Res* 2020;287:S0165-1781(20)30540-0.
- 17 Sahu P. Closure of Universities Due to Coronavirus Disease 2019 (COVID-19): impact on education and mental health of students and academic staff. *Cureus* 2020;12:e7541.
- 18 Foster JH, Rankin S. Self-reported sleep during the COVID lockdown in a sample of UK university students and staff. *Healthcare (Basel)* 2022;10:2083.
- 19 Gadi N, Saleh S, Johnson J-A, et al. The impact of the COVID-19 pandemic on the lifestyle and behaviours, mental health and education of students studying healthcare-related courses at a British university. *BMC Med Educ* 2022;22:115.
- 20 Savage MJ, Hennis PJ, Magistro D, et al. Nine months into the COVID-19 pandemic: a longitudinal study showing mental health and movement behaviours are impaired in UK students. *Int J Environ Res Public Health* 2021;18:2930.
- 21 Bell M, Duncan MJ, Patte KA, et al. Changes in body mass, physical activity, and dietary intake during the COVID-19 pandemic lockdowns in Canadian university students. *Biology (Basel)* 2023;12:326.
- 22 Salway R, Foster C, de Vocht F, et al. Accelerometer-measured physical activity and sedentary time among children and their



- parents in the UK before and after COVID-19 lockdowns: a natural experiment. *Int J Behav Nutr Phys Act* 2022;19:51.
- 23 Meloni A, Cadeddu C, Cugusi L, *et al.* Gender differences and cardiometabolic risk: the importance of the risk factors. *Int J Mol Sci* 2023;24:1588.
  - 24 Razieh C, Zaccardi F, Miksza J, *et al.* Differences in the risk of cardiovascular disease across ethnic groups: UK Biobank observational study. *Nutr Metab Cardiovasc Dis* 2022;32:2594–602.
  - 25 Ervin RB. Prevalence of metabolic syndrome among adults 20 years of age and over, by sex, age, race and ethnicity, and body mass index: United States, 2003. Available: <https://stacks.cdc.gov/view/cdc/5448>
  - 26 Deforche B, Van Dyck D, Deliens T, *et al.* Changes in weight, physical activity, sedentary behaviour and dietary intake during the transition to higher education: a prospective study. *Int J Behav Nutr Phys Act* 2015;12:16.
  - 27 Elm E von, Altman DG, Egger M, *et al.* Strengthening the reporting of observational studies in epidemiology (STROBE) statement: guidelines for reporting observational studies. *BMJ* 2007;335:806–8.
  - 28 Stewart A, Marfell-Jones M, Olds T, *et al.* International Standards for Anthropometric Assessment. 2011;137.
  - 29 Komori T, Eguchi K, Hoshida S, *et al.* Comparison of wrist-type and arm-type 24-h blood pressure monitoring devices for ambulatory use. *Blood Press Monit* 2013;18:57–62.
  - 30 NHS England. How to tell if you're living with obesity, 2017. Available: <https://www.nhs.uk/conditions/obesity/>
  - 31 WHO. Waist circumference and waist-hip ratio: report of a WHO expert consultation, Available: <https://www.who.int/publications-detail-redirect/9789241501491>
  - 32 Browning LM, Hsieh SD, Ashwell M. A systematic review of waist-to-height ratio as a screening tool for the prediction of cardiovascular disease and diabetes: 0.5 could be a suitable global boundary value. *Nutr Res Rev* 2010;23:247–69.
  - 33 Williams B, Poulter NR, Brown MJ, *et al.* British Hypertension Society guidelines for hypertension management 2004 (BHS-IV): summary. *BMJ* 2004;328:634–40.
  - 34 NICE. Overview | Type 2 diabetes: prevention in people at high risk | Guidance, Available: <https://www.nice.org.uk/guidance/ph38>
  - 35 Glass GV, Peckham PD, Sanders JR. Consequences of failure to meet assumptions underlying the fixed effects analyses of variance and covariance. *Rev Educ Res* 1972;42:237.
  - 36 Lumley T, Diehr P, Emerson S, *et al.* The importance of the normality assumption in large public health data sets. *Annu Rev Public Health* 2002;23:151–69.
  - 37 Bakeman R. Recommended effect size statistics for repeated measures designs. *Behav Res Methods* 2005;37:379–84.
  - 38 Cohen J. A power primer. *Psychol Bull* 1992;112:155–9.
  - 39 Unger T, Borghi C, Charchar F, *et al.* 2020 International Society of Hypertension Global Hypertension Practice Guidelines. *Hypertension* 2020;75:1334–57.
  - 40 NHS England. Health Survey for England 2019 [NS], Available: <https://digital.nhs.uk/data-and-information/publications/statistical/health-survey-for-england/2019>
  - 41 Campbell E, Macey E, Kneebone-Hopkins C, *et al.* Sociodemographic and health-related differences in non-diabetic hyperglycaemia and undiagnosed type 2 diabetes in the health survey for england 2013–2019. *Endocrinology (including Diabetes Mellitus and Metabolic Disease)* [Preprint] 2024. 10.1101/2024.02.12.24302519 Available: <https://www.medrxiv.org/content/10.1101/2024.02.12.24302519v1>
  - 42 Bailey CP, Sharma S, Economos CD, *et al.* College campuses' influence on student weight and related behaviours: A review of observational and intervention research. *Obes Sci Pract* 2020;6:694–707.
  - 43 Alhawari HH, Al-Shelleh S, Alhawari HH, *et al.* Blood pressure and its association with gender, body mass index, smoking, and family history among university students. *Int J Hypertens* 2018;2018:4186496.
  - 44 Aynaci G, Akdemir O. The relationship between lifestyle, health promotion lifestyle profile II and high blood pressure in university students. *Open Access Maced J Med Sci* 2018;6:1756–61.
  - 45 Gülsen Ş, Deniz KE, Başak C, *et al.* The effect of age and gender on HbA1c levels in adults without diabetes mellitus. *J Med Biochem* 2023;42:714–21.
  - 46 Peters SAE, Huxley RR, Woodward M. Sex differences in body anthropometry and composition in individuals with and without diabetes in the UK Biobank. *BMJ Open* 2016;6:e010007.
  - 47 Janssen I, Heymsfield SB, Wang ZM, *et al.* Skeletal muscle mass and distribution in 468 men and women aged 18–88 yr. *J Appl Physiol* (1985) 2000;89:81–8.
  - 48 Reckelhoff JF. Gender differences in the regulation of blood pressure. *Hypertension* 2001;37:1199–208.
  - 49 LaBrie JW, Ehret PJ, Hummer JF, *et al.* Poor adjustment to college life mediates the relationship between drinking motives and alcohol consequences: a look at college adjustment, drinking motives, and drinking outcomes. *Addict Behav* 2012;37:379–86.
  - 50 Yahia N, Wang D, Rapley M, *et al.* Assessment of weight status, dietary habits and beliefs, physical activity, and nutritional knowledge among university students. *Perspect Public Health* 2016;136:231–44.
  - 51 Becker SP, Jarrett MA, Luebke AM, *et al.* Sleep in a large, multi-university sample of college students: sleep problem prevalence, sex differences, and mental health correlates. *Sleep Health* 2018;4:174–81.
  - 52 Heller AT, Berg SS, Prichard JR. University housing reinforces the negative relationship between interpersonal violence, psychological distress, and suicidality in undergraduates, particularly among gender diverse students. *J Am Coll Health* 2023;71:102–10.
  - 53 Leslie-Walker A, Mulvenna C, Bramble O. Exploring the motivations and challenges West African female students experience engaging in university sport. *Curriculum Studies in Health and Physical Education* 2023;14:190–206.
  - 54 Boek S, Bianco-Simeral S, Chan K, *et al.* Gender and race are significant determinants of students' food choices on a college campus. *J Nutr Educ Behav* 2012;44:372–8.
  - 55 Sa J, Choe S, Cho B-Y, *et al.* Relationship between sleep and obesity among U.S. and South Korean college students. *BMC Public Health* 2020;20:96.
  - 56 Hardy ST, Chen L, Cherrington AL, *et al.* Racial and ethnic differences in blood pressure among US adults, 1999–2018. *Hypertension* 2021;78:1730–41.
  - 57 Vella-Zarb RA, Elgar FJ. Predicting the 'freshman 15': Environmental and psychological predictors of weight gain in first-year university students. *Health Educ J* 2010;69:321–32.
  - 58 Fejes I, Ábrahám G, Légrády P. The effect of an exam period as a stress situation on baroreflex sensitivity among healthy university students. *Blood Press* 2020;29:175–81.
  - 59 Serlachius A, Hamer M, Wardle J. Stress and weight change in university students in the United Kingdom. *Physiol Behav* 2007;92:548–53.
  - 60 Arain M, Haque M, Johal L, *et al.* Maturation of the adolescent brain. *Neuropsychiatr Dis Treat* 2013;9:449–61.
  - 61 Kemmler W, von Stengel S, Kohl M, *et al.* Impact of exercise changes on body composition during the college years - a five year randomized controlled study. *BMC Public Health* 2015;16:50.
  - 62 Winpenny EM, Smith M, Penney T, *et al.* Changes in physical activity, diet, and body weight across the education and employment transitions of early adulthood: A systematic review and meta-analysis. *Obes Rev* 2020;21:e12962.
  - 63 Burnatowska E, Surma S, Olszaneczka-Glinianowicz M. Relationship between Mental Health and Emotional Eating during the COVID-19 Pandemic: A Systematic Review. *Nutrients* 2022;14:3989.
  - 64 Jehi T, Khan R, Halawani R, *et al.* Effect of COVID-19 outbreak on the diet, body weight and food security status of students of higher education: a systematic review. *Br J Nutr* 2023;129:1916–28.
  - 65 Technology OA, Bermingham R. Parliamentary office of science and technology; screen use and health in young people, 2020. Available: <https://post.parliament.uk/research-briefings/post-pn-0635>
  - 66 Shah RV, Murthy VL, Colangelo LA, *et al.* Association of fitness in young adulthood with survival and cardiovascular risk: The Coronary Artery Risk Development in Young Adults (CARDIA) study. *JAMA Intern Med* 2016;176:87–95.